
Loredana Cleris

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1810611/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Selective cytotoxicity of betulinic acid on tumor cell lines, but not on normal cells. Cancer Letters, 2002, 175, 17-25.	7.2	441
2	In vitro and In vivo Activity of SKI-606, a Novel Src-Abl Inhibitor, against Imatinib-Resistant Bcr-Abl+ Neoplastic Cells. Cancer Research, 2006, 66, 11314-11322.	0.9	352
3	In Vivo Eradication of Human BCR/ABL-Positive Leukemia Cells With an ABL Kinase Inhibitor. Journal of the National Cancer Institute, 1999, 91, 163-168.	6.3	341
4	Growth-inhibitory effect of STI571 on cells transformed by theCOL1A1/PDGFB rearrangement. International Journal of Cancer, 2001, 92, 354-360.	5.1	114
5	Induction of apoptosis by fenretinide (4HPR) in human ovarian carcinoma cells and its association with retinoic acid receptor expression. , 1996, 65, 491-497.		106
6	TIMP3 regulates migration, invasion and in vivo tumorigenicity of thyroid tumor cells. Oncogene, 2011, 30, 3011-3023.	5.9	78
7	IGFBP7: an oncosuppressor gene in thyroid carcinogenesis. Oncogene, 2010, 29, 3835-3844.	5.9	69
8	Inhibition of RET tyrosine kinase by SU5416. Journal of Molecular Endocrinology, 2006, 37, 199-212.	2.5	68
9	Metallothionein 1G acts as an oncosupressor in papillary thyroid carcinoma. Laboratory Investigation, 2008, 88, 474-481.	3.7	60
10	Age- and irradiation-associated loss of bone marrow hematopoietic function in mice is reversed by recombinant human growth hormone. Experimental Hematology, 2004, 32, 171-178.	0.4	48
11	BIM upregulation and ROS-dependent necroptosis mediate the antitumor effects of the HDACi Givinostat and Sorafenib in Hodgkin lymphoma cell line xenografts. Leukemia, 2014, 28, 1861-1871.	7.2	48
12	Colorectal Tumors Are Effectively Eradicated by Combined Inhibition of β-Catenin, KRAS, and the Oncogenic Transcription Factor ITF2. Cancer Research, 2010, 70, 7253-7263.	0.9	45
13	Primary cross-resistance to BRAFV600E-, MEK1/2- and PI3K/mTOR-specific inhibitors in BRAF-mutant melanoma cells counteracted by dual pathway blockade. Oncotarget, 2016, 7, 3947-3965.	1.8	45
14	Role of retinoic acid receptor overexpression in sensitivity to fenretinide and tumorigenicity of human ovarian carcinoma cells. , 1999, 81, 829-834.		42
15	S100A11 Overexpression Contributes to the Malignant Phenotype of Papillary Thyroid Carcinoma. Journal of Clinical Endocrinology and Metabolism, 2013, 98, E1591-E1600.	3.6	40
16	AMPK activators inhibit the proliferation of human melanomas bearing the activated MAPK pathway. Melanoma Research, 2012, 22, 341-350.	1.2	38
17	The Anti–Human Leukocyte Antigen-DR Monoclonal Antibody 1D09C3 Activates the Mitochondrial Cell Death Pathway and Exerts a Potent Antitumor Activity in Lymphoma-Bearing Nonobese Diabetic/Severe Combined Immunodeficient Mice. Cancer Research, 2006, 66, 1799-1808.	0.9	37
18	Sorafenib Inhibits Lymphoma Xenografts by Targeting MAPK/ERK and AKT Pathways in Tumor and Vascular Cells. PLoS ONE, 2013, 8, e61603.	2.5	34

LOREDANA CLERIS

#	Article	IF	CITATIONS
19	Decrease in drug accumulation and in tumour aggressiveness marker expression in a fenretinide-induced resistant ovarian tumour cell line. British Journal of Cancer, 2001, 84, 1528-1534.	6.4	33
20	Antitumor Activity of Human CD34+Cells Expressing Membrane-Bound Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand. Human Gene Therapy, 2006, 17, 1225-1240.	2.7	33
21	Human CD34+ cells engineered to express membrane-bound tumor necrosis factor–related apoptosis-inducing ligand target both tumor cells and tumor vasculature. Blood, 2010, 115, 2231-2240.	1.4	32
22	Bcl-XL down-regulation suppresses the tumorigenic potential of NPM/ALK in vitro and in vivo. Blood, 2004, 103, 2787-2794.	1.4	30
23	Valproic acid enhances bosutinib cytotoxicity in colon cancer cells. International Journal of Cancer, 2009, 124, 1990-1996.	5.1	29
24	Role of TFG sequences outside the coiled-coil domain in TRK-T3 oncogenic activation. Oncogene, 2003, 22, 807-818.	5.9	28
25	Perifosine and sorafenib combination induces mitochondrial cell death and antitumor effects in NOD/SCID mice with Hodgkin lymphoma cell line xenografts. Leukemia, 2013, 27, 1677-1687.	7.2	26
26	CD52 antigen expressed by malignant plasma cells can be targeted by alemtuzumab in vivo in NOD/SCID mice. Experimental Hematology, 2006, 34, 721-727.	0.4	25
27	Therapeutic effects of the combination of fenretinide and all-trans-retinoic acid and of the two retinoids with cisplatin in a human ovarian carcinoma xenograft and in a cisplatin-resistant sub-line. European Journal of Cancer, 2000, 36, 2411-2419.	2.8	22
28	Immunomodulatory Factors Control the Fate of Melanoma Tumor Initiating Cells. Stem Cells, 2016, 34, 2449-2460.	3.2	21
29	The Detection and Morphological Analysis of Circulating Tumor and Host Cells in Breast Cancer Xenograft Models. Cells, 2019, 8, 683.	4.1	21
30	Effect of verapamil on doxorubicin activity and pharmacokinetics in mice bearing resistant and sensitive solid tumors. Cancer Chemotherapy and Pharmacology, 1988, 21, 329-36.	2.3	19
31	IFN-γ Enhances the Antimyeloma Activity of the Fully Human Anti–Human Leukocyte Antigen-DR Monoclonal Antibody 1D09C3. Cancer Research, 2007, 67, 3269-3275.	0.9	18
32	Defibrotide in combination with granulocyte colony-stimulating factor significantly enhances the mobilization of primitive and committed peripheral blood progenitor cells in mice. Cancer Research, 2002, 62, 6152-7.	0.9	18
33	Targeting COPZ1 non-oncogene addiction counteracts the viability of thyroid tumor cells. Cancer Letters, 2017, 410, 201-211.	7.2	15
34	Gene signatures of circulating breast cancer cell models are a source of novel molecular determinants of metastasis and improve circulating tumor cell detection in patients. Journal of Experimental and Clinical Cancer Research, 2022, 41, 78.	8.6	15
35	Characterization of compound 584, an Abl kinase inhibitor with lasting effects. Haematologica, 2008, 93, 653-661.	3.5	14
36	Placental Growth Factor-1 Potentiates Hematopoietic Progenitor Cell Mobilization Induced by Granulocyte Colony-Stimulating Factor in Mice and Nonhuman Primates. Stem Cells, 2007, 25, 252-261.	3.2	12

LOREDANA CLERIS

#	Article	IF	CITATIONS
37	Effect of imatinib on haematopoietic recovery following idarubicin exposure. Leukemia, 2003, 17, 298-304.	7.2	11
38	D Quantification of Tumor Vasculature in Lymphoma Xenografts in NOD/SCID Mice Allows to Detect Differences among Vascular-Targeted Therapies. PLoS ONE, 2013, 8, e59691.	2.5	9
39	Verapamil potentiation of doxorubicin resistance development in B16 melanoma cells both in vitro and in vivo. British Journal of Cancer, 1988, 57, 343-347.	6.4	8
40	Mobilization of primitive and committed hematopoietic progenitors in nonhuman primates treated with defibrotide and recombinant human granulocyte colony-stimulating factor. Experimental Hematology, 2004, 32, 68-75.	0.4	7
41	Induction of death receptor 5 expression in tumor vasculature by perifosine restores the vascular disruption activity of TRAIL-expressing CD34+ cells. Angiogenesis, 2013, 16, 707-722.	7.2	5
42	Sodium 4-Carboxymethoxyimino-(4-HPR) a Novel Water-Soluble Derivative of 4-Oxo-4-HPR Endowed with In Vivo Anticancer Activity on Solid Tumors. Frontiers in Pharmacology, 2017, 8, 226.	3.5	5
43	Postsurgical adjuvant chemoimmunotherapy with recombinant interleukin-2 and 1,3-bis-(2-chloroethyl)-1-nitrosurea on spontaneous metastases of a non-immunogenic murine tumour. Cancer Immunology, Immunotherapy, 1992, 34, 383-388.	4.2	4
44	Senescent Thyrocytes, Similarly to Thyroid Tumor Cells, Elicit M2-like Macrophage Polarization In Vivo. Biology, 2021, 10, 985.	2.8	3
45	Waterâ€soluble derivatives of 4â€oxoâ€ <i>N</i> â€(4â€hydroxyphenyl) retinamide: synthesis and biological activity. Chemical Biology and Drug Design, 2016, 88, 608-614.	3.2	2
46	Antitumor Activity of Human CD34+Cells Expressing Membrane-Bound Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand. Human Gene Therapy, 2006, .	2.7	0