

Delphine Rea

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/181517/publications.pdf>

Version: 2024-02-01

134
papers

10,728
citations

50170

46
h-index

31759

101
g-index

139
all docs

139
docs citations

139
times ranked

7784
citing authors

#	ARTICLE	IF	CITATIONS
1	Discontinuation of imatinib in patients with chronic myeloid leukaemia who have maintained complete molecular remission for at least 2 years: the prospective, multicentre Stop Imatinib (STIM) trial. <i>Lancet Oncology</i> , The, 2010, 11, 1029-1035.	5.1	1,359
2	Imatinib mesylate discontinuation in patients with chronic myelogenous leukemia in complete molecular remission for more than 2 years. <i>Blood</i> , 2007, 109, 58-60.	0.6	505
3	Intermittent Target Inhibition With Dasatinib 100 mg Once Daily Preserves Efficacy and Improves Tolerability in Imatinib-Resistant and -Intolerant Chronic-Phase Chronic Myeloid Leukemia. <i>Journal of Clinical Oncology</i> , 2008, 26, 3204-3212.	0.8	458
4	Discontinuation of tyrosine kinase inhibitor therapy in chronic myeloid leukaemia (EURO-SKI): a prespecified interim analysis of a prospective, multicentre, non-randomised, trial. <i>Lancet Oncology</i> , The, 2018, 19, 747-757.	5.1	444
5	Ponatinib efficacy and safety in Philadelphia chromosomeâ€“positive leukemia: final 5-year results of the phase 2 PACE trial. <i>Blood</i> , 2018, 132, 393-404.	0.6	392
6	Long-Term Follow-Up of the French Stop Imatinib (STIM1) Study in Patients With Chronic Myeloid Leukemia. <i>Journal of Clinical Oncology</i> , 2017, 35, 298-305.	0.8	380
7	Imatinib plus Peginterferon Alfa-2a in Chronic Myeloid Leukemia. <i>New England Journal of Medicine</i> , 2010, 363, 2511-2521.	13.9	362
8	Loss of Major Molecular Response As a Trigger for Restarting Tyrosine Kinase Inhibitor Therapy in Patients With Chronic-Phase Chronic Myelogenous Leukemia Who Have Stopped Imatinib After Durable Undetectable Disease. <i>Journal of Clinical Oncology</i> , 2014, 32, 424-430.	0.8	355
9	Imatinib combined with induction or consolidation chemotherapy in patients with de novo Philadelphia chromosomeâ€“positive acute lymphoblastic leukemia: results of the GRAAPH-2003 study. <i>Blood</i> , 2007, 109, 1408-1413.	0.6	300
10	BCR-ABL1 Compound Mutations Combining Key Kinase Domain Positions Confer Clinical Resistance to Ponatinib in Ph Chromosome-Positive Leukemia. <i>Cancer Cell</i> , 2014, 26, 428-442.	7.7	292
11	Discontinuation of dasatinib or nilotinib in chronic myeloid leukemia: interim analysis of the STOP 2G-TKI study. <i>Blood</i> , 2017, 129, 846-854.	0.6	268
12	Asciminib in Chronic Myeloid Leukemia after ABL Kinase Inhibitor Failure. <i>New England Journal of Medicine</i> , 2019, 381, 2315-2326.	13.9	257
13	Vascular safety issues in CML patients treated with BCR/ABL1 kinase inhibitors. <i>Blood</i> , 2015, 125, 901-906.	0.6	239
14	Lung Abnormalities after Dasatinib Treatment for Chronic Myeloid Leukemia. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2007, 176, 814-818.	2.5	189
15	Expression of the Serpin Serine Protease Inhibitor 6 Protects Dendritic Cells from Cytotoxic T Lymphocyteâ€“Induced Apoptosis. <i>Journal of Experimental Medicine</i> , 2001, 194, 657-668.	4.2	187
16	Immature Dendritic Cells Acquire Cd8+Cytotoxic T Lymphocyte Priming Capacity upon Activation by T Helper Cellâ€“Independent orâ€“Dependent Stimuli. <i>Journal of Experimental Medicine</i> , 2000, 192, 145-150.	4.2	173
17	Glucocorticoids transform CD40-triggering of dendritic cells into an alternative activation pathway resulting in antigen-presenting cells that secrete IL-10. <i>Blood</i> , 2000, 95, 3162-3167.	0.6	154
18	Highly Efficient Transduction of Human Monocyte-Derived Dendritic Cells with Subgroup B Fiber-Modified Adenovirus Vectors Enhances Transgene-Encoded Antigen Presentation to Cytotoxic T Cells. <i>Journal of Immunology</i> , 2001, 166, 5236-5244.	0.4	149

#	ARTICLE	IF	CITATIONS
19	A phase 3, open-label, randomized study of asciminib, a STAMP inhibitor, vs bosutinib in CML after 2 or more prior TKIs. <i>Blood</i> , 2021, 138, 2031-2041.	0.6	147
20	Combined Treatment With Arsenic Trioxide and All-Trans-Retinoic Acid in Patients With Relapsed Acute Promyelocytic Leukemia. <i>Journal of Clinical Oncology</i> , 2003, 21, 2326-2334.	0.8	146
21	Severe Peripheral Arterial Disease During Nilotinib Therapy. <i>Journal of the National Cancer Institute</i> , 2011, 103, 1347-1348.	3.0	145
22	Dasatinib in imatinib-resistant or intolerant chronic-phase, chronic myeloid leukemia patients: 7-year follow-up of study CA180034. <i>American Journal of Hematology</i> , 2016, 91, 869-874.	2.0	145
23	Adenoviruses Activate Human Dendritic Cells without Polarization toward a T-Helper Type 1-Inducing Subset. <i>Journal of Virology</i> , 1999, 73, 10245-10253.	1.5	145
24	Long-Term Follow-Up of the Imatinib GRAAPH-2003 Study in Newly Diagnosed Patients with De Novo Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia: A GRAALL Study. <i>Biology of Blood and Marrow Transplantation</i> , 2013, 19, 150-155.	2.0	140
25	Combining the Allosteric Inhibitor Asciminib with Ponatinib Suppresses Emergence of and Restores Efficacy against Highly Resistant BCR-ABL1 Mutants. <i>Cancer Cell</i> , 2019, 36, 431-443.e5.	7.7	137
26	BCR/ABL Oncogene Directly Controls MHC Class I Chain-Related Molecule A Expression in Chronic Myelogenous Leukemia. <i>Journal of Immunology</i> , 2006, 176, 5108-5116.	0.4	126
27	Phase 2 study of subcutaneous omacetaxine mepesuccinate after TKI failure in patients with chronic-phase CML with T315I mutation. <i>Blood</i> , 2012, 120, 2573-2580.	0.6	123
28	In vivo production of interleukin-10 by malignant cells in AIDS lymphomas. <i>European Journal of Immunology</i> , 1992, 22, 2937-2942.	1.6	116
29	Recurrent ETNK1 mutations in atypical chronic myeloid leukemia. <i>Blood</i> , 2015, 125, 499-503.	0.6	115
30	Early onset hypercholesterolemia induced by the 2nd-generation tyrosine kinase inhibitor nilotinib in patients with chronic phase-chronic myeloid leukemia. <i>Haematologica</i> , 2014, 99, 1197-1203.	1.7	114
31	Natural killer-cell counts are associated with molecular relapse-free survival after imatinib discontinuation in chronic myeloid leukemia: the IMMUNOSTIM study. <i>Haematologica</i> , 2017, 102, 1368-1377.	1.7	114
32	Monomethylfumarate affects polarization of monocyte-derived dendritic cells resulting in down-regulated Th1 lymphocyte responses. <i>European Journal of Immunology</i> , 2004, 34, 565-575.	1.6	99
33	Clinical features of pulmonary arterial hypertension in patients receiving dasatinib. <i>American Journal of Hematology</i> , 2015, 90, 1060-1064.	2.0	98
34	Treatment-free remission with first- and second-generation tyrosine kinase inhibitors. <i>American Journal of Hematology</i> , 2019, 94, 346-357.	2.0	96
35	Dasatinib discontinuation in patients with chronic-phase chronic myeloid leukemia and stable deep molecular response: the DASFREE study. <i>Leukemia and Lymphoma</i> , 2020, 61, 650-659.	0.6	93
36	Second tyrosine kinase inhibitor discontinuation attempt in patients with chronic myeloid leukemia. <i>Cancer</i> , 2017, 123, 4403-4410.	2.0	85

#	ARTICLE	IF	CITATIONS
37	Deep molecular responses achieved in patients with CML-CP who are switched to nilotinib after long-term imatinib. <i>Blood</i> , 2014, 124, 729-736.	0.6	84
38	Evaluation of Residual Disease and TKI Duration Are Critical Predictive Factors for Molecular Recurrence after Stopping Imatinib First-line in Chronic Phase CML Patients. <i>Clinical Cancer Research</i> , 2019, 25, 6606-6613.	3.2	82
39	Defective blood dendritic cells in chronic myeloid leukemia correlate with high plasmatic VEGF and are not normalized by imatinib mesylate. <i>Leukemia</i> , 2004, 18, 1656-1661.	3.3	79
40	Prolongation of skin graft survival by modulation of the alloimmune response with alternatively activated dendritic cells ¹ . <i>Transplantation</i> , 2003, 76, 1608-1615.	0.5	71
41	High-dose imatinib mesylate combined with vincristine and dexamethasone (DIV regimen) as induction therapy in patients with resistant Philadelphia-positive acute lymphoblastic leukemia and lymphoid blast crisis of chronic myeloid leukemia. <i>Leukemia</i> , 2006, 20, 400-403.	3.3	67
42	Discontinuation of tyrosine kinase inhibitors in chronic myeloid leukemia: Recommendations for clinical practice from the French Chronic Myeloid Leukemia Study Group. <i>Cancer</i> , 2018, 124, 2956-2963.	2.0	63
43	Expert opinion on management of chronic myeloid leukemia after resistance to second-generation tyrosine kinase inhibitors. <i>Leukemia</i> , 2020, 34, 1495-1502.	3.3	63
44	Incidence, outcomes, and risk factors of pleural effusion in patients receiving dasatinib therapy for Philadelphia chromosome-positive leukemia. <i>Haematologica</i> , 2019, 104, 93-101.	1.7	62
45	Leukemic stem cell persistence in chronic myeloid leukemia patients in deep molecular response induced by tyrosine kinase inhibitors and the impact of therapy discontinuation. <i>Oncotarget</i> , 2016, 7, 35293-35301.	0.8	54
46	Adherence to oral tyrosine kinase inhibitor therapies in chronic myeloid leukemia. <i>Leukemia Research</i> , 2012, 36, 817-825.	0.4	51
47	Management of adverse events associated with tyrosine kinase inhibitors in chronic myeloid leukemia. <i>Annals of Hematology</i> , 2015, 94, 149-158.	0.8	48
48	Nilotinib and peginterferon alfa-2a for newly diagnosed chronic-phase chronic myeloid leukaemia (NiloPeg): a multicentre, non-randomised, open-label phase 2 study. <i>Lancet Haematology</i> , 2015, 2, e37-e46.	2.2	45
49	Final analysis of the efficacy and safety of omacetaxine mepesuccinate in patients with chronic or accelerated phase chronic myeloid leukemia: Results with 24 months of follow-up. <i>Cancer</i> , 2015, 121, 1637-1644.	2.0	44
50	Deterioration of pulmonary hypertension and pleural effusion with bosutinib following dasatinib lung toxicity. <i>European Respiratory Journal</i> , 2016, 48, 1517-1519.	3.1	44
51	Second transplant with two unrelated cord blood units for early graft failure after haematopoietic stem cell transplantation. <i>British Journal of Haematology</i> , 2007, 137, 248-251.	1.2	41
52	Preliminary Report Of The STIM2 Study: A Multicenter Stop Imatinib Trial For Chronic Phase Chronic Myeloid Leukemia De Novo Patients On Imatinib. <i>Blood</i> , 2013, 122, 654-654.	0.6	41
53	Intermediate maturation of Mycobacterium tuberculosis LAM-activated human dendritic cells. <i>Cellular Microbiology</i> , 2007, 9, 1412-1425.	1.1	40
54	Usefulness of the 2012 European CVD risk assessment model to identify patients at high risk of cardiovascular events during nilotinib therapy in chronic myeloid leukemia. <i>Leukemia</i> , 2015, 29, 1206-1209.	3.3	38

#	ARTICLE	IF	CITATIONS
55	Ponatinib in chronic myeloid leukemia (CML): Consensus on patient treatment and management from a European expert panel. <i>Critical Reviews in Oncology/Hematology</i> , 2017, 120, 52-59.	2.0	38
56	First-line imatinib mesylate in patients with newly diagnosed accelerated phase-chronic myeloid leukemia. <i>Leukemia</i> , 2012, 26, 2254-2259.	3.3	37
57	Treatment by Lenalidomide in lower risk myelodysplastic syndrome with 5q deletion – The GFM experience. <i>Leukemia Research</i> , 2011, 35, 1444-1448.	0.4	36
58	Treatment-free remission in patients with chronic myeloid leukemia. <i>International Journal of Hematology</i> , 2018, 108, 355-364.	0.7	35
59	ABO-mismatched marrow processing for transplantation: results of 114 procedures and analysis of immediate adverse events and hematopoietic recovery. <i>Transfusion</i> , 2006, 46, 398-402.	0.8	34
60	Ponatinib evaluation and safety in real-life chronic myelogenous leukemia patients failing more than two tyrosine kinase inhibitors: the PEARL observational study. <i>Experimental Hematology</i> , 2018, 67, 41-48.	0.2	34
61	Longer treatment duration and history of osteoarticular symptoms predispose to tyrosine kinase inhibitor withdrawal syndrome. <i>British Journal of Haematology</i> , 2019, 187, 337-346.	1.2	31
62	A specific time course for mobilization of peripheral blood CD34+ cells after plerixafor injection in very poor mobilizer patients: impact on the timing of the apheresis procedure. <i>Transfusion</i> , 2013, 53, 564-569.	0.8	30
63	Impact of age on efficacy and toxicity of nilotinib in patients with chronic myeloid leukemia in chronic phase: ENEST1st subanalysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 1585-1596.	1.2	29
64	The addition of daunorubicin to imatinib mesylate in combination with cytarabine improves the response rate and the survival of patients with myeloid blast crisis chronic myelogenous leukemia (AFRO1 study). <i>Leukemia Research</i> , 2011, 35, 777-782.	0.4	27
65	How I manage relapse of chronic myeloid leukaemia after stopping tyrosine kinase inhibitor therapy. <i>British Journal of Haematology</i> , 2018, 180, 24-32.	1.2	27
66	Cartridge-based automated BCR-ABL1 mRNA quantification: solving the issues of standardization, at what cost?. <i>Haematologica</i> , 2011, 96, 664-671.	1.7	25
67	Long-term safety and efficacy of imatinib mesylate (Gleevec®) in elderly patients with chronic phase chronic myelogenous leukemia: Results of the AFR04 study. <i>American Journal of Hematology</i> , 2013, 88, 1-4.	2.0	25
68	Rapid onset of peripheral artery disease in a chronic myeloid leukemia patient without prior arterial disorder: direct relationship with nilotinib exposure and clinical outcome. <i>European Journal of Haematology</i> , 2015, 94, 363-367.	1.1	25
69	Imatinib Increases Serum Creatinine by Inhibiting Its Tubular Secretion in a Reversible Fashion in Chronic Myeloid Leukemia. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2016, 16, 169-174.	0.2	25
70	Curing Chronic Myeloid Leukemia. <i>Current Hematologic Malignancy Reports</i> , 2012, 7, 103-108.	1.2	24
71	Impact of NFE2 mutations on AML transformation and overall survival in patients with myeloproliferative neoplasms. <i>Blood</i> , 2021, 138, 2142-2148.	0.6	23
72	Treatment-Free Remission (TFR) in Patients with Chronic Phase Chronic Myeloid Leukemia (CML-CP) and in Stable Deep Molecular Response (DMR) to Dasatinib - the Dasfree Study. <i>Blood</i> , 2016, 128, 1895-1895.	0.6	23

#	ARTICLE	IF	CITATIONS
73	Omacetaxine mepesuccinate for patients with accelerated phase chronic myeloid leukemia with resistance or intolerance to two or more tyrosine kinase inhibitors. <i>Haematologica</i> , 2013, 98, e78-e79.	1.7	22
74	ETNK1 mutations induce a mutator phenotype that can be reverted with phosphoethanolamine. <i>Nature Communications</i> , 2020, 11, 5938.	5.8	22
75	Dasatinib dose optimisation based on therapeutic drug monitoring reduces pleural effusion rates in chronic myeloid leukaemia patients. <i>British Journal of Haematology</i> , 2021, 194, 393-402.	1.2	22
76	ABL001, a Potent, Allosteric Inhibitor of BCR-ABL, Exhibits Safety and Promising Single-Agent Activity in a Phase I Study of Patients with CML with Failure of Prior TKI Therapy. <i>Blood</i> , 2015, 126, 138-138.	0.6	22
77	Development of asciminib, a novel allosteric inhibitor of BCR-ABL1. <i>Critical Reviews in Oncology/Hematology</i> , 2022, 171, 103580.	2.0	21
78	Dasatinib-induced lupus. <i>Lancet</i> , The, 2008, 372, 713-714.	6.3	20
79	Expanded Phase 1 Study of ABL001, a Potent, Allosteric Inhibitor of BCR-ABL, Reveals Significant and Durable Responses in Patients with CML-Chronic Phase with Failure of Prior TKI Therapy. <i>Blood</i> , 2016, 128, 625-625.	0.6	20
80	Imatinib mesylate minimally affects bcr-abl+ and normal monocyte-derived dendritic cells but strongly inhibits T cell expansion despite reciprocal dendritic cell-T cell activation. <i>Journal of Leukocyte Biology</i> , 2006, 79, 747-756.	1.5	19
81	Bone marrow mesenchymal stromal cell (MSC) gene profiling in chronic myeloid leukemia (CML) patients at diagnosis and in deep molecular response induced by tyrosine kinase inhibitors (TKIs). <i>Leukemia Research</i> , 2017, 60, 94-102.	0.4	19
82	Strategies for improved antigen delivery into dendritic cells. <i>Trends in Molecular Medicine</i> , 2001, 7, 91-94.	3.5	17
83	COVID-19 in Patients (pts) with Chronic Myeloid Leukemia (CML): Results from the International CML Foundation (iCMLf) CML and COVID-19 (CANDID) Study. <i>Blood</i> , 2020, 136, 46-47.	0.6	17
84	Prospective flow cytometric evaluation of nucleated red blood cells in cord blood units and relationship with nucleated and CD34+ cell quantification. <i>Transfusion</i> , 2006, 46, 403-406.	0.8	16
85	Tolerability and efficacy of pegylated interferon- α 2a in combination with imatinib for patients with chronic-phase chronic myeloid leukemia. <i>Cancer</i> , 2013, 119, 4284-4289.	2.0	16
86	Recombinant adenovirus-transduced human dendritic cells engineered to secrete interleukin-10 (IL-10) suppress Th1-type responses while selectively activating IL-10-producing CD4+ T cells. <i>Human Immunology</i> , 2004, 65, 1344-1355.	1.2	15
87	Long-term outcome of imatinib 400 mg compared to imatinib 600 mg or imatinib 400 mg daily in combination with cytarabine or pegylated interferon alpha 2a for chronic myeloid leukaemia: results from the French SPIRIT phase III randomised trial. <i>Leukemia</i> , 2021, 35, 2332-2345.	3.3	15
88	Undetectable molecular residual disease after omacetaxine and nilotinib combination therapy in an imatinib-resistant chronic myeloid leukaemia patient harbouring the BCR-ABL1 T315I gatekeeper mutation. <i>British Journal of Haematology</i> , 2012, 157, 407-410.	1.2	14
89	Reversible lymph node follicular hyperplasia associated with dasatinib treatment of chronic myeloid leukemia in chronic phase. <i>Blood</i> , 2013, 122, 3082-3084.	0.6	14
90	Integrated Genomic, Functional, and Prognostic Characterization of Atypical Chronic Myeloid Leukemia. <i>HemaSphere</i> , 2020, 4, e497.	1.2	14

#	ARTICLE	IF	CITATIONS
91	Osteoarticular Pain after Discontinuation of Tyrosine Kinase Inhibitors (TKI): A French Cohort. <i>Blood</i> , 2015, 126, 137-137.	0.6	14
92	Insulin resistance is an underlying mechanism of impaired glucose metabolism during nilotinib therapy. <i>American Journal of Hematology</i> , 2018, 93, E342-E345.	2.0	13
93	Killer immunoglobulin-like receptor genotypes and chronic myeloid leukemia outcomes after imatinib cessation for treatment-free remission. <i>Cancer Medicine</i> , 2019, 8, 4976-4985.	1.3	13
94	The self peptide annexin II (208-223) presented by dendritic cells sensitizes autologous CD4+ T lymphocytes to recognize melanoma cells. <i>Cancer Immunology, Immunotherapy</i> , 2001, 49, 671-678.	2.0	11
95	Reply to J. Richter et al. <i>Journal of Clinical Oncology</i> , 2014, 32, 2823-2825.	0.8	11
96	ENESTPath: A Phase 3 Study to Assess the Effect of Nilotinib Treatment Duration on Treatment-Free Remission (TFR) in Patients with Chronic Myeloid Leukemia in Chronic Phase (CML-CP) Previously Treated with Imatinib: 24-Month Analysis of the First 300 Patients in the Induction/Consolidation Phase. <i>Blood</i> , 2016, 128, 3094-3094.	0.6	11
97	Calibration of BCR-ABL1 mRNA quantification methods using genetic reference materials is a valid strategy to report results on the international scale. <i>Clinical Biochemistry</i> , 2014, 47, 1333-1336.	0.8	10
98	Towards a Personalized Treatment of Patients with Chronic Myeloid Leukemia. <i>Current Hematologic Malignancy Reports</i> , 2019, 14, 492-500.	1.2	10
99	Combination of Dasatinib and Peg-Interferon Alpha 2b in Chronic Phase Chronic Myeloid Leukemia (CP-CML) First Line: Preliminary Results of a Phase II Trial, from the French Intergroup of CML (Fi-LMC). <i>Blood</i> , 2015, 126, 134-134.	0.6	10
100	Nilotinib first-line therapy in patients with Philadelphia chromosome-negative/BCR-ABL-positive chronic myeloid leukemia in chronic phase: ENEST1st sub-analysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 1225-1233.	1.2	9
101	Imatinib Sensitizes T-cell Lymphocytes From Chronic Myeloid Leukemia Patients to FasL-induced Cell Death. <i>Journal of Immunotherapy</i> , 2012, 35, 154-158.	1.2	8
102	Ponatinib In Patients (pts) With Chronic Myeloid Leukemia (CML) and Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia (Ph+ ALL) Resistant Or Intolerant To Dasatinib Or Nilotinib, Or With The T315I BCR-ABL Mutation: 2-Year Follow-Up Of The PACE Trial. <i>Blood</i> , 2013, 122, 650-650.	0.6	8
103	Recommandations du groupe FI-LMC pour la prise en charge des patients présentant des mutations du domaine tyrosine kinase de BCR-ABL dans les leucémies malignes à chromosome Philadelphia. <i>Hématologie</i> , 2010, 16, 65-79.	0.0	7
104	Efficacy and Safety of Ponatinib in CP-CML Patients By Number of Prior Tyrosine Kinase Inhibitors: 4-Year Follow-up of the Phase 2 PACE Trial. <i>Blood</i> , 2015, 126, 4025-4025.	0.6	7
105	Enestpath: A Phase III Study to Assess the Effect of Nilotinib Treatment Duration on Treatment-Free Remission (TFR) in Chronic Phase-Chronic Myeloid Leukemia (CP-CML) Patients (pts) Previously Treated with Imatinib: Interim Analysis from the First Year of Induction Phase. <i>Blood</i> , 2015, 126, 4040-4040.	0.6	7
106	Chronic Myeloid Leukemia Diagnosed during Pregnancy: Therapy, Outcomes and Follow-up. <i>Blood</i> , 2018, 132, 4255-4255.	0.6	6
107	COVID-19 in Patients with Chronic Myeloid Leukemia: Poor Outcomes for Patients with Comorbidities, Older Age, Advanced Phase Disease, and Those from Low-Income Countries: An Update of the Candid Study. <i>Blood</i> , 2021, 138, 634-634.	0.6	5
108	Handling challenging questions in the management of chronic myeloid leukemia: when is it safe to stop tyrosine kinase inhibitors?. <i>Blood Advances</i> , 2020, 4, 5589-5594.	2.5	4

#	ARTICLE	IF	CITATIONS
109	Ponatinib long-term follow-up of efficacy and safety in CP-CML patients in real world settings in France: The POST-PACE study. <i>Leukemia Research</i> , 2021, 104, 106541.	0.4	4
110	Combined Chemotherapy (daunorubicin + cytarabine) and Dasatinib as Salvage Therapy of Chronic Myeloid Leukemia (CML) in Myeloid Blast Crisis, a Pilot Study.. <i>Blood</i> , 2009, 114, 2195-2195.	0.6	4
111	<i>SF3B1</i> mutations in the Driver Clone Increase the Risk of Evolution to Myelofibrosis in Patients with Myeloproliferative Neoplasms (MPN). <i>Blood</i> , 2020, 136, 1-1.	0.6	4
112	Novel fusion between the breakpoint cluster region and platelet-derived growth factor receptor-alpha genes in a patient with chronic myeloid leukemia-like neoplasm: undetectable residual disease after imatinib therapy. <i>European Journal of Haematology</i> , 2015, 95, 480-483.	1.1	2
113	Pegylated Interferon-Alpha 2a in Combination with Nilotinib As First-Line Therapy in Newly Diagnosed Chronic Phase Chronic Myelogenous Leukemia (Nilopeg trial). Four-Year Follow-up Results. <i>Blood</i> , 2015, 126, 1578-1578.	0.6	2
114	NFE2 Mutations Impact AML Transformation and Overall Survival in Patients with Myeloproliferative Neoplasms (MPN). <i>Blood</i> , 2020, 136, 36-36.	0.6	2
115	Ruxolitinib Treatment Is Associated with Increased Incidence of Infections and Higher Risk of HSV/Vzv Recurrence in Patients with Myeloproliferative Neoplasm (MPN) Related Myelofibrosis (MF). <i>Blood</i> , 2020, 136, 8-8.	0.6	2
116	Quantification of nucleated red blood cells in allogeneic marrow graft and impact of processing on recovery. <i>Transfusion</i> , 2007, 47, 266-271.	0.8	1
117	Aspects pratiques des traitements par inhibiteurs de tyrosine kinase dans la leucémie myéloïde chronique. <i>Hématologie</i> , 2009, 15, 197-202.	0.0	1
118	Association of Vemurafenib and Pipobroman Enhances BRAF-CRAF Dimerization in Squamous Cell Carcinoma. <i>Journal of Investigative Dermatology</i> , 2016, 136, 1302-1305.	0.3	1
119	Thrombocytapheresis and sequential chemotherapy for extreme symptomatic thrombocytosis secondary to myelofibrosis: a case report. <i>Annals of Hematology</i> , 2020, 99, 897-898.	0.8	1
120	Impact of Age on Efficacy and Toxicity of Nilotinib in Patients with Chronic Myeloid Leukemia in Chronic Phase (CML-CP): ENEST1st Sub-Analysis. <i>Blood</i> , 2015, 126, 479-479.	0.6	1
121	What is treatment free remission in chronic myeloid leukemia?. <i>Oncotarget</i> , 2018, 9, 4279-4279.	0.8	1
122	ETNK1 Is an Early Event and SETBP1 a Late Event in Atypical Chronic Myeloid Leukemia. <i>Blood</i> , 2015, 126, 3652-3652.	0.6	1
123	ETNK1 Mutations in Atypical Chronic Myeloid Leukemia Induce a Mutator Phenotype That Can be Reverted with Phosphoethanolamine. <i>Blood</i> , 2020, 136, LBA-5-LBA-5.	0.6	1
124	Management of Adverse Events Associated with ATP-Competitive BCR-ABL1 Tyrosine Kinase Inhibitors in Chronic Myeloid Leukemia. <i>Hematologic Malignancies</i> , 2016, , 71-87.	0.2	0
125	Management of ITK pulmonary and pleural adverse effects: Fi-LMC guidelines. <i>Hématologie</i> , 2018, 24, 134-144.	0.0	0
126	Handling challenging questions in the management of chronic myeloid leukemia: when is it safe to stop tyrosine kinase inhibitors?. <i>Hematology American Society of Hematology Education Program</i> , 2020, 2020, 243-247.	0.9	0

#	ARTICLE	IF	CITATIONS
127	Adverse Events Associated with ATP-Competitive BCR-ABL1 Tyrosine Kinase Inhibitors in Chronic Myeloid Leukemia. Hematologic Malignancies, 2021, , 77-91.	0.2	0
128	Evidence of ETNK1 Somatic Variants in Atypical Chronic Myeloid Leukemia. Blood, 2014, 124, 2212-2212.	0.6	0
129	Evaluation of the Benefit/Risk Profile of Ponatinib in CP-CML Patients over Time: 4-Year Follow-up of the Phase 2 PACE Study. Blood, 2015, 126, 5142-5142.	0.6	0
130	Molecular Response with Nilotinib in Patients with Philadelphia Negative (Ph-) Chronic Myeloid Leukemia in Chronic Phase (CML-CP): ENEST1st Sub-Analysis. Blood, 2015, 126, 4054-4054.	0.6	0
131	Ponatinib for Chronic Phase (CP) CML Failing Two or More Tyrosine Kinase Inhibitors (TKI) or Harboring a T315I Mutation in the Real Life: Pearl Observational Study. Blood, 2015, 126, 4039-4039.	0.6	0
132	<i>The Outcome of Treatment-Free Remission after First-Line Nilotinib or Dasatinib in Chronic Phase Chronic Myeloid Leukemia Patients Is Different</i>. Blood, 2021, 138, 2552-2552.	0.6	0
133	Treatment Free Survival (TFS) in Patients (pts) with Chronic Myeloid Leukemia (CML) Carrying Atypical BCR-ABL1 Fusion Transcripts: The French CML Group (Fi-LMC) Experience. Blood, 2021, 138, 3604-3604.	0.6	0
134	Patient and Physician Perspectives of Unmet Needs in CML - Designing the CML SUN Survey. Blood, 2021, 138, 4986-4986.	0.6	0