

Feibo B Wu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1819544/publications.pdf>

Version: 2024-02-01

145
papers

8,695
citations

36203

51
h-index

49773

87
g-index

147
all docs

147
docs citations

147
times ranked

7637
citing authors

#	ARTICLE	IF	CITATIONS
1	An miR156-regulated nucleobase-ascorbate transporter 2 confers cadmium tolerance via enhanced anti-oxidative capacity in barley. <i>Journal of Advanced Research</i> , 2023, 44, 23-37.	4.4	11
2	Identification of low grain cadmium accumulation genotypes and its physiological mechanism in maize (<i>Zea mays</i> L.). <i>Environmental Science and Pollution Research</i> , 2022, 29, 20721-20730.	2.7	8
3	The genome and gene editing system of sea barleygrass provide a novel platform for cereal domestication and stress tolerance studies. <i>Plant Communications</i> , 2022, 3, 100333.	3.6	8
4	An ATP binding cassette transporter HvABCB25 confers aluminum detoxification in wild barley. <i>Journal of Hazardous Materials</i> , 2021, 401, 123371.	6.5	33
5	Metalloid hazards: From plant molecular evolution to mitigation strategies. <i>Journal of Hazardous Materials</i> , 2021, 409, 124495.	6.5	29
6	Evolution of rapid blue-light response linked to explosive diversification of ferns in angiosperm forests. <i>New Phytologist</i> , 2021, 230, 1201-1213.	3.5	33
7	Metabolome Analysis Revealed the Mechanism of Exogenous Glutathione to Alleviate Cadmium Stress in Maize (<i>Zea mays</i> L.) Seedlings. <i>Plants</i> , 2021, 10, 105.	1.6	23
8	Comparative transcriptome and tolerance mechanism analysis in the two contrasting wheat (<i>Triticum</i>) Tj ETQq0 0 0 rgBT /Overlock 10 T 101-114.	1.8	28
9	Strigolactone GR24 improves cadmium tolerance by regulating cadmium uptake, nitric oxide signaling and antioxidant metabolism in barley (<i>Hordeum vulgare</i> L.). <i>Environmental Pollution</i> , 2021, 273, 116486.	3.7	54
10	Differences in Grain Microstructure and Proteomics of a Broad Bean (<i>Vicia faba</i> L.) Landrace Cixidabaican in China Compared with Lingxiyicun Introduced from Japan. <i>Plants</i> , 2021, 10, 1385.	1.6	4
11	Mechanistic Insights into Potassium-Conferred Drought Stress Tolerance in Cultivated and Tibetan Wild Barley: Differential Osmoregulation, Nutrient Retention, Secondary Metabolism and Antioxidative Defense Capacity. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13100.	1.8	7
12	Genome-Wide Discovery of miRNAs with Differential Expression Patterns in Responses to Salinity in the Two Contrasting Wheat Cultivars. <i>International Journal of Molecular Sciences</i> , 2021, 22, 12556.	1.8	10
13	Silicon regulates the expression of vacuolar H ⁺ -pyrophosphatase 1 and decreases cadmium accumulation in rice (<i>Oryza sativa</i> L.). <i>Chemosphere</i> , 2020, 240, 124907.	4.2	40
14	Genotypic difference in secondary metabolism-related enzyme activities and their relative gene expression patterns, osmolyte and plant hormones in wheat. <i>Physiologia Plantarum</i> , 2020, 168, 921-933.	2.6	6
15	Overexpression of HvAKT1 improves drought tolerance in barley by regulating root ion homeostasis and ROS and NO signaling. <i>Journal of Experimental Botany</i> , 2020, 71, 6587-6600.	2.4	31
16	Breeding for low cadmium accumulation cereals. <i>Journal of Zhejiang University: Science B</i> , 2020, 21, 442-459.	1.3	16
17	Optimized Protocol for OnGuard2 Software in Studying Guard Cell Membrane Transport and Stomatal Physiology. <i>Frontiers in Plant Science</i> , 2020, 11, 131.	1.7	0
18	CO ₂ enrichment using CRAM fermentation improves growth, physiological traits and yield of cherry tomato (<i>Solanum lycopersicum</i> L.). <i>Saudi Journal of Biological Sciences</i> , 2020, 27, 1041-1048.	1.8	15

#	ARTICLE	IF	CITATIONS
19	The Barley S-Adenosylmethionine Synthetase 3 Gene HvSAMS3 Positively Regulates the Tolerance to Combined Drought and Salinity Stress in Tibetan Wild Barley. <i>Cells</i> , 2020, 9, 1530.	1.8	20
20	HvHOX9, a novel homeobox leucine zipper transcription factor, positively regulates aluminum tolerance in Tibetan wild barley. <i>Journal of Experimental Botany</i> , 2020, 71, 6057-6073.	2.4	19
21	Comparison of Biochemical, Anatomical, Morphological, and Physiological Responses to Salinity Stress in Wheat and Barley Genotypes Deferring in Salinity Tolerance. <i>Agronomy</i> , 2020, 10, 127.	1.3	119
22	HvAKT2 and HvHAK1 confer drought tolerance in barley through enhanced leaf mesophyll H ⁺ homeostasis. <i>Plant Biotechnology Journal</i> , 2020, 18, 1683-1696.	4.1	54
23	Genome-Wide Identification and Characterization of Drought Stress Responsive microRNAs in Tibetan Wild Barley. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2795.	1.8	29
24	Resemblance and Difference of Seedling Metabolic and Transporter Gene Expression in High Tolerance Wheat and Barley Cultivars in Response to Salinity Stress. <i>Plants</i> , 2020, 9, 519.	1.6	18
25	Agriculture organic wastes fermentation CO ₂ enrichment in greenhouse and the fermentation residues improve growth, yield and fruit quality in tomato. <i>Journal of Cleaner Production</i> , 2020, 275, 123885.	4.6	29
26	Comparative physiological analysis in the tolerance to salinity and drought individual and combination in two cotton genotypes with contrasting salt tolerance. <i>Physiologia Plantarum</i> , 2019, 165, 155-168.	2.6	46
27	Exogenous hydrogen sulfide reduces cadmium uptake and alleviates cadmium toxicity in barley. <i>Plant Growth Regulation</i> , 2019, 89, 227-237.	1.8	48
28	Genome-Wide Identification, Characterization and Expression Analysis of Xyloglucan Endotransglucosylase/Hydrolase Genes Family in Barley (<i>Hordeum vulgare</i>). <i>Molecules</i> , 2019, 24, 1935.	1.7	29
29	Foliar application of betaine improves water-deficit stress tolerance in barley (<i>Hordeum vulgare</i> L.). <i>Plant Growth Regulation</i> , 2019, 89, 109-118.	1.8	22
30	Transient silencing of an expansin HvEXPA1 inhibits root cell elongation and reduces Al accumulation in root cell wall of Tibetan wild barley. <i>Environmental and Experimental Botany</i> , 2019, 165, 120-128.	2.0	17
31	Chloride transport at plant-soil Interface modulates barley cd tolerance. <i>Plant and Soil</i> , 2019, 441, 409-421.	1.8	12
32	Genome-wide characterization of drought stress responsive long non-coding RNAs in Tibetan wild barley. <i>Environmental and Experimental Botany</i> , 2019, 164, 124-134.	2.0	31
33	Exogenous Glycinebetaine Reduces Cadmium Uptake and Mitigates Cadmium Toxicity in Two Tobacco Genotypes Differing in Cadmium Tolerance. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1612.	1.8	33
34	HvPAA1 Encodes a P-Type ATPase, a Novel Gene for Cadmium Accumulation and Tolerance in Barley (<i>Hordeum vulgare</i> L.). <i>International Journal of Molecular Sciences</i> , 2019, 20, 1732.	1.8	20
35	Response of Tibetan Wild Barley Genotypes to Drought Stress and Identification of Quantitative Trait Loci by Genome-Wide Association Analysis. <i>International Journal of Molecular Sciences</i> , 2019, 20, 791.	1.8	15
36	Evolution of chloroplast retrograde signaling facilitates green plant adaptation to land. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 5015-5020.	3.3	138

#	ARTICLE	IF	CITATIONS
37	Leaf epidermis transcriptome reveals drought-Induced hormonal signaling for stomatal regulation in wild barley. <i>Plant Growth Regulation</i> , 2019, 87, 39-54.	1.8	29
38	Genotypic differences in leaf secondary metabolism, plant hormones and yield under alone and combined stress of drought and salinity in cotton genotypes. <i>Physiologia Plantarum</i> , 2019, 165, 343-355.	2.6	71
39	Differences in physiological and biochemical characteristics in response to single and combined drought and salinity stresses between wheat genotypes differing in salt tolerance. <i>Physiologia Plantarum</i> , 2019, 165, 134-143.	2.6	66
40	Application of sulfur fertilizer reduces cadmium accumulation and toxicity in tobacco seedlings (<i>Nicotiana tabacum</i>). <i>Plant Growth Regulation</i> , 2018, 85, 165-170.	1.8	16
41	Tolerance to Drought, Low pH and Al Combined Stress in Tibetan Wild Barley Is Associated with Improvement of ATPase and Modulation of Antioxidant Defense System. <i>International Journal of Molecular Sciences</i> , 2018, 19, 3553.	1.8	12
42	Genotypic difference in the influence of aluminum and low pH on ion flux, rhizospheric pH and ATPase activity between Tibetan wild and cultivated barley. <i>Environmental and Experimental Botany</i> , 2018, 156, 16-24.	2.0	7
43	Genotype-dependent effects of phosphorus supply on physiological and biochemical responses to Al-stress in cultivated and Tibetan wild barley. <i>Plant Growth Regulation</i> , 2017, 82, 259-270.	1.8	6
44	Genotype-dependent alleviation effects of exogenous GSH on salinity stress in cotton is related to improvement in chlorophyll content, photosynthetic performance, and leaf/root ultrastructure. <i>Environmental Science and Pollution Research</i> , 2017, 24, 9417-9427.	2.7	27
45	Genotypic-dependent effects of N fertilizer, glutathione, silicon, zinc, and selenium on proteomic profiles, amino acid contents, and quality of rice genotypes with contrasting grain Cd accumulation. <i>Functional and Integrative Genomics</i> , 2017, 17, 387-397.	1.4	9
46	Exploration and Utilization of Drought-Tolerant Barley Germplasm. , 2016, , 115-152.		3
47	The regulation of root growth in response to phosphorus deficiency mediated by phytohormones in a Tibetan wild barley accession. <i>Acta Physiologiae Plantarum</i> , 2016, 38, 1.	1.0	16
48	Identification of the differentially accumulated proteins associated with low phosphorus tolerance in a Tibetan wild barley accession. <i>Journal of Plant Physiology</i> , 2016, 198, 10-22.	1.6	10
49	Identification and comparative analysis of the microRNA transcriptome in roots of two contrasting tobacco genotypes in response to cadmium stress. <i>Scientific Reports</i> , 2016, 6, 32805.	1.6	37
50	Drought Tolerant Wild Species Are the Important Sources of Genes and Molecular Mechanisms Studies: Implication for Developing Drought Tolerant Crops. , 2016, , 401-426.		0
51	Physiological and molecular analysis on root growth associated with the tolerance to aluminum and drought individual and combined in Tibetan wild and cultivated barley. <i>Planta</i> , 2016, 243, 973-985.	1.6	22
52	K ⁺ Uptake, H ⁺ -ATPase pumping activity and Ca ²⁺ efflux mechanism are involved in drought tolerance of barley. <i>Environmental and Experimental Botany</i> , 2016, 129, 57-66.	2.0	43
53	Cr-induced changes in leaf protein profile, ultrastructure and photosynthetic traits in the two contrasting tobacco genotypes. <i>Plant Growth Regulation</i> , 2016, 79, 147-156.	1.8	25
54	DNA microarray revealed and RNAi plants confirmed key genes conferring low Cd accumulation in barley grains. <i>BMC Plant Biology</i> , 2015, 15, 259.	1.6	28

#	ARTICLE	IF	CITATIONS
55	Genome-wide identification of chromium stress-responsive micro RNAs and their target genes in tobacco (<i>Nicotiana tabacum</i>) roots. <i>Environmental Toxicology and Chemistry</i> , 2015, 34, 2573-2582.	2.2	50
56	<i>HvEXPB7</i> , a novel β -expansin gene revealed by the root hair transcriptome of Tibetan wild barley, improves root hair growth under drought stress. <i>Journal of Experimental Botany</i> , 2015, 66, 7405-7419.	2.4	94
57	Genotypic differences in photosynthetic performance, antioxidant capacity, ultrastructure and nutrients in response to combined stress of salinity and Cd in cotton. <i>BioMetals</i> , 2015, 28, 1063-1078.	1.8	29
58	Differences in photosynthesis, yield and grain cadmium accumulation as affected by exogenous cadmium and glutathione in the two rice genotypes. <i>Plant Growth Regulation</i> , 2015, 75, 715-723.	1.8	84
59	Physiological and biochemical responses to drought stress in cultivated and Tibetan wild barley. <i>Plant Growth Regulation</i> , 2015, 75, 567-574.	1.8	71
60	Difference in physiological and biochemical responses to salt stress between Tibetan wild and cultivated barleys. <i>Acta Physiologiae Plantarum</i> , 2015, 37, 1.	1.0	9
61	Tolerance to Combined Stress of Drought and Salinity in Barley. , 2015, , 93-121.		6
62	Comparative proteomic analysis of drought tolerance in the two contrasting Tibetan wild genotypes and cultivated genotype. <i>BMC Genomics</i> , 2015, 16, 432.	1.2	57
63	Modulation of exogenous selenium in cadmium-induced changes in antioxidative metabolism, cadmium uptake, and photosynthetic performance in the 2 tobacco genotypes differing in cadmium tolerance. <i>Environmental Toxicology and Chemistry</i> , 2015, 34, 92-99.	2.2	70
64	Secondary metabolism and antioxidants are involved in the tolerance to drought and salinity, separately and combined, in Tibetan wild barley. <i>Environmental and Experimental Botany</i> , 2015, 111, 1-12.	2.0	129
65	Comparative proteomic analysis of two tobacco (<i>Nicotiana tabacum</i>) genotypes differing in Cd tolerance. <i>BioMetals</i> , 2014, 27, 1277-1289.	1.8	16
66	Transcriptome profiling reveals mosaic genomic origins of modern cultivated barley. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 13403-13408.	3.3	74
67	N-acetyl-cysteine alleviates Cd toxicity and reduces Cd uptake in the two barley genotypes differing in Cd tolerance. <i>Plant Growth Regulation</i> , 2014, 74, 93-105.	1.8	18
68	Variation in β -amylase activity and thermostability in Tibetan annual wild and cultivated barley genotypes. <i>Journal of Zhejiang University: Science B</i> , 2014, 15, 801-808.	1.3	5
69	Genotypic and environmental variation in cadmium, chromium, lead and copper in rice and approaches for reducing the accumulation. <i>Science of the Total Environment</i> , 2014, 496, 275-281.	3.9	81
70	MALDI-TOF mass spectrometry provides an efficient approach to monitoring protein modification in the malting process. <i>International Journal of Mass Spectrometry</i> , 2014, 371, 8-16.	0.7	8
71	Genome-wide transcriptome and functional analysis of two contrasting genotypes reveals key genes for cadmium tolerance in barley. <i>BMC Genomics</i> , 2014, 15, 611.	1.2	101
72	Physiological and proteomic alterations in rice (<i>Oryza sativa</i> L.) seedlings under hexavalent chromium stress. <i>Planta</i> , 2014, 240, 291-308.	1.6	59

#	ARTICLE	IF	CITATIONS
73	Differences in physiological features associated with aluminum tolerance in Tibetan wild and cultivated barleys. <i>Plant Physiology and Biochemistry</i> , 2014, 75, 36-44.	2.8	14
74	The changes in physiological and biochemical traits of Tibetan wild and cultivated barley in response to low phosphorus stress. <i>Soil Science and Plant Nutrition</i> , 2014, 60, 832-842.	0.8	15
75	Development and Characterization of Polymorphic EST-SSR and Genomic SSR Markers for Tibetan Annual Wild Barley. <i>PLoS ONE</i> , 2014, 9, e94881.	1.1	42
76	Genotypic differences in physiological characteristics in the tolerance to drought and salinity combined stress between Tibetan wild and cultivated barley. <i>Plant Physiology and Biochemistry</i> , 2013, 63, 49-60.	2.8	219
77	Differential changes in grain ultrastructure, amylase, protein and amino acid profiles between Tibetan wild and cultivated barleys under drought and salinity alone and combined stress. <i>Food Chemistry</i> , 2013, 141, 2743-2750.	4.2	66
78	Genotypic and Environmental Variations of Arabinoxylan Content and Endoxylanase Activity in Barley Grains. <i>Journal of Integrative Agriculture</i> , 2013, 12, 1489-1494.	1.7	12
79	Alleviation of chromium toxicity in rice seedlings by applying exogenous glutathione. <i>Journal of Plant Physiology</i> , 2013, 170, 772-779.	1.6	67
80	Comparative study of alleviating effects of GSH, Se and Zn under combined contamination of cadmium and chromium in rice (<i>Oryza sativa</i>). <i>BioMetals</i> , 2013, 26, 297-308.	1.8	50
81	Tissue Metabolic Responses to Salt Stress in Wild and Cultivated Barley. <i>PLoS ONE</i> , 2013, 8, e55431.	1.1	186
82	Characteristics of Photosynthetic Performance, Antioxidant Capacity and Nutrient Concentration of Tibetan Wild Barley in Response to Aluminium Stress. <i>Asian Journal of Chemistry</i> , 2013, 25, 7727-7731.	0.1	2
83	Difference in Yield and Physiological Features in Response to Drought and Salinity Combined Stress during Anthesis in Tibetan Wild and Cultivated Barleys. <i>PLoS ONE</i> , 2013, 8, e77869.	1.1	116
84	Differences in Grain Ultrastructure, Phytochemical and Proteomic Profiles between the Two Contrasting Grain Cd-Accumulation Barley Genotypes. <i>PLoS ONE</i> , 2013, 8, e79158.	1.1	19
85	Comparative Proteomic Analysis of Aluminum Tolerance in Tibetan Wild and Cultivated Barleys. <i>PLoS ONE</i> , 2013, 8, e63428.	1.1	30
86	Selenium reduces cadmium uptake and mitigates cadmium toxicity in rice. <i>Journal of Hazardous Materials</i> , 2012, 235-236, 343-351.	6.5	259
87	Glutathione-Mediated Alleviation of Chromium Toxicity in Rice Plants. <i>Biological Trace Element Research</i> , 2012, 148, 255-263.	1.9	79
88	Alleviation of aluminum toxicity by hydrogen sulfide is related to elevated ATPase, and suppressed aluminum uptake and oxidative stress in barley. <i>Journal of Hazardous Materials</i> , 2012, 209-210, 121-128.	6.5	151
89	Subcellular distribution and chemical forms of chromium in rice plants suffering from different levels of chromium toxicity. <i>Journal of Plant Nutrition and Soil Science</i> , 2011, 174, 249-256.	1.1	89
90	Difference in response to aluminum stress among Tibetan wild barley genotypes. <i>Journal of Plant Nutrition and Soil Science</i> , 2011, 174, 952-960.	1.1	33

#	ARTICLE	IF	CITATIONS
91	Alleviation of Chromium Toxicity by Silicon Addition in Rice Plants. <i>Agricultural Sciences in China</i> , 2011, 10, 1188-1196.	0.6	70
92	Genotypic dependent effect of exogenous glutathione on Cd-induced changes in proteins, ultrastructure and antioxidant defense enzymes in rice seedlings. <i>Journal of Hazardous Materials</i> , 2011, 192, 1056-1066.	6.5	72
93	The ecotoxicological and interactive effects of chromium and aluminum on growth, oxidative damage and antioxidant enzymes on two barley genotypes differing in Al tolerance. <i>Environmental and Experimental Botany</i> , 2011, 70, 185-191.	2.0	84
94	The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants. <i>Environmental Pollution</i> , 2011, 159, 84-91.	3.7	970
95	Genotypic differences in callus induction and plant regeneration from mature embryos of barley (<i>Hordeum vulgare</i> L.). <i>Journal of Zhejiang University: Science B</i> , 2011, 12, 399-407.	1.3	20
96	Evaluation of salinity tolerance and analysis of allelic function of HvHKT1 and HvHKT2 in Tibetan wild barley. <i>Theoretical and Applied Genetics</i> , 2011, 122, 695-703.	1.8	123
97	Effects of Cadmium, Chromium and Lead on Growth, Metal Uptake and Antioxidative Capacity in <i>Typha angustifolia</i> . <i>Biological Trace Element Research</i> , 2011, 142, 77-92.	1.9	48
98	Modulation of Exogenous Glutathione in Phytochelatins and Photosynthetic Performance Against Cd Stress in the Two Rice Genotypes Differing in Cd Tolerance. <i>Biological Trace Element Research</i> , 2011, 143, 1159-1173.	1.9	76
99	Modulation of Exogenous Glutathione in Ultrastructure and Photosynthetic Performance Against Cd Stress in the Two Barley Genotypes Differing in Cd Tolerance. <i>Biological Trace Element Research</i> , 2011, 144, 1275-1288.	1.9	78
100	Interactive effects of aluminum and chromium stresses on the uptake of nutrients and the metals in barley. <i>Soil Science and Plant Nutrition</i> , 2011, 57, 68-79.	0.8	17
101	Identification of QTLs for yield and yield components of barley under different growth conditions. <i>Journal of Zhejiang University: Science B</i> , 2010, 11, 169-176.	1.3	58
102	Genotype-Dependent Effect of Exogenous Nitric Oxide on Cd-induced Changes in Antioxidative Metabolism, Ultrastructure, and Photosynthetic Performance in Barley Seedlings (<i>Hordeum vulgare</i>). <i>Journal of Plant Growth Regulation</i> , 2010, 29, 394-408.	2.8	88
103	Comparative proteomic analysis of <i>Typha angustifolia</i> leaf under chromium, cadmium and lead stress. <i>Journal of Hazardous Materials</i> , 2010, 184, 191-203.	6.5	72
104	Difference in response to drought stress among Tibet wild barley genotypes. <i>Euphytica</i> , 2010, 172, 395-403.	0.6	101
105	Modulation of exogenous glutathione in antioxidant defense system against Cd stress in the two barley genotypes differing in Cd tolerance. <i>Plant Physiology and Biochemistry</i> , 2010, 48, 663-672.	2.8	249
106	Effects of chromium stress on the subcellular distribution and chemical form of Ca, Mg, Fe, and Zn in two rice genotypes. <i>Journal of Plant Nutrition and Soil Science</i> , 2010, 173, 135-148.	1.1	27
107	Hormonal changes in grains of cv. Triumph and its mutant TL43 as affected by nitrogen fertilizer at heading time. <i>Journal of Cereal Science</i> , 2009, 49, 246-249.	1.8	6
108	The effect of H ₂ O ₂ and abscisic acid (ABA) interaction on α -amylase activity under osmotic stress during grain development in barley. <i>Plant Physiology and Biochemistry</i> , 2009, 47, 778-784.	2.8	16

#	ARTICLE	IF	CITATIONS
109	Genetic mapping of quantitative trait loci associated with α -amylase and limit dextrinase activities and β -glucan and protein fraction contents in barley. <i>Journal of Zhejiang University: Science B</i> , 2009, 10, 839-846.	1.3	13
110	The Variation of α -amylase Activity and Protein Fractions in Barley Grains as Affected by Genotypes and Post-anthesis Temperatures. <i>Journal of the Institute of Brewing</i> , 2009, 115, 208-213.	0.8	21
111	Comparison of EDTA- and Citric Acid-Enhanced Phytoextraction of Heavy Metals in Artificially Metal Contaminated Soil by <i>Typha Angustifolia</i> . <i>International Journal of Phytoremediation</i> , 2009, 11, 558-574.	1.7	90
112	Identification of Barley Varieties Tolerant to Cadmium Toxicity. <i>Biological Trace Element Research</i> , 2008, 121, 171-179.	1.9	43
113	Genotypic and environmental variation in chromium, cadmium and lead concentrations in rice. <i>Environmental Pollution</i> , 2008, 153, 309-314.	3.7	154
114	Changes of organic acid exudation and rhizosphere pH in rice plants under chromium stress. <i>Environmental Pollution</i> , 2008, 155, 284-289.	3.7	131
115	Identification of barley genotypes with low grain Cd accumulation and its interaction with four microelements. <i>Chemosphere</i> , 2007, 67, 2082-2088.	4.2	102
116	Differences in yield components and kernel Cd accumulation in response to Cd toxicity in four barley genotypes. <i>Chemosphere</i> , 2007, 70, 83-92.	4.2	64
117	Characterization of Leaf Photosynthetic Properties for No-Tillage Rice. <i>Rice Science</i> , 2007, 14, 283-288.	1.7	23
118	Influence of Aluminum and Cadmium Stresses on Mineral Nutrition and Root Exudates in Two Barley Cultivars. <i>Pedosphere</i> , 2007, 17, 505-512.	2.1	63
119	A Chromium-Tolerant Plant Growing in Cr-Contaminated Land. <i>International Journal of Phytoremediation</i> , 2007, 9, 167-179.	1.7	49
120	Differences in Mn uptake and subcellular distribution in different barley genotypes as a response to Cd toxicity. <i>Science of the Total Environment</i> , 2007, 385, 228-234.	3.9	23
121	Cadmium translocation and accumulation in developing barley grains. <i>Planta</i> , 2007, 227, 223-232.	1.6	54
122	Genotypic Difference in the Responses of Seedling Growth and Cd Toxicity in Rice (<i>Oryza sativa</i> L.). <i>Agricultural Sciences in China</i> , 2006, 5, 68-76.	0.6	43
123	Differences in Physiological Traits Among Salt-Stressed Barley Genotypes. <i>Communications in Soil Science and Plant Analysis</i> , 2006, 37, 557-570.	0.6	29
124	Influence of cadmium on antioxidant capacity and four microelement concentrations in tomato seedlings (<i>Lycopersicon esculentum</i>). <i>Chemosphere</i> , 2006, 64, 1659-1666.	4.2	244
125	Heterosis for Yield and some Physiological Traits in Hybrid Cotton Cikangza 1. <i>Euphytica</i> , 2006, 151, 71-77.	0.6	16
126	Interaction of Salinity and Cadmium Stresses on Antioxidant Enzymes, Sodium, and Cadmium Accumulation in Four Barley Genotypes. <i>Journal of Plant Nutrition</i> , 2006, 29, 2215-2225.	0.9	16

#	ARTICLE	IF	CITATIONS
127	Heterosis in CMS hybrids of cotton for photosynthetic and chlorophyll fluorescence parameters. <i>Euphytica</i> , 2005, 144, 353-361.	0.6	13
128	Effect of cadmium on growth and photosynthesis of tomato seedlings. <i>Journal of Zhejiang University Science B</i> , 2005, 6B, 974-980.	0.4	91
129	Zinc alleviates growth inhibition and oxidative stress caused by cadmium in rice. <i>Journal of Plant Nutrition and Soil Science</i> , 2005, 168, 255-261.	1.1	121
130	Response of Cadmium Uptake in Different Barley Genotypes to Cadmium Level. <i>Journal of Plant Nutrition</i> , 2005, 28, 2201-2209.	0.9	13
131	Subcellular distribution and chemical form of Cd and Cd-Zn interaction in different barley genotypes. <i>Chemosphere</i> , 2005, 60, 1437-1446.	4.2	228
132	Effect of Cadmium on Uptake and Translocation of Three Microelements in Cotton. <i>Journal of Plant Nutrition</i> , 2005, 27, 2019-2032.	0.9	14
133	Protein and hordein fraction content in barley seeds as affected by sowing date and their relations to malting quality. <i>Journal of Zhejiang University Science B</i> , 2005, 6B, 1069-1075.	0.4	34
134	Differences in growth and yield in response to cadmium toxicity in cotton genotypes. <i>Journal of Plant Nutrition and Soil Science</i> , 2004, 167, 85-90.	1.1	36
135	Genotypic Difference in Effect of Cadmium on Development and Mineral Concentrations of Cotton. <i>Communications in Soil Science and Plant Analysis</i> , 2004, 35, 285-299.	0.6	17
136	Effects of aluminum and cadmium toxicity on growth and antioxidant enzyme activities of two barley genotypes with different Al resistance. <i>Plant and Soil</i> , 2004, 258, 241-248.	1.8	157
137	The changes of β -glucan content and β -glucanase activity in barley before and after malting and their relationships to malt qualities. <i>Food Chemistry</i> , 2004, 86, 223-228.	4.2	87
138	Effect of cadmium on free amino acid, glutathione and ascorbic acid concentrations in two barley genotypes (<i>Hordeum vulgare</i> L.) differing in cadmium tolerance. <i>Chemosphere</i> , 2004, 57, 447-454.	4.2	107
139	Four barley genotypes respond differently to cadmium: lipid peroxidation and activities of antioxidant capacity. <i>Environmental and Experimental Botany</i> , 2003, 50, 67-78.	2.0	322
140	Genotypic and environmental variation in barley beta-amylase activity and its relation to protein content. <i>Food Chemistry</i> , 2003, 83, 163-165.	4.2	16
141	Interaction of Cadmium and Four Microelements for Uptake and Translocation in Different Barley Genotypes. <i>Communications in Soil Science and Plant Analysis</i> , 2003, 34, 2003-2020.	0.6	53
142	Genotypic Differences in Effect of Cadmium on Growth Parameters of Barley During Ontogenesis. <i>Communications in Soil Science and Plant Analysis</i> , 2003, 34, 2021-2034.	0.6	8
143	GENOTYPIC VARIATION IN KERNEL HEAVY METAL CONCENTRATIONS IN BARLEY AND AS AFFECTED BY SOIL FACTORS. <i>Journal of Plant Nutrition</i> , 2002, 25, 1163-1173.	0.9	64
144	ALLEVIATION OF CADMIUM-TOXICITY BY APPLICATION OF ZINC AND ASCORBIC ACID IN BARLEY. <i>Journal of Plant Nutrition</i> , 2002, 25, 2745-2761.	0.9	64

#	ARTICLE	IF	CITATIONS
145	Effect of Cadmium on Uptake and Translocation of Three Microelements in Cotton. , 0, .		1