

# Satomi Kobayashi

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/1820249/satomi-kobayashi-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

33  
papers

1,391  
citations

14  
h-index

37  
g-index

37  
ext. papers

1,802  
ext. citations

5.8  
avg, IF

4.21  
L-index

#	Paper	IF	Citations
33	Nutrient and Food Group Prediction as Orchestrated by an Automated Image Recognition System in a Smartphone App (CALO mama): Validation Study.. <i>JMIR Formative Research</i> , <b>2022</b> , 6, e31875	2.5	1
32	Identifying the most influential gene expression profile in distinguishing ANCA-associated vasculitis from healthy controls. <i>Journal of Autoimmunity</i> , <b>2021</b> , 119, 102617	15.5	1
31	Dynamic landscape of immune cell-specific gene regulation in immune-mediated diseases. <i>Cell</i> , <b>2021</b> , 184, 3006-3021.e17	56.2	28
30	Associations of Education With Overall Diet Quality Are Explained by Different Food Groups in Middle-aged and Old Japanese Women. <i>Journal of Epidemiology</i> , <b>2021</b> , 31, 280-286	3.4	0
29	Integrated bulk and single-cell RNA-sequencing identified disease-relevant monocytes and a gene network module underlying systemic sclerosis. <i>Journal of Autoimmunity</i> , <b>2021</b> , 116, 102547	15.5	7
28	Serum Amphiregulin and Heparin-Binding Epidermal Growth Factor as Biomarkers in Patients with Idiopathic Inflammatory Myopathy. <i>Journal of Clinical Medicine</i> , <b>2021</b> , 10,	5.1	1
27	Intake of Coffee Associated With Decreased Depressive Symptoms Among Elderly Japanese Women: A Multi-Center Cross-Sectional Study. <i>Journal of Epidemiology</i> , <b>2020</b> , 30, 338-344	3.4	6
26	Association of alcohol consumption with prevalence of fatty liver after adjustment for dietary patterns: Cross-sectional analysis of Japanese middle-aged adults. <i>Clinical Nutrition</i> , <b>2020</b> , 39, 1580-1586	5.9	1
25	Current status of education and research on public health nutrition in Japan: comparison with South Korea, Taiwan, and mainland China. <i>BMC Nutrition</i> , <b>2019</b> , 5, 10	2.5	1
24	No association between fruits or vegetables and non-alcoholic fatty liver disease in middle-aged men and women. <i>Nutrition</i> , <b>2019</b> , 61, 119-124	4.8	17
23	Relative validity of brief-type self-administered diet history questionnaire among very old Japanese aged 80 years or older. <i>Public Health Nutrition</i> , <b>2019</b> , 22, 212-222	3.3	17
22	Dietary non-enzymatic antioxidant capacity and DNA damage in a working population. <i>Nutrition</i> , <b>2018</b> , 47, 63-68	4.8	6
21	Dietary phosphorus intake estimated by 4-day dietary records and two 24-hour urine collections and their associated factors in Japanese adults. <i>European Journal of Clinical Nutrition</i> , <b>2018</b> , 72, 517-525	5.2	8
20	Association between habitual tryptophan intake and depressive symptoms in young and middle-aged women. <i>Journal of Affective Disorders</i> , <b>2018</b> , 231, 44-50	6.6	8
19	Higher dietary acid load is associated with a higher prevalence of frailty, particularly slowness/weakness and low physical activity, in elderly Japanese women. <i>European Journal of Nutrition</i> , <b>2018</b> , 57, 1639-1650	5.2	7
18	Higher dietary glycemic index, but not glycemic load, is associated with a lower prevalence of depressive symptoms in a cross-sectional study of young and middle-aged Japanese women. <i>European Journal of Nutrition</i> , <b>2018</b> , 57, 2261-2273	5.2	7
17	Urine 24-Hour Sodium Excretion Decreased between 1953 and 2014 in Japan, but Estimated Intake Still Exceeds the WHO Recommendation. <i>Journal of Nutrition</i> , <b>2017</b> , 147, 390-397	4.1	10

16	Living status and frequency of eating out-of-home foods in relation to nutritional adequacy in 4,017 Japanese female dietetic students aged 18-20 years: A multicenter cross-sectional study. <i>Journal of Epidemiology</i> , <b>2017</b> , 27, 287-293	3.4	18
15	Two cases of very elderly onset male lupus patient; the characteristics and sex differences of elderly onset systemic lupus erythematosus patients. <i>Modern Rheumatology Case Reports</i> , <b>2017</b> , 1, 84-88 <sup>0.4</sup>		
14	Effect of excess iodine intake on thyroid diseases in different populations: A systematic review and meta-analyses including observational studies. <i>PLoS ONE</i> , <b>2017</b> , 12, e0173722	3.7	55
13	No Association of Caffeinated Beverage or Caffeine Intake with Prevalence of Urinary Incontinence Among Middle-Aged Japanese Women: A Multicenter Cross-Sectional Study. <i>Journal of Womens Health</i> , <b>2017</b> , 26, 860-869	3	1
12	Food-based diet quality score in relation to depressive symptoms in young and middle-aged Japanese women. <i>British Journal of Nutrition</i> , <b>2017</b> , 117, 1674-1681	3.6	21
11	Nutritional correlates of monetary diet cost in young, middle-aged and older Japanese women. <i>Journal of Nutritional Science</i> , <b>2017</b> , 6, e22	2.7	2
10	Diet with a combination of high protein and high total antioxidant capacity is strongly associated with low prevalence of frailty among old Japanese women: a multicenter cross-sectional study. <i>Nutrition Journal</i> , <b>2017</b> , 16, 29	4.3	30
9	Association between rice, bread, and noodle intake and the prevalence of non-alcoholic fatty liver disease in Japanese middle-aged men and women. <i>Clinical Nutrition</i> , <b>2017</b> , 36, 1601-1608	5.9	20
8	A Japanese diet with low glycaemic index and glycaemic load is associated with both favourable and unfavourable aspects of dietary intake patterns in three generations of women. <i>Public Health Nutrition</i> , <b>2017</b> , 20, 649-659	3.3	9
7	Factors Influencing Exclusive Breastfeeding in Early Infancy: A Prospective Study in North Central Nigeria. <i>Maternal and Child Health Journal</i> , <b>2016</b> , 20, 363-75	2.4	16
6	Development of a food-based diet quality score for Japanese: associations of the score with nutrient intakes in young, middle-aged and older Japanese women. <i>Journal of Nutritional Science</i> , <b>2016</b> , 5, e41	2.7	15
5	Cohabitational effect of grandparents on dietary intake among young Japanese women and their mothers living together. A multicenter cross-sectional study. <i>Appetite</i> , <b>2015</b> , 91, 287-97	4.5	8
4	High protein intake is associated with low prevalence of frailty among old Japanese women: a multicenter cross-sectional study. <i>Nutrition Journal</i> , <b>2013</b> , 12, 164	4.3	112
3	Both comprehensive and brief self-administered diet history questionnaires satisfactorily rank nutrient intakes in Japanese adults. <i>Journal of Epidemiology</i> , <b>2012</b> , 22, 151-9	3.4	455
2	Dietary total antioxidant capacity from different assays in relation to serum C-reactive protein among young Japanese women. <i>Nutrition Journal</i> , <b>2012</b> , 11, 91	4.3	36
1	Comparison of relative validity of food group intakes estimated by comprehensive and brief-type self-administered diet history questionnaires against 16 d dietary records in Japanese adults. <i>Public Health Nutrition</i> , <b>2011</b> , 14, 1200-11	3.3	467