

# Ikuko Hara-Nishimura

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1827563/publications.pdf>

Version: 2024-02-01

198  
papers

14,311  
citations

12330

69  
h-index

24982

109  
g-index

205  
all docs

205  
docs citations

205  
times ranked

10694  
citing authors

#	ARTICLE	IF	CITATIONS
1	A rich and bountiful harvest: Key discoveries in plant cell biology. <i>Plant Cell</i> , 2022, 34, 53-71.	6.6	7
2	Aerial (+)-borneol modulates root morphology, auxin signalling and meristematic activity in <i>Arabidopsis</i> roots. <i>Biology Letters</i> , 2022, 18, 20210629.	2.3	2
3	Plant ESCRT protein ALIX coordinates with retromer complex in regulating receptor-mediated sorting of soluble vacuolar proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, e2200492119.	7.1	12
4	The <i>Arabidopsis</i> transcription factor NAI1 activates the <i>NAI2</i> promoter by binding to the G-box motifs. <i>Plant Signaling and Behavior</i> , 2021, 16, 1846928.	2.4	3
5	Excess sterol accumulation affects seed morphology and physiology in <i>Arabidopsis thaliana</i> . <i>Plant Signaling and Behavior</i> , 2021, 16, 1872217.	2.4	10
6	In vitro assembly of nuclear envelope in tobacco cultured cells. <i>Nucleus</i> , 2021, 12, 82-89.	2.2	0
7	Regulation and Physiological Significance of the Nuclear Shape in Plants. <i>Frontiers in Plant Science</i> , 2021, 12, 673905.	3.6	7
8	Galactoglucomannan structure of <i>Arabidopsis</i> seed coat mucilage in <i>GDP</i> mannose synthesis impaired mutants. <i>Physiologia Plantarum</i> , 2021, 173, 1244-1252.	5.2	9
9	<i>Arabidopsis</i> FLYING SAUCER 2 Functions Redundantly with FLY1 to Establish Normal Seed Coat Mucilage. <i>Plant and Cell Physiology</i> , 2020, 61, 308-317.	3.1	9
10	Vacuolar processing enzymes in the plant life cycle. <i>New Phytologist</i> , 2020, 226, 21-31.	7.3	51
11	NAI2 and TSA1 Drive Differentiation of Constitutive and Inducible ER Body Formation in Brassicaceae. <i>Plant and Cell Physiology</i> , 2020, 61, 722-734.	3.1	8
12	Higher Stomatal Density Improves Photosynthetic Induction and Biomass Production in <i>Arabidopsis</i> Under Fluctuating Light. <i>Frontiers in Plant Science</i> , 2020, 11, 589603.	3.6	69
13	Dynamic Capture and Release of Endoplasmic Reticulum Exit Sites by Golgi Stacks in <i>Arabidopsis</i> . <i>IScience</i> , 2020, 23, 101265.	4.1	11
14	Generation of <i>Arabidopsis</i> lines with a red fluorescent marker for endoplasmic reticulum using a tail-anchored protein cytochrome b5-B. <i>Plant Signaling and Behavior</i> , 2020, 15, 1790196.	2.4	3
15	Subnuclear gene positioning through lamina association affects copper tolerance. <i>Nature Communications</i> , 2020, 11, 5914.	12.8	37
16	The nuclear envelope protein KAKU4 determines the migration order of the vegetative nucleus and sperm cells in pollen tubes. <i>Journal of Experimental Botany</i> , 2020, 71, 6273-6281.	4.8	20
17	Characterization of rhizome transcriptome and identification of a rhizomatous ER body in the clonal plant <i>Cardamine leucantha</i> . <i>Scientific Reports</i> , 2020, 10, 13291.	3.3	4
18	Excess sterols disrupt plant cellular activity by inducing stress-responsive gene expression. <i>Journal of Plant Research</i> , 2020, 133, 383-392.	2.4	8

#	ARTICLE	IF	CITATIONS
19	Arabidopsis ECHIDNA protein is involved in seed coloration, protein trafficking to vacuoles, and vacuolar biogenesis. <i>Journal of Experimental Botany</i> , 2020, 71, 3999-4009.	4.8	10
20	Endoplasmic reticulum-derived bodies enable a single-cell chemical defense in Brassicaceae plants. <i>Communications Biology</i> , 2020, 3, 21.	4.4	26
21	A space-saving visual screening method, <i>Glycine max</i> FAST, for generating transgenic soybean. <i>Plant Signaling and Behavior</i> , 2020, 15, 1722911.	2.4	1
22	HIGH STEROL ESTER 1 is a key factor in plant sterol homeostasis. <i>Nature Plants</i> , 2019, 5, 1154-1166.	9.3	26
23	How to Investigate the Role of the Actin-Myosin Cytoskeleton in Organ Straightening. <i>Methods in Molecular Biology</i> , 2019, 1924, 215-221.	0.9	0
24	Polar Localization of the Borate Exporter BOR1 Requires AP2-Dependent Endocytosis. <i>Plant Physiology</i> , 2019, 179, 1569-1580.	4.8	58
25	Leaf Endoplasmic Reticulum Bodies Identified in Arabidopsis Rosette Leaves Are Involved in Defense against Herbivory. <i>Plant Physiology</i> , 2019, 179, 1515-1524.	4.8	58
26	A Genotypic Comparison Reveals That the Improvement in Nitrogen Remobilization Efficiency in Oilseed Rape Leaves Is Related to Specific Patterns of Senescence-Associated Protease Activities and Phytohormones. <i>Frontiers in Plant Science</i> , 2019, 10, 46.	3.6	13
27	Biogenesis of leaf endoplasmic reticulum body is regulated by both jasmonate-dependent and independent pathways. <i>Plant Signaling and Behavior</i> , 2019, 14, 1622982.	2.4	6
28	tRNA Wobble Modification Affects Leaf Cell Development in Arabidopsis thaliana. <i>Plant and Cell Physiology</i> , 2019, 60, 2026-2039.	3.1	14
29	Comprehensive nuclear proteome of Arabidopsis obtained by sequential extraction. <i>Nucleus</i> , 2019, 10, 81-92.	2.2	28
30	Identification of Periplasmic Root-Cap Mucilage in Developing Columella Cells of Arabidopsis thaliana. <i>Plant and Cell Physiology</i> , 2019, 60, 1296-1303.	3.1	13
31	Sucrose Starvation Induces Microautophagy in Plant Root Cells. <i>Frontiers in Plant Science</i> , 2019, 10, 1604.	3.6	27
32	ANGUSTIFOLIA Regulates Actin Filament Alignment for Nuclear Positioning in Leaves. <i>Plant Physiology</i> , 2019, 179, 233-247.	4.8	18
33	Measurement of the Caspase-1-Like Activity of Vacuolar Processing Enzyme in Plants. <i>Methods in Molecular Biology</i> , 2018, 1743, 163-171.	0.9	3
34	Plant Vacuoles. <i>Annual Review of Plant Biology</i> , 2018, 69, 123-145.	18.7	94
35	Involvement of Adapter Protein Complex 4 in Hypersensitive Cell Death Induced by Avirulent Bacteria. <i>Plant Physiology</i> , 2018, 176, 1824-1834.	4.8	25
36	Membrane Dynamics and Multiple Functions of Oil Bodies in Seeds and Leaves. <i>Plant Physiology</i> , 2018, 176, 199-207.	4.8	73

#	ARTICLE	IF	CITATIONS
37	Tissue-specific and intracellular localization of indican synthase from <i>Polygonum tinctorium</i> . <i>Plant Physiology and Biochemistry</i> , 2018, 132, 138-144.	5.8	11
38	GREEN FLUORESCENT SEED, to Evaluate Vacuolar Trafficking in Arabidopsis Seeds. <i>Methods in Molecular Biology</i> , 2018, 1789, 1-7.	0.9	1
39	Endoplasmic Reticulum (ER) Membrane Proteins (LUNAPARKs) are Required for Proper Configuration of the Cortical ER Network in Plant Cells. <i>Plant and Cell Physiology</i> , 2018, 59, 1931-1941.	3.1	8
40	Pectin RG-I rhamnosyltransferases represent a novel plant-specific glycosyltransferase family. <i>Nature Plants</i> , 2018, 4, 669-676.	9.3	111
41	Specialized Vacuoles of Myrosin Cells: Chemical Defense Strategy in Brassicales Plants. <i>Plant and Cell Physiology</i> , 2018, 59, 1309-1316.	3.1	54
42	Stress granule formation is induced by a threshold temperature rather than a temperature difference in Arabidopsis. <i>Journal of Cell Science</i> , 2018, 131, .	2.0	27
43	The AP-1 Complex is Required for Proper Mucilage Formation in Arabidopsis Seeds. <i>Plant and Cell Physiology</i> , 2018, 59, 2331-2338.	3.1	15
44	Synaptotagmin-Associated Endoplasmic Reticulum-Plasma Membrane Contact Sites Are Localized to Immobile ER Tubules. <i>Plant Physiology</i> , 2018, 178, 641-653.	4.8	27
45	The Multifaceted Roles of Plant Vacuoles. <i>Plant and Cell Physiology</i> , 2018, 59, 1285-1287.	3.1	8
46	Nup82 functions redundantly with Nup136 in a salicylic acid-dependent defense response of <i>Arabidopsis thaliana</i> . <i>Nucleus</i> , 2017, 8, 301-311.	2.2	16
47	Inhibition of cell polarity establishment in stomatal asymmetric cell division using the chemical compound bubblin. <i>Development (Cambridge)</i> , 2017, 144, 499-506.	2.5	11
48	HSP90 stabilizes auxin receptor TIR1 and ensures plasticity of auxin responses. <i>Plant Signaling and Behavior</i> , 2017, 12, e1311439.	2.4	16
49	Polar Localization of the NIP5;1 Boric Acid Channel Is Maintained by Endocytosis and Facilitates Boron Transport in Arabidopsis Roots. <i>Plant Cell</i> , 2017, 29, 824-842.	6.6	107
50	Isolation of Protein Storage Vacuoles and Their Membranes. <i>Methods in Molecular Biology</i> , 2017, 1511, 163-168.	0.9	0
51	PYK10 myrosinase reveals a functional coordination between endoplasmic reticulum bodies and glucosinolates in <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 2017, 89, 204-220.	5.7	128
52	An efficient <i>Agrobacterium</i> -mediated transformation method for switchgrass genotypes using Type I callus. <i>Plant Biotechnology</i> , 2016, 33, 19-26.	1.0	4
53	FAMA: A Molecular Link between Stomata and Myrosin Cells. <i>Trends in Plant Science</i> , 2016, 21, 861-871.	8.8	24
54	The $\frac{1}{4}$ Subunit of <i>Arabidopsis</i> Adaptor Protein-2 Is Involved in Effector-Triggered Immunity Mediated by Membrane-Localized Resistance Proteins. <i>Molecular Plant-Microbe Interactions</i> , 2016, 29, 345-351.	2.6	24

#	ARTICLE	IF	CITATIONS
55	HSP90 Stabilizes Auxin-Responsive Phenotypes by Masking a Mutation in the Auxin Receptor TIR1. <i>Plant and Cell Physiology</i> , 2016, 57, 2245-2254.	3.1	28
56	Plant Nuclei Move to Escape Ultraviolet-Induced DNA Damage and Cell Death. <i>Plant Physiology</i> , 2016, 170, 678-685.	4.8	22
57	The Adaptor Complex AP-4 Regulates Vacuolar Protein Sorting at the trans-Golgi Network by Interacting with VACUOLAR SORTING RECEPTOR1. <i>Plant Physiology</i> , 2016, 170, 211-219.	4.8	72
58	Phosphorylation of the C Terminus of RHD3 Has a Critical Role in Homotypic ER Membrane Fusion in Arabidopsis. <i>Plant Physiology</i> , 2016, 170, 867-880.	4.8	31
59	Myrosin cells are differentiated directly from ground meristem cells and are developmentally independent of the vasculature in Arabidopsis leaves. <i>Plant Signaling and Behavior</i> , 2016, 11, e1150403.	2.4	13
60	An ABC transporter B family protein, ABCB19, is required for cytoplasmic streaming and gravitropism of the inflorescence stems. <i>Plant Signaling and Behavior</i> , 2016, 11, e1010947.	2.4	21
61	Decreased Expression of a Gene Caused by a T-DNA Insertion in an Adjacent Gene in Arabidopsis. <i>PLoS ONE</i> , 2016, 11, e0147911.	2.5	5
62	A directionâ€selective localâ€thresholding method, <scp>DSL</scp>, in combination with a dyeâ€based method for automated threeâ€dimensional segmentation of cells and airspaces in developing leaves. <i>Plant Journal</i> , 2015, 81, 357-366.	5.7	15
63	Vacuolar processing enzyme in plant programmed cell death. <i>Frontiers in Plant Science</i> , 2015, 6, 234.	3.6	182
64	Methyl Jasmonate Affects Morphology, Number and Activity of Endoplasmic Reticulum Bodies in <i>Raphanus sativus</i> Root Cells. <i>Plant and Cell Physiology</i> , 2015, 56, 61-72.	3.1	14
65	Retromer Contributes to Immunity-Associated Cell Death in Arabidopsis. <i>Plant Cell</i> , 2015, 27, 463-479.	6.6	67
66	Recent advances in understanding plant nuclear envelope proteins involved in nuclear morphology. <i>Journal of Experimental Botany</i> , 2015, 66, 1641-1647.	4.8	28
67	Leaf oil bodies are subcellular factories producing antifungal oxylipins. <i>Current Opinion in Plant Biology</i> , 2015, 25, 145-150.	7.1	40
68	BEACH-Domain Proteins Act Together in a Cascade to Mediate Vacuolar Protein Trafficking and Disease Resistance in Arabidopsis. <i>Molecular Plant</i> , 2015, 8, 389-398.	8.3	27
69	Regulation of organ straightening and plant posture by an actinâ€myosin XI cytoskeleton. <i>Nature Plants</i> , 2015, 1, 15031.	9.3	60
70	Oil body-mediated defense against fungi: From tissues to ecology. <i>Plant Signaling and Behavior</i> , 2015, 10, e989036.	2.4	27
71	Functions of plant-specific myosin XI: from intracellular motility to plant postures. <i>Current Opinion in Plant Biology</i> , 2015, 28, 30-38.	7.1	44
72	Effects of stomatal density and leaf water content on the <sup>18</sup><scp>O</scp> enrichment of leaf water. <i>New Phytologist</i> , 2015, 206, 141-151.	7.3	21

#	ARTICLE	IF	CITATIONS
73	Nucleoporin 75 Is Involved in the Ethylene-Mediated Production of Phytoalexin for the Resistance of <i>Nicotiana benthamiana</i> to <i>Phytophthora infestans</i> . <i>Molecular Plant-Microbe Interactions</i> , 2014, 27, 1318-1330.	2.6	27
74	Functional insights of nucleocytoplasmic transport in plants. <i>Frontiers in Plant Science</i> , 2014, 5, 118.	3.6	50
75	ER bodies in plants of the Brassicales order: biogenesis and association with innate immunity. <i>Frontiers in Plant Science</i> , 2014, 5, 73.	3.6	93
76	Myosin Cell Development Is Regulated by Endocytosis Machinery and PIN1 Polarity in Leaf Primordia of <i>Arabidopsis thaliana</i> . <i>Plant Cell</i> , 2014, 26, 4448-4461.	6.6	12
77	Microtubules Contribute to Tubule Elongation and Anchoring of Endoplasmic Reticulum, Resulting in High Network Complexity in <i>Arabidopsis</i> . <i>Plant Physiology</i> , 2014, 166, 1869-1876.	4.8	55
78	GFS9 contributes to intracellular membrane trafficking and flavonoid accumulation in <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 2014, 80, 410-423.	5.7	63
79	CONTINUOUS VASCULAR RING (COV1) is a trans-Golgi Network-Localized Membrane Protein Required for Golgi Morphology and Vacuolar Protein Sorting. <i>Plant and Cell Physiology</i> , 2014, 55, 764-772.	3.1	32
80	The Novel Nuclear Envelope Protein KAKU4 Modulates Nuclear Morphology in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2014, 26, 2143-2155.	6.6	81
81	CRISPR/Cas9-Mediated Targeted Mutagenesis in the Liverwort <i>Marchantia polymorpha</i> L.. <i>Plant and Cell Physiology</i> , 2014, 55, 475-481.	3.1	262
82	Leaf Oil Body Functions as a Subcellular Factory for the Production of a Phytoalexin in <i>Arabidopsis</i> . <i>Plant Physiology</i> , 2014, 164, 105-118.	4.8	98
83	<i>Arabidopsis</i> mutants affecting oxylipin signaling in photo-oxidative stress responses. <i>Plant Physiology and Biochemistry</i> , 2014, 81, 90-95.	5.8	16
84	FAMA Is an Essential Component for the Differentiation of Two Distinct Cell Types, Myosin Cells and Guard Cells, in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2014, 26, 4039-4052.	6.6	50
85	Seed storage albumins: biosynthesis, trafficking and structures. <i>Functional Plant Biology</i> , 2014, 41, 671.	2.1	37
86	Evaluation of Defective Endosomal Trafficking to the Vacuole by Monitoring Seed Storage Proteins in <i>Arabidopsis thaliana</i> . <i>Methods in Molecular Biology</i> , 2014, 1209, 131-142.	0.9	2
87	Myosin XI-i Links the Nuclear Membrane to the Cytoskeleton to Control Nuclear Movement and Shape in <i>Arabidopsis</i> . <i>Current Biology</i> , 2013, 23, 1776-1781.	3.9	193
88	The molecular architecture of the plant nuclear pore complex. <i>Journal of Experimental Botany</i> , 2013, 64, 823-832.	4.8	78
89	MAIGO5 Functions in Protein Export from Golgi-Associated Endoplasmic Reticulum Exit Sites in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2013, 25, 4658-4675.	6.6	53
90	Trafficking of Vacuolar Proteins: The Crucial Role of <i>Arabidopsis</i> Vacuolar Protein Sorting 29 in Recycling Vacuolar Sorting Receptor. <i>Plant Cell</i> , 2013, 24, 5058-5073.	6.6	41

#	ARTICLE	IF	CITATIONS
91	The AP-1 $\mu$ Adaptin is Required for KNOLLE Localization at the Cell Plate to Mediate Cytokinesis in <i>Arabidopsis</i> . <i>Plant and Cell Physiology</i> , 2013, 54, 838-847.	3.1	79
92	Spatiotemporal Secretion of PEROXIDASE36 Is Required for Seed Coat Mucilage Extrusion in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2013, 25, 1355-1367.	6.6	85
93	<sc>MAG</sc>2 and three <sc>MAG</sc>2<sc>INTERACTING PROTEIN</sc>s form an <sc>ER</sc>-localized complex to facilitate storage protein transport in <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 2013, 76, 781-791.	5.7	34
94	Enhancement of leaf photosynthetic capacity through increased stomatal density in <i>Arabidopsis</i> . <i>New Phytologist</i> , 2013, 198, 757-764.	7.3	223
95	Identification and Dynamics of <i>Arabidopsis</i> Adaptor Protein-2 Complex and Its Involvement in Floral Organ Development. <i>Plant Cell</i> , 2013, 25, 2958-2969.	6.6	121
96	Stomagen/EPFL9. , 2013, , 67-70.		0
97	Plant Legumain, Asparaginyl Endopeptidase, Vacuolar Processing Enzyme. , 2013, , 2314-2320.		2
98	Identification of Two Novel Endoplasmic Reticulum Body-Specific Integral Membrane Proteins $\hat{\hat{A}}$ . <i>Plant Physiology</i> , 2012, 161, 108-120.	4.8	51
99	ERMO3/MVP1/GOLD36 Is Involved in a Cell Type-Specific Mechanism for Maintaining ER Morphology in <i>Arabidopsis thaliana</i> . <i>PLoS ONE</i> , 2012, 7, e49103.	2.5	22
100	Positive and negative peptide signals control stomatal density. <i>Cellular and Molecular Life Sciences</i> , 2011, 68, 2081-2088.	5.4	63
101	Identification and Characterization of Nuclear Pore Complex Components in <i>Arabidopsis thaliana</i> . <i>Plant Cell</i> , 2011, 22, 4084-4097.	6.6	256
102	A non-destructive screenable marker, OsFAST, for identifying transgenic rice seeds. <i>Plant Signaling and Behavior</i> , 2011, 6, 1454-1456.	2.4	9
103	Involvement of the nuclear pore complex in morphology of the plant nucleus. <i>Nucleus</i> , 2011, 2, 168-172.	2.2	63
104	Unique Defense Strategy by the Endoplasmic Reticulum Body in Plants. <i>Plant and Cell Physiology</i> , 2011, 52, 2039-2049.	3.1	76
105	Myosin XI-Dependent Formation of Tubular Structures from Endoplasmic Reticulum Isolated from Tobacco Cultured BY-2 Cells $\hat{\hat{A}}$ . <i>Plant Physiology</i> , 2011, 156, 129-143.	4.8	46
106	Oil-Body-Membrane Proteins and Their Physiological Functions in Plants. <i>Biological and Pharmaceutical Bulletin</i> , 2010, 33, 360-363.	1.4	102
107	æç%©ã®ç°èfžæ»ã,'áã,<é*ã¼ã&ãšã½ ©ã*ãf†ã,1ãf—ãfãf†ã,çãf¼ã,¼ã®ã½1ã%²ã,'æžçã,ç. <i>Kagaku To Seibutsu</i> , 2010, 48, 734-736		
108	A rapid and non-destructive screenable marker, FAST, for identifying transformed seeds of <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 2010, 61, 519-528.	5.7	325

#	ARTICLE	IF	CITATIONS
109	Arabidopsis Qa-SNARE SYP2 proteins localized to different subcellular regions function redundantly in vacuolar protein sorting and plant development. <i>Plant Journal</i> , 2010, 64, 924-935.	5.7	46
110	Stomagen positively regulates stomatal density in Arabidopsis. <i>Nature</i> , 2010, 463, 241-244.	27.8	382
111	Ectopic Expression of an Esterase, Which is a Candidate for the Unidentified Plant Cutinase, Causes Cuticular Defects in Arabidopsis thaliana. <i>Plant and Cell Physiology</i> , 2010, 51, 123-131.	3.1	105
112	MAG4/Atp115 is a Golgi-Localized Tethering Factor that Mediates Efficient Anterograde Transport in Arabidopsis. <i>Plant and Cell Physiology</i> , 2010, 51, 1777-1787.	3.1	33
113	The cystatin M/Î²cathepsin L balance is essential for tissue homeostasis in epidermis, hair follicles, and cornea. <i>FASEB Journal</i> , 2010, 24, 3744-3755.	0.5	37
114	Myosin-dependent endoplasmic reticulum motility and F-actin organization in plant cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 6894-6899.	7.1	306
115	Two vacuole-mediated defense strategies in plants. <i>Plant Signaling and Behavior</i> , 2010, 5, 1568-1570.	2.4	50
116	Vacuolar Processing Enzyme plays an Essential Role in the Crystalline Structure of Glutelin in Rice Seed. <i>Plant and Cell Physiology</i> , 2010, 51, 38-46.	3.1	74
117	Vacuolar SNAREs Function in the Formation of the Leaf Vascular Network by Regulating Auxin Distribution. <i>Plant and Cell Physiology</i> , 2009, 50, 1319-1328.	3.1	52
118	A novel membrane fusion-mediated plant immunity against bacterial pathogens. <i>Genes and Development</i> , 2009, 23, 2496-2506.	5.9	244
119	The ER body, a new organelle in <i>Arabidopsis thaliana</i> , requires NAI2 for its formation and accumulates specific Î²-glucosidases. <i>Plant Signaling and Behavior</i> , 2009, 4, 849-852.	2.4	23
120	GNOM-LIKE1/ERMO1 and SEC24a/ERMO2 Are Required for Maintenance of Endoplasmic Reticulum Morphology in <i>Arabidopsis thaliana</i> . <i>Plant Cell</i> , 2009, 21, 3672-3685.	6.6	92
121	An isoform of myosin XI is responsible for the translocation of endoplasmic reticulum in tobacco cultured BY-2 cells. <i>Journal of Experimental Botany</i> , 2009, 60, 197-212.	4.8	59
122	Dynamic Aspects of Ion Accumulation by Vesicle Traffic Under Salt Stress in Arabidopsis. <i>Plant and Cell Physiology</i> , 2009, 50, 2023-2033.	3.1	130
123	Quantitative Analysis of ER Body Morphology in an Arabidopsis Mutant. <i>Plant and Cell Physiology</i> , 2009, 50, 2015-2022.	3.1	29
124	Constitutive and Inducible ER Bodies of Arabidopsis thaliana Accumulate Distinct Î²-Glucosidases. <i>Plant and Cell Physiology</i> , 2009, 50, 480-488.	3.1	68
125	A novel role for oleosins in freezing tolerance of oilseeds in <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 2008, 55, 798-809.	5.7	184
126	AtMap1: a DNA microarray for genomic deletion mapping in <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 2008, 56, 1058-1065.	5.7	10



#	ARTICLE	IF	CITATIONS
127	Neuroprotective Actions of PIKE-L by Inhibition of SET Proteolytic Degradation by Asparagine Endopeptidase. <i>Molecular Cell</i> , 2008, 29, 665-678.	9.7	116
128	Antagonistic Jacalin-Related Lectins Regulate the Size of ER Body-Type $\beta$ -Glucosidase Complexes in <i>Arabidopsis thaliana</i> . <i>Plant and Cell Physiology</i> , 2008, 49, 969-980.	3.1	85
129	<i>Arabidopsis</i> VPS35, a Retromer Component, is Required for Vacuolar Protein Sorting and Involved in Plant Growth and Leaf Senescence. <i>Plant and Cell Physiology</i> , 2008, 49, 142-156.	3.1	105
130	<i>Arabidopsis</i> VPS35, a Retromer Component, is Required for Vacuolar Protein Sorting and Involved in Plant Growth and Leaf Senescence. <i>Plant and Cell Physiology</i> , 2008, 49, 678-678.	3.1	0
131	NAI2 Is an Endoplasmic Reticulum Body Component That Enables ER Body Formation in <i>Arabidopsis thaliana</i> . <i>Plant Cell</i> , 2008, 20, 2529-2540.	6.6	62
132	An Asparaginyl Endopeptidase Mediates in Vivo Protein Backbone Cyclization. <i>Journal of Biological Chemistry</i> , 2007, 282, 29721-29728.	3.4	207
133	<i>Arabidopsis</i> KAM2/GRV2 Is Required for Proper Endosome Formation and Functions in Vacuolar Sorting and Determination of the Embryo Growth Axis. <i>Plant Cell</i> , 2007, 19, 320-332.	6.6	83
134	MAIGO2 Is Involved in Exit of Seed Storage Proteins from the Endoplasmic Reticulum in <i>Arabidopsis thaliana</i> . <i>Plant Cell</i> , 2007, 18, 3535-3547.	6.6	79
135	<i>Arabidopsis</i> Vacuolar Sorting Mutants (green fluorescent seed) Can Be Identified Efficiently by Secretion of Vacuole-Targeted Green Fluorescent Protein in Their Seeds. <i>Plant Cell</i> , 2007, 19, 597-609.	6.6	87
136	AtVAM3 is Required for Normal Specification of Idioblasts, Myrosin Cells. <i>Plant and Cell Physiology</i> , 2006, 47, 164-175.	3.1	91
137	AtVPS29, a Putative Component of a Retromer Complex, is Required for the Efficient Sorting of Seed Storage Proteins. <i>Plant and Cell Physiology</i> , 2006, 47, 1187-1194.	3.1	135
138	Induction of Specialized Compartments from the ER. <i>Plant Cell Monographs</i> , 2006, , 141-154.	0.4	2
139	A VPE family supporting various vacuolar functions in plants. <i>Physiologia Plantarum</i> , 2005, 123, 369-375.	5.2	86
140	Endosomal proteases facilitate the fusion of endosomes with vacuoles at the final step of the endocytotic pathway. <i>Plant Journal</i> , 2005, 41, 888-898.	5.7	52
141	Vacuolar processing enzyme: an executor of plant cell death. <i>Current Opinion in Plant Biology</i> , 2005, 8, 404-408.	7.1	223
142	Activation of an ER-body-localized $\beta$ -Glucosidase via a Cytosolic Binding Partner in Damaged Tissues of <i>Arabidopsis thaliana</i> . <i>Plant and Cell Physiology</i> , 2005, 46, 1140-1148.	3.1	72
143	Identification of an Allele of VAM3/SYP22 that Confers a Semi-dwarf Phenotype in <i>Arabidopsis thaliana</i> . <i>Plant and Cell Physiology</i> , 2005, 46, 1358-1365.	3.1	41
144	Vacuolar Processing Enzyme Is Essential for Mycotoxin-induced Cell Death in <i>Arabidopsis thaliana</i> . <i>Journal of Biological Chemistry</i> , 2005, 280, 32914-32920.	3.4	196

#	ARTICLE	IF	CITATIONS
145	Asparagine Endopeptidase Is Not Essential for Class II MHC Antigen Presentation but Is Required for Processing of Cathepsin L in Mice. <i>Journal of Immunology</i> , 2005, 174, 7066-7074.	0.8	98
146	A Vacuolar Processing Enzyme, VPE, Is Involved in Seed Coat Formation at the Early Stage of Seed Development. <i>Plant Cell</i> , 2005, 17, 876-887.	6.6	199
147	KATAMARI1/MURUS3 Is a Novel Golgi Membrane Protein That Is Required for Endomembrane Organization in Arabidopsis. <i>Plant Cell</i> , 2005, 17, 1764-1776.	6.6	134
148	An ER-Localized Form of PV72, a Seed-Specific Vacuolar Sorting Receptor, Interferes the Transport of an NPIR-Containing Proteinase in Arabidopsis Leaves. <i>Plant and Cell Physiology</i> , 2004, 45, 9-17.	3.1	64
149	NAI1 Gene Encodes a Basic-Helix-Loop-Helix-Type Putative Transcription Factor That Regulates the Formation of an Endoplasmic Reticulum-Derived Structure, the ER Body. <i>Plant Cell</i> , 2004, 16, 1536-1549.	6.6	99
150	Endoplasmic reticulum-resident proteins are constitutively transported to vacuoles for degradation. <i>Plant Journal</i> , 2004, 39, 393-402.	5.7	53
151	The slow wound-response of VPE is regulated by endogenous salicylic acid in Arabidopsis. <i>Planta</i> , 2004, 218, 599-605.	3.2	36
152	Diversity and Formation of Endoplasmic Reticulum-Derived Compartments in Plants. Are These Compartments Specific to Plant Cells?. <i>Plant Physiology</i> , 2004, 136, 3435-3439.	4.8	61
153	A Plant Vacuolar Protease, VPE, Mediates Virus-Induced Hypersensitive Cell Death. <i>Science</i> , 2004, 305, 855-858.	12.6	579
154	A wound-inducible organelle derived from endoplasmic reticulum: a plant strategy against environmental stresses?. <i>Current Opinion in Plant Biology</i> , 2003, 6, 583-588.	7.1	53
155	A novel ER-derived compartment, the ER body, selectively accumulates a $\beta$ -glucosidase with an ER-retention signal in Arabidopsis. <i>Plant Journal</i> , 2003, 33, 493-502.	5.7	172
156	Why green fluorescent fusion proteins have not been observed in the vacuoles of higher plants. <i>Plant Journal</i> , 2003, 35, 545-555.	5.7	226
157	Biosynthetic Processing of Cathepsins and Lysosomal Degradation Are Abolished in Asparaginyl Endopeptidase-deficient Mice. <i>Journal of Biological Chemistry</i> , 2003, 278, 33194-33199.	3.4	181
158	The ER Body, a Novel Endoplasmic Reticulum-Derived Structure in Arabidopsis. <i>Plant and Cell Physiology</i> , 2003, 44, 661-666.	3.1	92
159	C-Terminal KDEL Sequence of A KDEL-Tailed Cysteine Proteinase (Sulfhydryl-Endopeptidase) Is Involved in Formation of KDEL Vesicle and in Efficient Vacuolar Transport of Sulfhydryl-Endopeptidase. <i>Plant Physiology</i> , 2003, 132, 1892-1900.	4.8	56
160	Vacuolar Processing Enzymes Are Essential for Proper Processing of Seed Storage Proteins in Arabidopsis thaliana. <i>Journal of Biological Chemistry</i> , 2003, 278, 32292-32299.	3.4	189
161	Vacuolar sorting receptor for seed storage proteins in Arabidopsis thaliana. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 16095-16100.	7.1	235
162	Behavior of Vacuoles during Microspore and Pollen Development in Arabidopsis thaliana. <i>Plant and Cell Physiology</i> , 2003, 44, 1192-1201.	3.1	99

#	ARTICLE	IF	CITATIONS
163	A Vacuolar Sorting Receptor PV72 on the Membrane of Vesicles that Accumulate Precursors of Seed Storage Proteins (PAC Vesicles). <i>Plant and Cell Physiology</i> , 2002, 43, 1086-1095.	3.1	74
164	Calcium-mediated Association of a Putative Vacuolar Sorting Receptor PV72 with a Propeptide of 2S Albumin. <i>Journal of Biological Chemistry</i> , 2002, 277, 8708-8715.	3.4	64
165	An Endoplasmic Reticulum-Derived Structure That Is Induced under Stress Conditions in Arabidopsis. <i>Plant Physiology</i> , 2002, 130, 1807-1814.	4.8	147
166	Molecular Characterization of an Arabidopsis Acyl-Coenzyme A Synthetase Localized on Glyoxysomal Membranes. <i>Plant Physiology</i> , 2002, 130, 2019-2026.	4.8	40
167	Activation of Arabidopsis Vacuolar Processing Enzyme by Self-Catalytic Removal of an Auto-Inhibitory Domain of the C-Terminal Propeptide. <i>Plant and Cell Physiology</i> , 2002, 43, 143-151.	3.1	112
168	A Novel Membrane Protein That Is Transported to Protein Storage Vacuoles via Precursor-Accumulating Vesicles. <i>Plant Cell</i> , 2001, 13, 2361-2372.	6.6	31
169	A Slow Maturation of a Cysteine Protease with a Granulin Domain in the Vacuoles of Senescing Arabidopsis Leaves. <i>Plant Physiology</i> , 2001, 127, 1626-1634.	4.8	158
170	A Proteinase-Storing Body that Prepares for Cell Death or Stresses in the Epidermal Cells of Arabidopsis. <i>Plant and Cell Physiology</i> , 2001, 42, 894-899.	3.1	208
171	Characterization of an Arabidopsis cDNA Encoding a Subunit of Serine Palmitoyltransferase, the Initial Enzyme in Sphingolipid Biosynthesis. <i>Plant and Cell Physiology</i> , 2001, 42, 1274-1281.	3.1	36
172	Characterization of Organelles in the Vacuolar-Sorting Pathway by Visualization with GFP in Tobacco BY-2 Cells. <i>Plant and Cell Physiology</i> , 2000, 41, 993-1001.	3.1	138
173	Multiple Functional Proteins Are Produced by Cleaving Asn-Gln Bonds of a Single Precursor by Vacuolar Processing Enzyme. <i>Journal of Biological Chemistry</i> , 1999, 274, 2563-2570.	3.4	98
174	Chloroplast Cpn20 forms a tetrameric structure in Arabidopsis thaliana. <i>Plant Journal</i> , 1999, 17, 467-477.	5.7	46
175	Vacuolar processing enzyme is up-regulated in the lytic vacuoles of vegetative tissues during senescence and under various stressed conditions. <i>Plant Journal</i> , 1999, 19, 43-53.	5.7	215
176	Vacuolar processing enzyme is self-catalytically activated by sequential removal of the C-terminal and N-terminal propeptides. <i>FEBS Letters</i> , 1999, 447, 213-216.	2.8	101
177	Vacuolar processing enzymes in protein-storage vacuoles and lytic vacuoles. <i>Journal of Plant Physiology</i> , 1998, 152, 668-674.	3.5	78
178	Transport of Storage Proteins to Protein Storage Vacuoles Is Mediated by Large Precursor-Accumulating Vesicles. <i>Plant Cell</i> , 1998, 10, 825-836.	6.6	307
179	The AtVAM3 Encodes a Syntaxin-related Molecule Implicated in the Vacuolar Assembly in Arabidopsis thaliana. <i>Journal of Biological Chemistry</i> , 1997, 272, 24530-24535.	3.4	109
180	An Aspartic Endopeptidase is Involved in the Breakdown of Propeptides of Storage Proteins in Protein-Storage Vacuoles of Plants. <i>FEBS Journal</i> , 1997, 246, 133-141.	0.2	104

#	ARTICLE	IF	CITATIONS
181	Expression and activation of the vacuolar processing enzyme in <i>Saccharomyces cerevisiae</i> . <i>Plant Journal</i> , 1997, 12, 819-829.	5.7	51
182	Isolation and characterization of a cDNA encoding mitochondrial chaperonin 10 from <i>Arabidopsis thaliana</i> by functional complementation of an <i>Escherichia coli</i> groES mutant. <i>Plant Journal</i> , 1996, 10, 1119-1125.	5.7	18
183	Molecular characterization of proteins in protein-body membrane that disappear most rapidly during transformation of protein bodies into vacuoles. <i>Plant Journal</i> , 1995, 7, 235-243.	5.7	54
184	Homologues of a vacuolar processing enzyme that are expressed in different organs in <i>Arabidopsis thaliana</i> . <i>Plant Molecular Biology</i> , 1995, 29, 81-89.	3.9	65
185	Vacuolar Processing Enzyme Responsible for Maturation of Seed Proteins. <i>Journal of Plant Physiology</i> , 1995, 145, 632-640.	3.5	125
186	Immunological analysis of aconitase in pumpkin cotyledons: the absence of aconitase in glyoxysomes. <i>Physiologia Plantarum</i> , 1994, 90, 757-762.	5.2	28
187	Vacuolar Processing Enzyme of Soybean That Converts Proproteins to the Corresponding Mature Forms. <i>Plant and Cell Physiology</i> , 1994, 35, 713-718.	3.1	87
188	Amino acid sequence surrounding the retinal-binding site in retinochrome of the squid, <i>Todarodes pacificus</i> . <i>FEBS Letters</i> , 1993, 335, 94-98.	2.8	12
189	Cloning and nucleotide sequence of cDNA for rhodopsin of the squid <i>Todarodes pacificus</i> . <i>FEBS Letters</i> , 1993, 317, 5-11.	2.8	37
190	Molecular Characterization of a Vacuolar Processing Enzyme Related to a Putative Cysteine Proteinase of <i>Schistosoma mansoni</i> . <i>Plant Cell</i> , 1993, 5, 1651.	6.6	34
191	A unique vacuolar processing enzyme responsible for conversion of several proprotein precursors into the mature forms. <i>FEBS Letters</i> , 1991, 294, 89-93.	2.8	229
192	Pumpkin malate synthase. Cloning and sequencing of the cDNA and Northern blot analysis. <i>FEBS Journal</i> , 1991, 197, 331-336.	0.2	53
193	Cloning and nucleotide sequence of cDNA for retinochrome, retinal photoisomerase from the squid retina. <i>FEBS Letters</i> , 1990, 271, 106-110.	2.8	39
194	Nucleotide sequence of cloned cDNA coding for pumpkin 11-S globulin beta subunit. <i>FEBS Journal</i> , 1988, 172, 627-632.	0.2	30
195	Effect of monensin on intracellular transport and posttranslational processing of 11 S globulin precursors in developing pumpkin cotyledons. <i>FEBS Letters</i> , 1988, 238, 197-200.	2.8	7
196	Proglobulin Processing Enzyme in Vacuoles Isolated from Developing Pumpkin Cotyledons. <i>Plant Physiology</i> , 1987, 85, 440-445.	4.8	104
197	Biosynthesis and Intracellular Transport of 11S Globulin in Developing Pumpkin Cotyledons. <i>Plant Physiology</i> , 1985, 77, 747-752.	4.8	56
198	Suborganellar Localization of Proteinase Catalyzing the Limited Hydrolysis of Pumpkin Globulin. <i>Plant Physiology</i> , 1982, 70, 699-703.	4.8	39