

# Ali Reza Heidari-Bakavoli

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1834843/publications.pdf>

Version: 2024-02-01

28  
papers

685  
citations

687363

13  
h-index

610901

24  
g-index

29  
all docs

29  
docs citations

29  
times ranked

1181  
citing authors

#	ARTICLE	IF	CITATIONS
1	Mashhad stroke and heart atherosclerotic disorder (MASHAD) study: design, baseline characteristics and 10-year cardiovascular risk estimation. <i>International Journal of Public Health</i> , 2015, 60, 561-572.	2.3	114
2	Depression and anxiety both associate with serum level of hs-CRP: A gender-stratified analysis in a population-based study. <i>Psychoneuroendocrinology</i> , 2017, 81, 63-69.	2.7	95
3	Depression and anxiety symptoms are associated with white blood cell count and red cell distribution width: A sex-stratified analysis in a population-based study. <i>Psychoneuroendocrinology</i> , 2017, 84, 101-108.	2.7	78
4	Association of Serum hs-CRP Levels With the Presence of Obesity, Diabetes Mellitus, and Other Cardiovascular Risk Factors. <i>Journal of Clinical Laboratory Analysis</i> , 2016, 30, 672-676.	2.1	58
5	Nutrient patterns and their relationship to metabolic syndrome in Iranian adults. <i>European Journal of Clinical Investigation</i> , 2016, 46, 840-852.	3.4	51
6	Association of hematocrit with blood pressure and hypertension. <i>Journal of Clinical Laboratory Analysis</i> , 2017, 31, .	2.1	51
7	Association between dietary inflammatory index and risk of cardiovascular disease in the Mashhad stroke and heart atherosclerotic disorder study population. <i>IUBMB Life</i> , 2020, 72, 706-715.	3.4	36
8	Oxidative stress and inflammation, two features associated with a high percentage body fat, and that may lead to diabetes mellitus and metabolic syndrome. <i>BioFactors</i> , 2019, 45, 35-42.	5.4	33
9	Interaction between a variant of CDKN2A/B-gene with lifestyle factors in determining dyslipidemia and estimated cardiovascular risk: A step toward personalized nutrition. <i>Clinical Nutrition</i> , 2018, 37, 254-261.	5.0	27
10	Prevalence of combined and noncombined dyslipidemia in an Iranian population. <i>Journal of Clinical Laboratory Analysis</i> , 2018, 32, e22579.	2.1	22
11	Association between serum uric acid, high sensitive C-reactive protein and pro-oxidant-antioxidant balance in patients with metabolic syndrome. <i>BioFactors</i> , 2018, 44, 263-271.	5.4	20
12	Changes in Plasma Level of Heat Shock Protein 27 After Acute Coronary Syndrome. <i>Angiology</i> , 2012, 63, 12-16.	1.8	17
13	Hookah smoking is strongly associated with diabetes mellitus, metabolic syndrome and obesity: a population-based study. <i>Diabetology and Metabolic Syndrome</i> , 2018, 10, 33.	2.7	17
14	Evaluating of associated risk factors of metabolic syndrome by using decision tree. <i>Comparative Clinical Pathology</i> , 2018, 27, 215-223.	0.7	13
15	There is an association between body fat percentage and metabolic abnormality in normal weight subjects: Iranian large population. <i>Translational Metabolic Syndrome Research</i> , 2019, 2, 11-16.	0.8	10
16	Lack of association between LXR <sup>1</sup> and LXR <sup>2</sup> gene polymorphisms and prevalence of metabolic syndrome: A case-control study of an Iranian population. <i>Gene</i> , 2013, 532, 288-293.	2.2	7
17	The relationship between dietary intake and other cardiovascular risk factors with blood pressure in individuals without a history of a cardiovascular event: Evidence based study with 5670 subjects. <i>Diabetes and Metabolic Syndrome: Clinical Research and Reviews</i> , 2017, 11, S65-S71.	3.6	7
18	Relationship between platelet count and platelet width distribution and serum uric acid concentrations in patients with untreated essential hypertension. <i>BioFactors</i> , 2018, 44, 532-538.	5.4	5

#	ARTICLE	IF	CITATIONS
19	Prevalence of ABO and Rh blood groups and their association with demographic and anthropometric factors in an Iranian population: Mashad study. <i>Eastern Mediterranean Health Journal</i> , 2020, 26, 916-922.	0.8	5
20	Cardiovascular Risk Factors and Nutritional Intake are not Associated with Ultrasound-defined Increased Carotid Intima Media Thickness in Individuals Without a History of Cardiovascular Events. <i>International Journal of Preventive Medicine</i> , 2014, 5, 1412-21.	0.4	5
21	Prevalence of hypertension, pre-hypertension and undetected hypertension in Mashhad, Iran. <i>Mediterranean Journal of Nutrition and Metabolism</i> , 2017, 9, 213-223.	0.5	4
22	The interaction between a HSP-70 gene variant with dietary calories in determining serum markers of inflammation and cardiovascular risk. <i>Clinical Nutrition</i> , 2018, 37, 2122-2126.	5.0	4
23	Is there any association between Serum anti-HSP27 antibody level and the presence of metabolic syndrome; population based case-control study. <i>Romanian Journal of Laboratory Medicine</i> , 2019, 27, 179-187.	0.2	3
24	Macronutrient intake and physical activity levels in individuals with and without metabolic syndrome: An observational study in an urban population. <i>ARYA Atherosclerosis</i> , 2019, 15, 136-145.	0.4	2
25	Distribution of obesity phenotypes and in a population-based sample of Iranian adults. <i>Mediterranean Journal of Nutrition and Metabolism</i> , 2017, 9, 203-212.	0.5	0
26	The association between a variant of the cyclin-dependent kinase inhibitor 2A/B gene and risk of cardiovascular disease. <i>Gene Reports</i> , 2022, 26, 101480.	0.8	0
27	Successful transvenous lead extraction of abandoned lead implanted through persistent left superior vena cava. <i>Future Cardiology</i> , 2022, 18, 185-190.	1.2	0
28	A Genetic Variant in Proline and Serine Rich Coiled-Coil 1 Gene Is Associated with the Risk of Cardiovascular Disease. <i>Reports of Biochemistry and Molecular Biology</i> , 2022, 10, 653-663.	1.4	0