

T EstefanÃ-a Saez Lancellotti

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1835458/publications.pdf>

Version: 2024-02-01

14
papers

285
citations

1162889

8
h-index

1058333

14
g-index

14
all docs

14
docs citations

14
times ranked

439
citing authors

#	ARTICLE	IF	CITATIONS
1	Impact of high fat diet on the sterol regulatory element-binding protein 2 cholesterol pathway in the testicle. <i>Molecular Human Reproduction</i> , 2021, 27, .	1.3	4
2	A chronic high-fat diet causes sperm head alterations in C57BL/6J mice. <i>Heliyon</i> , 2019, 5, e02868.	1.4	9
3	ODF1, sperm flagellar protein is expressed in kidney collecting ducts of rats. <i>Heliyon</i> , 2019, 5, e02932.	1.4	3
4	Olive oil addition to the high-fat diet reduces methylglyoxal (MG-H1) levels increased in hypercholesterolemic rabbits. <i>Mediterranean Journal of Nutrition and Metabolism</i> , 2019, 12, 13-21.	0.2	3
5	Pigment epithelium derived factor (PEDF) expression in the male tract of Wistar rats. <i>Biochemical and Biophysical Research Communications</i> , 2018, 504, 257-262.	1.0	4
6	Manchette-acrosome disorders and testicular efficiency decline observed in hypercholesterolemic rabbits are recovered with olive oil enriched diet. <i>PLoS ONE</i> , 2018, 13, e0202748.	1.1	8
7	Thiols of flagellar proteins are essential for progressive motility in human spermatozoa. <i>Reproduction, Fertility and Development</i> , 2017, 29, 1435.	0.1	9
8	Manchette-acrosome disorders during spermiogenesis and low efficiency of seminiferous tubules in hypercholesterolemic rabbit model. <i>PLoS ONE</i> , 2017, 12, e0172994.	1.1	15
9	Semen Quality and Sperm Function Loss by Hypercholesterolemic Diet Was Recovered by Addition of Olive Oil to Diet in Rabbit. <i>PLoS ONE</i> , 2013, 8, e52386.	1.1	42
10	SPINK3 modulates mouse sperm physiology through the reduction of nitric oxide level independently of its trypsin inhibitory activity. <i>Reproduction</i> , 2012, 143, 281-295.	1.1	25
11	Characterization of flagellar cysteine-rich sperm proteins involved in motility, by the combination of cellular fractionation, fluorescence detection, and mass spectrometry analysis. <i>Cytoskeleton</i> , 2011, 68, 491-500.	1.0	21
12	Protein fraction isolated from epididymal fluid re-associates sperm in vitro: Possible role of serpins in rat rosettes assembly. <i>Molecular Reproduction and Development</i> , 2010, 77, 410-419.	1.0	9
13	Hypercholesterolemia Impaired Sperm Functionality in Rabbits. <i>PLoS ONE</i> , 2010, 5, e13457.	1.1	83
14	Production of nerve growth factor by β 2-amyloid-stimulated astrocytes induces p75NTR-dependent tau hyperphosphorylation in cultured hippocampal neurons. <i>Journal of Neuroscience Research</i> , 2006, 84, 1098-1106.	1.3	50