

Cinthia Eyng

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1839666/publications.pdf>

Version: 2024-02-01

17
papers

61
citations

1937685

4
h-index

1720034

7
g-index

18
all docs

18
docs citations

18
times ranked

42
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of lipid source and emulsifier on productive and physiological parameters of broilers. <i>Animal Bioscience</i> , 2022, 35, 54-63.	2.0	6
2	Influence of Enzyme Supplementation in the Diets of Broiler Chickens Formulated with Different Corn Hybrids Dried at Various Temperatures. <i>Animals</i> , 2021, 11, 643.	2.3	4
3	Carnauba wax coating preserves the internal quality of commercial eggs during storage. <i>Semina:Ciencias Agrarias</i> , 2021, 42, 1229-1244.	0.3	6
4	Performance and biometry of the gastrointestinal tract in growing Japanese quails fed bee pollen. <i>Semina:Ciencias Agrarias</i> , 2021, 42, 1379-1386.	0.3	0
5	Different enzymatic associations in diets of broiler chickens formulated with corn dried at various temperatures. <i>Poultry Science</i> , 2021, 100, 101013.	3.4	5
6	Composition and availability of nutrients of three corn hybrids dried at different temperatures in the diets of broiler chickens. <i>Revista Brasileira De Zootecnia</i> , 2021, 50, .	0.8	1
7	Replacement of corn by rice grits, with or without phytase supplementation, in diets for silver catfish (<i>Rhamdia quelen</i>) juveniles. <i>Aquaculture Nutrition</i> , 2020, 26, 1584-1591.	2.7	1
8	Carcass characteristics and meat quality of slow-growing broilers fed diets containing dry residue of cassava, with or without the addition of carbohydrases. <i>Tropical Animal Health and Production</i> , 2020, 52, 2677-2685.	1.4	3
9	Valores energéticos e composição bromatológica do resíduo seco de fecularia associado a carboidrases para frangos de corte em fase de crescimento. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2020, 72, 1504-1510.	0.4	0
10	Dried Cassava Residue in Laying Quail Feeding. <i>Brazilian Journal of Poultry Science</i> , 2020, 22, .	0.7	3
11	Energy values of crude glycerin for broilers. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2020, 72, 2348-2354.	0.4	1
12	Energetic Values and Inclusion Levels of the Dry Residue of Cassava in Broiler Diet. <i>Brazilian Journal of Poultry Science</i> , 2020, 22, .	0.7	0
13	Technical Cashew Nutshell Liquid in Diets of Growing Meat-Type Quails. <i>Brazilian Journal of Poultry Science</i> , 2019, 21, .	0.7	0
14	Dietary inclusion of dehydrated bocaiuva pulp increases the antioxidant potential of quail eggs. <i>Journal of Animal Physiology and Animal Nutrition</i> , 2019, 103, 64-71.	2.2	6
15	Performance and Carcass Yield of Broilers Fed Crude Glycerin at Differing Inclusion Levels. <i>Brazilian Journal of Poultry Science</i> , 2018, 20, 797-804.	0.7	0
16	High levels of dietary phytase improves broiler performance. <i>Animal Feed Science and Technology</i> , 2018, 244, 56-65.	2.2	15
17	Cassava starch factory residues in the diet of slow-growing broilers. <i>Tropical Animal Health and Production</i> , 2014, 46, 1371-1381.	1.4	8