Mar Hernndez-Guillamon

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1841880/mar-hernandez-guillamon-publications-by-year.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

58
papers2,252
citations27
h-index47
g-index59
ext. papers2,753
ext. citations5.7
avg, IF4.7
L-index

#	Paper	IF	Citations
58	Circulating AQP4 Levels in Patients with Cerebral Amyloid Angiopathy-Associated Intracerebral Hemorrhage. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	3
57	Survival Bias and Crosstalk between Chronological and Behavioral Age: Age- and Genotype-Sensitivity Tests Define Behavioral Signatures in Middle-Aged, Old, and Long-Lived Mice with Normal and AD-Associated Aging. <i>Biomedicines</i> , 2021 , 9,	4.8	6
56	New candidate blood biomarkers potentially associated with white matter hyperintensities progression. <i>Scientific Reports</i> , 2021 , 11, 14324	4.9	O
55	Comparison of Plasma Lipoprotein Composition and Function in Cerebral Amyloid Angiopathy and Alzheimer U Disease. <i>Biomedicines</i> , 2021 , 9,	4.8	2
54	SSAO/VAP-1 in Cerebrovascular Disorders: A Potential Therapeutic Target for Stroke and Alzheimer u Disease. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	5
53	Association of CD2AP neuronal deposits with Braak neurofibrillary stage in Alzheimer u disease. <i>Brain Pathology</i> , 2021 , e13016	6	1
52	MFG-E8 (LACTADHERIN): a novel marker associated with cerebral amyloid angiopathy. <i>Acta Neuropathologica Communications</i> , 2021 , 9, 154	7.3	2
51	Circulating TIMP-1 is associated with hematoma volume in patients with spontaneous intracranial hemorrhage. <i>Scientific Reports</i> , 2020 , 10, 10329	4.9	1
50	CCL23: A Chemokine Associated with Progression from Mild Cognitive Impairment to Alzheimer u Disease. <i>Journal of Alzheimerus Disease</i> , 2020 , 73, 1585-1595	4.3	10
49	Cerebral amyloid angiopathy and Alzheimer disease - one peptide, two pathways. <i>Nature Reviews Neurology</i> , 2020 , 16, 30-42	15	171
48	Advancing diagnostic criteria for sporadic cerebral amyloid angiopathy: Study protocol for a multicenter MRI-pathology validation of Boston criteria v2.0. <i>International Journal of Stroke</i> , 2019 , 14, 956-971	6.3	18
47	Matrix metalloproteinases and ADAMs in stroke. Cellular and Molecular Life Sciences, 2019, 76, 3117-314	40 0.3	21
46	Peripheral administration of human recombinant ApoJ/clusterin modulates brain beta-amyloid levels in APP23 mice. <i>Alzheimerts Research and Therapy</i> , 2019 , 11, 42	9	15
45	Brain ApoA-I, ApoJ and ApoE Immunodetection in Cerebral Amyloid Angiopathy. <i>Frontiers in Neurology</i> , 2019 , 10, 187	4.1	11
44	PATJ Low Frequency Variants Are Associated With Worse Ischemic Stroke Functional Outcome. <i>Circulation Research</i> , 2019 , 124, 114-120	15.7	27
43	Absolute risk and predictors of the growth of acute spontaneous intracerebral haemorrhage: a systematic review and meta-analysis of individual patient data. <i>Lancet Neurology, The</i> , 2018 , 17, 885-894	4 ^{24.1}	142
42	Simvastatin blocks soluble SSAO/VAP-1 release in experimental models of cerebral ischemia: Possible benefits for stroke-induced inflammation control. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018 , 1864, 542-553	6.9	8

(2014-2017)

41	deposition in the APP23-transgenic mouse model of Alzheimer U disease. <i>Neurobiology of Aging</i> , 2017 , 60, 116-128	5.6	20
40	Brain hemorrhage recurrence, small vessel disease type, and cerebral microbleeds: A meta-analysis. <i>Neurology</i> , 2017 , 89, 820-829	6.5	115
39	Characterization of ApoJ-reconstituted high-density lipoprotein (rHDL) nanodisc for the potential treatment of cerebral Emyloidosis. <i>Scientific Reports</i> , 2017 , 7, 14637	4.9	22
38	Profiling and identification of new proteins involved in brain ischemia using MALDI-imaging-mass-spectrometry. <i>Journal of Proteomics</i> , 2017 , 152, 243-253	3.9	18
37	Plasmatic retinol-binding protein 4 and glial fibrillary acidic protein as biomarkers to differentiate ischemic stroke and intracerebral hemorrhage. <i>Journal of Neurochemistry</i> , 2016 , 136, 416-24	6	32
36	Identification of Plasma Biomarkers of Human Intracerebral Hemorrhage Subtypes through Microarray Technology. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2016 , 25, 665-71	2.8	2
35	Characterization of secretomes from a human blood brain barrier endothelial cells in-vitro model after ischemia by stable isotope labeling with aminoacids in cell culture (SILAC). <i>Journal of Proteomics</i> , 2016 , 133, 100-112	3.9	13
34	ApoA1, ApoJ and ApoE Plasma Levels and Genotype Frequencies in Cerebral Amyloid Angiopathy. <i>NeuroMolecular Medicine</i> , 2016 , 18, 99-108	4.6	16
33	Charge effect of a liposomal delivery system encapsulating simvastatin to treat experimental ischemic stroke in rats. <i>International Journal of Nanomedicine</i> , 2016 , 11, 3035-48	7.3	47
32	Modulation of Amyloid-11-40 Transport by ApoA1 and ApoJ Across an in vitro Model of the Blood-Brain Barrier. <i>Journal of Alzheimerts Disease</i> , 2016 , 53, 677-91	4.3	32
31	Cerebral amyloid angiopathy-related atraumatic convexal subarachnoid hemorrhage: an ARIA before the tsunami. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2015 , 35, 710-7	7.3	30
30	Sequential Amyloid-IDegradation by the Matrix Metalloproteases MMP-2 and MMP-9. <i>Journal of Biological Chemistry</i> , 2015 , 290, 15078-91	5.4	73
29	NURR1 involvement in recombinant tissue-type plasminogen activator treatment complications after ischemic stroke. <i>Stroke</i> , 2015 , 46, 477-84	6.7	13
28	Matrix Metalloproteinases in Alzheimerld Disease and Concurrent Cerebral Microbleeds. <i>Journal of Alzheimerls Disease</i> , 2015 , 48, 711-20	4.3	47
27	Mild hypothermia protects against oxygen glucose deprivation (OGD)-induced cell death in brain slices from adult mice. <i>Journal of Neural Transmission</i> , 2014 , 121, 113-7	4.3	4
26	Brain proteomics identifies potential simvastatin targets in acute phase of stroke in a rat embolic model. <i>Journal of Neurochemistry</i> , 2014 , 130, 301-12	6	19
25	Fluorescent molecular peroxidation products: a prognostic biomarker of early neurologic deterioration after thrombolysis. <i>Stroke</i> , 2014 , 45, 432-7	6.7	10
24	Rat middle cerebral artery occlusion is not a suitable model for the study of stroke-induced spontaneous infections. <i>PLoS ONE</i> , 2014 , 9, e99169	3.7	2

23	The angiogenic gene profile of circulating endothelial progenitor cells from ischemic stroke patients. <i>Vascular Cell</i> , 2013 , 5, 3	1	15
22	Cerebral ischaemia and matrix metalloproteinase-9 modulate the angiogenic function of early and late outgrowth endothelial progenitor cells. <i>Journal of Cellular and Molecular Medicine</i> , 2013 , 17, 1543-	5 5 .6	29
21	Combining statins with tissue plasminogen activator treatment after experimental and human stroke: a safety study on hemorrhagic transformation. <i>CNS Neuroscience and Therapeutics</i> , 2013 , 19, 863	3- 7 8	9
20	Genes involved in hemorrhagic transformations that follow recombinant t-PA treatment in stroke patients. <i>Pharmacogenomics</i> , 2013 , 14, 495-504	2.6	16
19	Factors secreted by endothelial progenitor cells enhance neurorepair responses after cerebral ischemia in mice. <i>PLoS ONE</i> , 2013 , 8, e73244	3.7	79
18	MMP-2/MMP-9 plasma level and brain expression in cerebral amyloid angiopathy-associated hemorrhagic stroke. <i>Brain Pathology</i> , 2012 , 22, 133-41	6	54
17	Evidence for the efficacy of statins in animal stroke models: a meta-analysis. <i>Journal of Neurochemistry</i> , 2012 , 122, 233-43	6	56
16	Differentiating ischemic from hemorrhagic stroke using plasma biomarkers: the S100B/RAGE pathway. <i>Journal of Proteomics</i> , 2012 , 75, 4758-65	3.9	47
15	Plasma Emyloid levels in cerebral amyloid angiopathy-associated hemorrhagic stroke. <i>Neurodegenerative Diseases</i> , 2012 , 10, 320-3	2.3	30
14	VAP-1/SSAO plasma activity and brain expression in human hemorrhagic stroke. <i>Cerebrovascular Diseases</i> , 2012 , 33, 55-63	3.2	36
13	ACE variants and risk of intracerebral hemorrhage recurrence in amyloid angiopathy. <i>Neurobiology of Aging</i> , 2011 , 32, 551.e13-22	5.6	21
12	A large screening of angiogenesis biomarkers and their association with neurological outcome after ischemic stroke. <i>Atherosclerosis</i> , 2011 , 216, 205-11	3.1	84
11	Matrix metalloproteinase 2 (MMP-2) degrades soluble vasculotropic amyloid-beta E22Q and L34V mutants, delaying their toxicity for human brain microvascular endothelial cells. <i>Journal of Biological Chemistry</i> , 2010 , 285, 27144-27158	5.4	39
10	Plasma VAP-1/SSAO activity predicts intracranial hemorrhages and adverse neurological outcome after tissue plasminogen activator treatment in stroke. <i>Stroke</i> , 2010 , 41, 1528-35	6.7	59
9	Mobilization, endothelial differentiation and functional capacity of endothelial progenitor cells after ischemic stroke. <i>Microvascular Research</i> , 2010 , 80, 317-23	3.7	63
8	The proteome of human brain after ischemic stroke. <i>Journal of Neuropathology and Experimental Neurology</i> , 2010 , 69, 1105-15	3.1	38
7	Neuronal TIMP-1 release accompanies astrocytic MMP-9 secretion and enhances astrocyte proliferation induced by beta-amyloid 25-35 fragment. <i>Journal of Neuroscience Research</i> , 2009 , 87, 2115	5- 2:1	32
6	Matrix metalloproteinase-13 is activated and is found in the nucleus of neural cells after cerebral ischemia. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2009 , 29, 398-410	7.3	59

LIST OF PUBLICATIONS

5	embrane-bound SSAO/VAP-1. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2008 , 1783, 1085 ⁴⁹ 4		24
4	Tissue plasminogen activator (t-PA) promotes neutrophil degranulation and MMP-9 release. Journal of Leukocyte Biology, 2008 , 84, 207-14	6.5	100
3	MMP-9-positive neutrophil infiltration is associated to blood-brain barrier breakdown and basal lamina type IV collagen degradation during hemorrhagic transformation after human ischemic stroke. <i>Stroke</i> , 2008 , 39, 1121-6	6.7	365
2	Fas system activation in perihematomal areas after spontaneous intracerebral hemorrhage. <i>Stroke</i> , 2008 , 39, 1730-4	6.7	34
1	Sodium bicarbonate enhances membrane-bound and soluble human semicarbazide-sensitive amine oxidase activity in vitro. <i>Journal of Biochemistry</i> , 2007 , 142, 571-6	3.1	5