Gianluca Tell

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1842669/gianluca-tell-publications-by-year.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

 169
 6,519
 43
 72

 papers
 citations
 h-index
 g-index

 175
 7,189
 5.3
 5.49

 ext. papers
 ext. citations
 avg, IF
 L-index

#	Paper	IF	Citations
169	Integrated multi-omics analyses on patient-derived CRC organoids highlight altered molecular pathways in colorectal cancer progression involving PTEN. <i>Journal of Experimental and Clinical Cancer Research</i> , 2021 , 40, 198	12.8	5
168	Enzymatically active apurinic/apyrimidinic endodeoxyribonuclease 1 is released by mammalian cells through exosomes. <i>Journal of Biological Chemistry</i> , 2021 , 296, 100569	5.4	5
167	Coping with RNA damage with a focus on APE1, a BER enzyme at the crossroad between DNA damage repair and RNA processing/decay. <i>DNA Repair</i> , 2021 , 104, 103133	4.3	4
166	Brain Exosomes: Friend or Foe in Alzheimer's Disease?. <i>Molecular Neurobiology</i> , 2021 , 58, 6610-6624	6.2	2
165	Role of phase partitioning in coordinating DNA damage response: focus on the Apurinic Apyrimidinic Endonuclease 1 interactome. <i>Biomolecular Concepts</i> , 2020 , 11, 209-220	3.7	2
164	Architecture of The Human Ape1 Interactome Defines Novel Cancers Signatures. <i>Scientific Reports</i> , 2020 , 10, 28	4.9	7
163	New perspectives in cancer biology from a study of canonical and non-canonical functions of base excision repair proteins with a focus on early steps. <i>Mutagenesis</i> , 2020 , 35, 129-149	2.8	14
162	Differential capacity of CD90+ cells in autophagy activation following chemotherapy in hepatocellular carcinoma. <i>Annals of Hepatology</i> , 2020 , 19, 645-652	3.1	O
161	Cleavage of the APE1 N-Terminal Domain in Acute Myeloid Leukemia Cells Is Associated with Proteasomal Activity. <i>Biomolecules</i> , 2020 , 10,	5.9	4
160	Somatostatin as Inflow Modulator in Liver-transplant Recipients With Severe Portal Hypertension: A Randomized Trial. <i>Annals of Surgery</i> , 2019 , 269, 1025-1033	7.8	23
159	Inhibition of APE1-endonuclease activity affects cell metabolism in colon cancer cells via a p53-dependent pathway. <i>DNA Repair</i> , 2019 , 82, 102675	4.3	16
158	Platinum Salts in Patients with Breast Cancer: A Focus on Predictive Factors. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	23
157	Unlike the counterpart, archaeal RNase HII cannot process ribose monophosphate abasic sites and oxidized ribonucleotides embedded in DNA. <i>Journal of Biological Chemistry</i> , 2019 , 294, 13061-13072	5.4	6
156	APE1 and NPM1 protect cancer cells from platinum compounds cytotoxicity and their expression pattern has a prognostic value in TNBC. <i>Journal of Experimental and Clinical Cancer Research</i> , 2019 , 38, 309	12.8	15
155	Serum AP-endonuclease 1 (sAPE1) as novel biomarker for hepatocellular carcinoma. <i>Oncotarget</i> , 2019 , 10, 383-394	3.3	19
154	Endonuclease and redox activities of human apurinic/apyrimidinic endonuclease 1 have distinctive and essential functions in IgA class switch recombination. <i>Journal of Biological Chemistry</i> , 2019 , 294, 51	98:420)7 ¹²
153	The redox-sensitive APE1 is a master cellular regulator for inflammatory pain condition. <i>IBRO Reports</i> , 2019 , 7, 52-53	2	2

(2015-2019)

152	Human AP-endonuclease (Ape1) activity on telomeric G4 structures is modulated by acetylatable lysine residues in the N-terminal sequence. <i>DNA Repair</i> , 2019 , 73, 129-143	4.3	24	
151	The regulatory role of APE1 in epithelial-to-mesenchymal transition and in determining EGFR-TKI responsiveness in non-small-cell lung cancer. <i>Cancer Medicine</i> , 2018 , 7, 4406-4419	4.8	17	
150	APE1/Ref-1 redox function contributes to inflammatory pain sensitization. <i>Experimental Neurology</i> , 2018 , 307, 1-11	5.7	8	
149	Unveiling the non-repair face of the Base Excision Repair pathway in RNA processing: A missing link between DNA repair and gene expression?. <i>DNA Repair</i> , 2017 , 56, 65-74	4.3	48	
148	Mammalian APE1 controls miRNA processing and its interactome is linked to cancer RNA metabolism. <i>Nature Communications</i> , 2017 , 8, 797	17.4	48	
147	DNA repair enzyme APE1 from evolutionarily ancient Hydra reveals redox activity exclusively found in mammalian APE1. <i>DNA Repair</i> , 2017 , 59, 44-56	4.3	6	
146	Abasic and oxidized ribonucleotides embedded in DNA are processed by human APE1 and not by RNase H2. <i>Nucleic Acids Research</i> , 2017 , 45, 11193-11212	20.1	29	
145	[Letter to the Editor] Isolation of mitochondria is necessary for precise quantification of mitochondrial DNA damage in human carcinoma samples. <i>BioTechniques</i> , 2017 , 62, 13-17	2.5	3	
144	Cardiac Cell Senescence and Redox Signaling. Frontiers in Cardiovascular Medicine, 2017, 4, 38	5.4	13	
143	The Abasic Endonuclease APE1: Much more than a DNA Repair Enzyme 2017 , 219-251		2	
142	Monoclonal antibodies against pools of mono- and polyacetylated peptides selectively recognize acetylated lysines within the context of the original antigen. <i>MAbs</i> , 2016 , 8, 1575-1589	6.6	3	
141	Inhibitors of the apurinic/apyrimidinic endonuclease 1 (APE1)/nucleophosmin (NPM1) interaction that display anti-tumor properties. <i>Molecular Carcinogenesis</i> , 2016 , 55, 688-704	5	48	
140	Placental aging and oxidation damage in a tissue micro-array model: an immunohistochemistry study. <i>Histochemistry and Cell Biology</i> , 2016 , 146, 191-204	2.4	26	
139	Destabilisation, aggregation, toxicity and cytosolic mislocalisation of nucleophosmin regions associated with acute myeloid leukemia. <i>Oncotarget</i> , 2016 , 7, 59129-59143	3.3	35	
138	Serum APE1 as a predictive marker for platinum-based chemotherapy of non-small cell lung cancer patients. <i>Oncotarget</i> , 2016 , 7, 77482-77494	3.3	21	
137	APE1 polymorphic variants cause persistent genomic stress and affect cancer cell proliferation. <i>Oncotarget</i> , 2016 , 7, 26293-306	3.3	21	
136	Isoforms of the Erythropoietin receptor in dopaminergic neurons of the Substantia Nigra. <i>Journal of Neurochemistry</i> , 2016 , 139, 596-609	6	8	
135	A review on protein-protein interaction network of APE1/Ref-1 and its associated biological functions. <i>Cell Biochemistry and Function</i> , 2015 , 33, 101-12	4.2	41	

134	Mitochondrial translocation of APE1 relies on the MIA pathway. Nucleic Acids Research, 2015, 43, 5451-6	4 0.1	42
133	Transcriptional Up-Regulation of APE1/Ref-1 in Hepatic Tumor: Role in Hepatocytes Resistance to Oxidative Stress and Apoptosis. <i>PLoS ONE</i> , 2015 , 10, e0143289	3.7	19
132	Impairment of enzymatic antioxidant defenses is associated with bilirubin-induced neuronal cell death in the cerebellum of Ugt1 KO mice. <i>Cell Death and Disease</i> , 2015 , 6, e1739	9.8	21
131	Cadmium treatment suppresses DNA polymerase latalytic subunit gene expression by acting on the p53 and Sp1 regulatory axis. <i>DNA Repair</i> , 2015 , 35, 90-105	4.3	17
130	The prion protein is critical for DNA repair and cell survival after genotoxic stress. <i>Nucleic Acids Research</i> , 2015 , 43, 904-16	20.1	33
129	Functional regulation of the apurinic/apyrimidinic endonuclease 1 by nucleophosmin: impact on tumor biology. <i>Oncogene</i> , 2014 , 33, 2876-87	9.2	39
128	A short review on the implications of base excision repair pathway for neurons: relevance to neurodegenerative diseases. <i>Mitochondrion</i> , 2014 , 16, 38-49	4.9	39
127	Expression and prognostic significance of APE1/Ref-1 and NPM1 proteins in high-grade ovarian serous cancer. <i>American Journal of Clinical Pathology</i> , 2014 , 141, 404-14	1.9	27
126	Base excision repair in Archaea: back to the future in DNA repair. DNA Repair, 2014, 21, 148-57	4.3	31
125	G-quadruplex DNA recognition by nucleophosmin: new insights from protein dissection. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2014 , 1840, 2050-9	4	44
124	The redox function of APE1 is involved in the differentiation process of stem cells toward a neuronal cell fate. <i>PLoS ONE</i> , 2014 , 9, e89232	3.7	25
123	SIRT1 gene expression upon genotoxic damage is regulated by APE1 through nCaRE-promoter elements. <i>Molecular Biology of the Cell</i> , 2014 , 25, 532-47	3.5	61
122	Nucleophosmin modulates stability, activity, and nucleolar accumulation of base excision repair proteins. <i>Molecular Biology of the Cell</i> , 2014 , 25, 1641-52	3.5	49
121	Emerging roles of the nucleolus in regulating the DNA damage response: the noncanonical DNA repair enzyme APE1/Ref-1 as a paradigmatical example. <i>Antioxidants and Redox Signaling</i> , 2014 , 20, 621-	- <mark>§</mark> 9	66
120	Alterations in the redox state and liver damage: hints from the EASL Basic School of Hepatology. Journal of Hepatology, 2013 , 58, 365-74	13.4	40
119	Role of the unstructured N-terminal domain of the hAPE1 (human apurinic/apyrimidinic endonuclease 1) in the modulation of its interaction with nucleic acids and NPM1 (nucleophosmin). <i>Biochemical Journal</i> , 2013 , 452, 545-57	3.8	31
118	Role of mutual interactions in the chemical and thermal stability of nucleophosmin NPM1 domains. <i>Biochemical and Biophysical Research Communications</i> , 2013 , 430, 523-8	3.4	17
117	Combining RNAi and in vivo confocal microscopy analysis of the photoconvertible fluorescent protein Dendra2 to study a DNA repair protein. <i>BioTechniques</i> , 2013 , 55, 198-203	2.5	6

(2009-2013)

116	Specific inhibition of the redox activity of ape1/ref-1 by e3330 blocks tnf-Induced activation of IL-8 production in liver cancer cell lines. <i>PLoS ONE</i> , 2013 , 8, e70909	3.7	36
115	Nucleolar accumulation of APE1 depends on charged lysine residues that undergo acetylation upon genotoxic stress and modulate its BER activity in cells. <i>Molecular Biology of the Cell</i> , 2012 , 23, 4079-96	3.5	73
114	Acetylation on critical lysine residues of Apurinic/apyrimidinic endonuclease 1 (APE1) in triple negative breast cancers. <i>Biochemical and Biophysical Research Communications</i> , 2012 , 424, 34-9	3.4	29
113	Human AP endonuclease/redox factor APE1/ref-1 modulates mitochondrial function after oxidative stress by regulating the transcriptional activity of NRF1. <i>Free Radical Biology and Medicine</i> , 2012 , 53, 237-48	7.8	29
112	Expression of Dicer and Drosha in triple-negative breast cancer. <i>Journal of Clinical Pathology</i> , 2012 , 65, 320-6	3.9	34
111	Nucleophosmin delocalization in thyroid tumour cells. <i>Endocrine Pathology</i> , 2011 , 22, 18-23	4.2	15
110	Shotgun proteomics analysis reveals new unsuspected molecular effectors of nitrogen-containing bisphosphonates in osteocytes. <i>Journal of Proteomics</i> , 2011 , 74, 1113-22	3.9	6
109	Knock-in reconstitution studies reveal an unexpected role of Cys-65 in regulating APE1/Ref-1 subcellular trafficking and function. <i>Molecular Biology of the Cell</i> , 2011 , 22, 3887-901	3.5	50
108	Critical lysine residues within the overlooked N-terminal domain of human APE1 regulate its biological functions. <i>Nucleic Acids Research</i> , 2010 , 38, 8239-56	20.1	83
107	Intrusion of a DNA repair protein in the RNome world: is this the beginning of a new era?. <i>Molecular and Cellular Biology</i> , 2010 , 30, 366-71	4.8	70
106	BRCA1 modulates the expression of hnRNPA2B1 and KHSRP. Cell Cycle, 2010, 9, 4666-73	4.7	17
105	Mass spectrometry approaches for the redox characterization of protein cysteine residues the case of the transcription factor Pax-8. <i>Methods in Enzymology</i> , 2010 , 473, 227-50	1.7	1
104	Nucleophosmin is overexpressed in thyroid tumors. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 397, 499-504	3.4	44
103	Two novel NPM1 mutations in a therapy-responder AML patient. <i>Hematological Oncology</i> , 2010 , 28, 151	-5 .3	6
102	Targeting DNA repair proteins for cancer treatment. Cellular and Molecular Life Sciences, 2010, 67, 3569	-72 .3	21
101	Understanding different functions of mammalian AP endonuclease (APE1) as a promising tool for cancer treatment. <i>Cellular and Molecular Life Sciences</i> , 2010 , 67, 3589-608	10.3	94
100	A proteomic approach to the bilirubin-induced toxicity in neuronal cells reveals a protective function of DJ-1 protein. <i>Proteomics</i> , 2010 , 10, 1645-57	4.8	26
99	APE1/Ref-1 interacts with NPM1 within nucleoli and plays a role in the rRNA quality control process. <i>Molecular and Cellular Biology</i> , 2009 , 29, 1834-54	4.8	169

98	Thyroid tumors: novel insights from proteomic studies. Expert Review of Proteomics, 2009, 6, 363-76	4.2	8
97	DeltaNp73alpha inhibits PTEN expression in thyroid cancer cells. <i>International Journal of Cancer</i> , 2009 , 124, 2539-48	7.5	30
96	Down-regulation of SM22/transgelin gene expression during H9c2 cells differentiation. <i>Molecular and Cellular Biochemistry</i> , 2009 , 327, 145-52	4.2	9
95	Genome-wide analysis and proteomic studies reveal APE1/Ref-1 multifunctional role in mammalian cells. <i>Proteomics</i> , 2009 , 9, 1058-74	4.8	79
94	Modern strategies to identify new molecular targets for the treatment of liver diseases: The promising role of Proteomics and Redox Proteomics investigations. <i>Proteomics - Clinical Applications</i> , 2009 , 3, 242-62	3.1	9
93	Unexpected phenotype of a typical NPM1 mutant. British Journal of Haematology, 2009, 147, 760-3	4.5	4
92	Aggresome-forming TTRAP mediates pro-apoptotic properties of Parkinson's disease-associated DJ-1 missense mutations. <i>Cell Death and Differentiation</i> , 2009 , 16, 428-38	12.7	49
91	Transcriptome and proteome analysis of osteocytes treated with nitrogen-containing bisphosphonates. <i>Journal of Proteome Research</i> , 2009 , 8, 1131-42	5.6	20
90	Differential proteomic analysis of subfractioned human hepatocellular carcinoma tissues. <i>Journal of Proteome Research</i> , 2009 , 8, 2273-84	5.6	13
89	APE1/Ref-1 in Alzheimer's disease: an immunohistochemical study. <i>Neuroscience Letters</i> , 2009 , 466, 124	· -3 .3	27
88	The many functions of APE1/Ref-1: not only a DNA repair enzyme. <i>Antioxidants and Redox Signaling</i> , 2009 , 11, 601-20	8.4	360
87	Identification of secondary targets of N-containing bisphosphonates in mammalian cells via parallel competition analysis of the barcoded yeast deletion collection. <i>Genome Biology</i> , 2009 , 10, R93	18.3	21
86	Redox state, oxidative stress, and molecular mechanisms of protective and toxic effects of bilirubin on cells. <i>Current Pharmaceutical Design</i> , 2009 , 15, 2908-14	3.3	23
85	H(2)O(2) modulates purinergic-dependent calcium signalling in osteoblast-like cells. <i>Cell Calcium</i> , 2008 , 43, 457-68	4	7
84	Molecular basis and mechanisms of progression of non-alcoholic steatohepatitis. <i>Trends in Molecular Medicine</i> , 2008 , 14, 72-81	11.5	324
83	The solution structure of DNA-free Pax-8 paired box domain accounts for redox regulation of		18
	transcriptional activity in the pax protein family. <i>Journal of Biological Chemistry</i> , 2008 , 283, 33321-8	5.4	
82	The neutrophil gelatinase-associated lipocalin (NGAL), a NF-kappaB-regulated gene, is a survival factor for thyroid neoplastic cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 14058-63	5.4	154

(2006-2008)

80	Molecular and functional changes associated with APE1/Ref-1 loss of expression reveal its multifunctional nature in mammalian cells. <i>FASEB Journal</i> , 2008 , 22, 289-289	0.9	
79	Oxidized transthyretin in amniotic fluid as an early marker of preeclampsia. <i>Journal of Proteome Research</i> , 2007 , 6, 160-70	5.6	60
78	RbAp48 is a target of nuclear factor-kappaB activity in thyroid cancer. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 1458-66	5.6	29
77	Subcellular localization of APE1/Ref-1 in human hepatocellular carcinoma: possible prognostic significance. <i>Molecular Medicine</i> , 2007 , 13, 89-96	6.2	79
76	Altered intracellular redox status in Gaucher disease fibroblasts and impairment of adaptive response against oxidative stress. <i>Journal of Cellular Physiology</i> , 2007 , 212, 223-35	7	39
75	Bilirubin-induced cell toxicity involves PTEN activation through an APE1/Ref-1-dependent pathway. <i>Journal of Molecular Medicine</i> , 2007 , 85, 1099-112	5.5	38
74	Structural studies on Pax-8 Prd domain/DNA complex. <i>Journal of Biomolecular Structure and Dynamics</i> , 2007 , 24, 429-41	3.6	4
73	Oxidative stress causes bone loss in estrogen-deficient mice through enhanced bone marrow dendritic cell activation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 15087-92	11.5	117
72	Redox proteomics and immunohistology to study molecular events during ischemia-reperfusion in human liver. <i>Transplantation Proceedings</i> , 2007 , 39, 1755-60	1.1	21
71	Detection of aromatic catabolic gene expression in heterogeneous organic matter used for reduction of volatile organic compounds (VOC) by biofiltration. <i>Biotechnology Letters</i> , 2007 , 29, 65-72	3	3
70	Biocrystallographic study on drug resistant variants of HIV protease with new inhibitors. <i>FASEB Journal</i> , 2007 , 21, A1011	0.9	
69	Nucleotide receptors stimulation by extracellular ATP controls Hsp90 expression through APE1/Ref-1 in thyroid cancer cells: a novel tumorigenic pathway. <i>Journal of Cellular Physiology</i> , 2006 , 209, 44-55	7	17
68	A potent HIV protease inhibitor identified in an epimeric mixture by high-resolution protein crystallography. <i>ChemMedChem</i> , 2006 , 1, 186-8	3.7	5
67	Early Molecular Events during Response to Oxidative Stress in Human Cells by Differential Proteomics 2006 , 369-397		3
66	Bisphosphonates activate nucleotide receptors signaling and induce the expression of Hsp90 in osteoblast-like cell lines. <i>Bone</i> , 2006 , 39, 739-53	4.7	24
65	Proteomic analysis of liver tissues subjected to early ischemia/reperfusion injury during human orthotopic liver transplantation. <i>Proteomics</i> , 2006 , 6, 3455-65	4.8	49
64	Molecular analysis of a human PAX6 homeobox mutant. <i>European Journal of Human Genetics</i> , 2006 , 14, 744-51	5.3	7
63	Differential proteomic analysis of nuclear extracts from thyroid cell lines. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2006 , 833, 41-50	3.2	16

62	Activation of APE1/Ref-1 is dependent on reactive oxygen species generated after purinergic receptor stimulation by ATP. <i>Nucleic Acids Research</i> , 2005 , 33, 4379-94	20.1	108
61	Cross-regulation between Egr-1 and APE/Ref-1 during early response to oxidative stress in the human osteoblastic HOBIT cell line: evidence for an autoregulatory loop. <i>Free Radical Research</i> , 2005 , 39, 269-81	4	39
60	Autocrine/paracrine stimulation of purinergic receptors in osteoblasts: contribution of vesicular ATP release. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 331, 1429-38	3.4	52
59	Extracellular nucleotides activate Runx2 in the osteoblast-like HOBIT cell line: a possible molecular link between mechanical stress and osteoblastsSresponse. <i>Bone</i> , 2005 , 36, 418-32	4.7	65
58	PTEN and Egr-1 expression in thyroid proliferative lesions. <i>Cancer Letters</i> , 2005 , 224, 105-9	9.9	13
57	The intracellular localization of APE1/Ref-1: more than a passive phenomenon?. <i>Antioxidants and Redox Signaling</i> , 2005 , 7, 367-84	8.4	311
56	Overoxidation of peroxiredoxins as an immediate and sensitive marker of oxidative stress in HepG2 cells and its application to the redox effects induced by ischemia/reperfusion in human liver. <i>Free Radical Research</i> , 2005 , 39, 255-68	4	54
55	Proteins as biomarkers of oxidative/nitrosative stress in diseases: the contribution of redox proteomics. <i>Mass Spectrometry Reviews</i> , 2005 , 24, 55-99	11	354
54	Functional analysis of a novel RUNX2 missense mutation found in a family with cleidocranial dysplasia. <i>Journal of Human Genetics</i> , 2005 , 50, 679-83	4.3	11
53	Activation of human T lymphocytes under conditions similar to those that occur during exposure to microgravity: a proteomics study. <i>Proteomics</i> , 2005 , 5, 1827-37	4.8	34
52	A differential proteomic approach to identify proteins associated with thyroid cell transformation. Journal of Molecular Endocrinology, 2005 , 34, 199-207	4.5	15
51	Effects of histone acetylation on sodium iodide symporter promoter and expression of thyroid-specific transcription factors. <i>Endocrinology</i> , 2005 , 146, 3967-74	4.8	68
50	Expression, regulation, and function of paired-box gene 8 in the human placenta and placental cancer cell lines. <i>Endocrinology</i> , 2005 , 146, 4009-15	4.8	16
49	The importance of redox state in liver damage. <i>Annals of Hepatology</i> , 2004 , 3, 86-92	3.1	139
48	Control of phosphatase and tensin homolog (PTEN) gene expression in normal and neoplastic thyroid cells. <i>Endocrinology</i> , 2004 , 145, 4660-6	4.8	19
47	Transcriptional regulation of human sodium/iodide symporter gene: a role for redox factor-1. <i>Endocrinology</i> , 2004 , 145, 1290-3	4.8	18
46	Proteomic evaluation of core biopsy specimens from breast lesions. <i>Cancer Letters</i> , 2004 , 204, 79-86	9.9	25
45	Galectin-3 expression in non-small cell lung carcinoma. <i>Cancer Letters</i> , 2004 , 212, 233-9	9.9	35

(2001-2004)

44	Functional interaction among thyroid-specific transcription factors: Pax8 regulates the activity of Hex promoter. <i>Molecular and Cellular Endocrinology</i> , 2004 , 214, 117-25	4.4	21
43	A proteomic approach to identify early molecular targets of oxidative stress in human epithelial lens cells. <i>Biochemical Journal</i> , 2004 , 378, 929-37	3.8	85
42	The importance of redox state in liver damage. <i>Annals of Hepatology</i> , 2004 , 3, 86-92	3.1	44
41	Extracellular ATP stimulates the early growth response protein 1 (Egr-1) via a protein kinase C-dependent pathway in the human osteoblastic HOBIT cell line. <i>Biochemical Journal</i> , 2003 , 373, 815-24	3.8	35
40	The PAX6 gene is activated by the basic helix-loop-helix transcription factor NeuroD/BETA2. <i>Biochemical Journal</i> , 2003 , 376, 707-15	3.8	26
39	Identification of different isoforms of eEF1A in the nuclear fraction of human T-lymphoblastic cancer cell line specifically binding to aptameric cytotoxic GT oligomers. <i>FEBS Journal</i> , 2003 , 270, 3251-6	62	27
38	Probing the dimeric structure of porcine aminoacylase 1 by mass spectrometric and modeling procedures. <i>Biochemistry</i> , 2003 , 42, 4430-43	3.2	41
37	Cytosine-block telomeric type DNA-binding activity of hnRNP proteins from human cell lines. <i>Archives of Biochemistry and Biophysics</i> , 2003 , 409, 305-14	4.1	26
36	Nuclear localization of Galectin-3 in transformed thyroid cells: a role in transcriptional regulation. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 302, 545-53	3.4	89
35	H(2)O(2) induces translocation of APE/Ref-1 to mitochondria in the Raji B-cell line. <i>Journal of Cellular Physiology</i> , 2002 , 193, 180-6	7	84
34	Expression and localization of the homeodomain-containing protein HEX in human thyroid tumors. Journal of Clinical Endocrinology and Metabolism, 2002 , 87, 1376-83	5.6	34
33	APE/Ref-1 is controlled by both redox and cAMP-dependent mechanisms in rat thyroid cells. <i>Hormone and Metabolic Research</i> , 2002 , 34, 303-10	3.1	7
32	Redox effector factor-1 regulates the activity of thyroid transcription factor 1 by controlling the redox state of the N transcriptional activation domain. <i>Journal of Biological Chemistry</i> , 2002 , 277, 14564	ı- 5 4	29
31	APE/Ref-1 is increased in nuclear fractions of human thyroid hyperfunctioning nodules. <i>Molecular and Cellular Endocrinology</i> , 2002 , 194, 71-6	4.4	6
30	Expression and Localization of the Homeodomain-Containing Protein HEX in Human Thyroid Tumors. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2002 , 87, 1376-1383	5.6	26
29	Prognostic role of Ape/Ref-1 subcellular expression in stage I-III breast carcinomas. <i>Oncology Reports</i> , 2002 , 9, 11-7	3.5	47
28	A chicken hnRNP of the A/B family recognizes the single-stranded d(CCCTAA)(n) telomeric repeated motif. <i>FEBS Journal</i> , 2001 , 268, 139-48		14
27	Missense mutations of human homeoboxes: A review. <i>Human Mutation</i> , 2001 , 18, 361-74	4.7	53

26	A unique combination of transcription factors controls differentiation of thyroid cells. <i>Progress in Molecular Biology and Translational Science</i> , 2001 , 66, 307-56		148
25	Mitochondrial localization of APE/Ref-1 in thyroid cells. <i>Mutation Research DNA Repair</i> , 2001 , 485, 143-	52	67
24	Monoclonal antibody detection of naphthalene dioxygenase from Pseudomonas aeruginosa 2NR. <i>Letters in Applied Microbiology</i> , 2000 , 31, 313-8	2.9	3
23	Calreticulin enhances the transcriptional activity of thyroid transcription factor-1 by binding to its homeodomain. <i>Journal of Biological Chemistry</i> , 1999 , 274, 4640-5	5.4	31
22	Adhesion to fibronectin promotes the activation of the p125(FAK)/Zap-70complex in human T cells. <i>Immunology</i> , 1999 , 98, 564-8	7.8	13
21	Molecular characterization of Pseudomonas aeruginosa 2NR degrading naphthalene. <i>Letters in Applied Microbiology</i> , 1999 , 29, 181-6	2.9	11
20	Comparative stability analysis of the thyroid transcription factor 1 and Antennapedia homeodomains: evidence for residue 54 in controlling the structural stability of the recognition helix. <i>International Journal of Biochemistry and Cell Biology</i> , 1999 , 31, 1339-53	5.6	5
19	Effect of phosphorothioate modifications on the ability of GTn oligodeoxynucleotides to specifically recognize single-stranded DNA-binding proteins and to affect human cancer cellular growth. <i>Biochimie</i> , 1999 , 81, 1115-22	4.6	8
18	Topology of the thyroid transcription factor 1 homeodomain-DNA complex. <i>Biochemistry</i> , 1999 , 38, 64-	-73 .2	24
17	Co-operation between the PAI and RED subdomains of Pax-8 in the interaction with the thyroglobulin promoter. <i>Biochemical Journal</i> , 1999 , 337, 253-262	3.8	27
16	Structural defects of a Pax8 mutant that give rise to congenital hypothyroidism. <i>Biochemical Journal</i> , 1999 , 341, 89-93	3.8	13
15	Co-operation between the PAI and RED subdomains of Pax-8 in the interaction with the thyroglobulin promoter. <i>Biochemical Journal</i> , 1999 , 337, 253	3.8	12
14	Structural defects of a Pax8 mutant that give rise to congenital hypothyroidism. <i>Biochemical Journal</i> , 1999 , 341, 89	3.8	6
13	Redox potential controls the structure and DNA binding activity of the paired domain. <i>Journal of Biological Chemistry</i> , 1998 , 273, 25062-72	5.4	86
12	Fibronectin binding promotes a PKC-dependent modulation of NF-kappa B in human T cells. <i>Biochemical and Biophysical Research Communications</i> , 1998 , 243, 732-7	3.4	20
11	Ref-1 controls pax-8 DNA-binding activity. <i>Biochemical and Biophysical Research Communications</i> , 1998 , 252, 178-83	3.4	80
10	Pax-8 protein levels regulate thyroglobulin gene expression. <i>Journal of Molecular Endocrinology</i> , 1998 , 21, 347-54	4.5	56
9	Structural and functional properties of the N transcriptional activation domain of thyroid transcription factor-1: similarities with the acidic activation domains. <i>Biochemical Journal</i> , 1998 , 329 (Pt 2), 395-403	3.8	30

LIST OF PUBLICATIONS

8	High mobility group I proteins interfere with the homeodomains binding to DNA. <i>Journal of Biological Chemistry</i> , 1997 , 272, 29904-10	5.4	23	
7	Biphasic control of NF-kappa B activation induced by the triggering of HLA-DR antigens expressed on B cells. <i>Cytokine</i> , 1997 , 9, 295-9	4	4	
6	Functional interference between contacting amino acids of homeodomains. <i>FEBS Letters</i> , 1997 , 407, 320-4	3.8	15	
5	Hydrogen-deuterium exchange studies of the rat thyroid transcription factor 1 homeodomain. <i>Journal of Biomolecular NMR</i> , 1997 , 9, 397-407	3	1	
4	Analysis of the solution structure of the homeodomain of rat thyroid transcription factor 1 by 1H-NMR spectroscopy and restrained molecular mechanics. <i>FEBS Journal</i> , 1996 , 241, 101-13		24	
3	Definition of the DNA-binding specificity of TTF-1 homeodomain by chromatographic selection of binding sequences. <i>Biochemical and Biophysical Research Communications</i> , 1995 , 213, 781-8	3.4	17	
2	Expression, purification and functional characterisation of a Kunitz-type module from chicken type VI collagen. <i>Biochemical and Biophysical Research Communications</i> , 1995 , 215, 1050-5	3.4	2	
1	Effect of salt concentration on TTF-1 HD binding to specific and non-specific DNA sequences. <i>Biochemical and Biophysical Research Communications</i> , 1993 , 197, 632-8	3.4	11	