

Subbaratnam Muthukrishnan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1846963/publications.pdf>

Version: 2024-02-01

62
papers

4,108
citations

101384

36
h-index

138251

58
g-index

62
all docs

62
docs citations

62
times ranked

2948
citing authors

#	ARTICLE	IF	CITATIONS
1	Chitin in insect cuticle. <i>Advances in Insect Physiology</i> , 2022, , 1-110.	1.1	5
2	AA15 lytic polysaccharide monooxygenase is required for efficient chitinous cuticle turnover during insect molting. <i>Communications Biology</i> , 2022, 5, .	2.0	10
3	Chitin deacetylases are necessary for insect femur muscle attachment and mobility. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, .	3.3	2
4	Yellow-y Functions in Egg Melanization and Chorion Morphology of the Asian Tiger Mosquito, <i>Aedes albopictus</i> . <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 769788.	1.8	10
5	Gene functions in adult cuticle pigmentation of the yellow mealworm, <i>Tenebrio molitor</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2020, 117, 103291.	1.2	37
6	Yellow-g and Yellow-g2 proteins are required for egg desiccation resistance and temporal pigmentation in the Asian tiger mosquito, <i>Aedes albopictus</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2020, 122, 103386.	1.2	46
7	Insect Cuticular Chitin Contributes to Form and Function. <i>Current Pharmaceutical Design</i> , 2020, 26, 3530-3545.	0.9	43
8	Chitin Organizing and Modifying Enzymes and Proteins Involved In Remodeling of the Insect Cuticle. <i>Advances in Experimental Medicine and Biology</i> , 2019, 1142, 83-114.	0.8	37
9	Future questions in insect chitin biology: A microreview. <i>Archives of Insect Biochemistry and Physiology</i> , 2018, 98, e21454.	0.6	14
10	Group I chitin deacetylases are essential for higher order organization of chitin fibers in beetle cuticle. <i>Journal of Biological Chemistry</i> , 2018, 293, 6985-6995.	1.6	34
11	A chitinase with two catalytic domains is required for organization of the cuticular extracellular matrix of a beetle. <i>PLoS Genetics</i> , 2018, 14, e1007307.	1.5	46
12	Functional analysis of TcCTLP-5C2, a chymotrypsin-like serine protease needed for molting in <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2017, 86, 20-28.	1.2	2
13	Development and ultrastructure of the rigid dorsal and flexible ventral cuticles of the elytron of the red flour beetle, <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2017, 91, 21-33.	1.2	36
14	Cuticle formation and pigmentation in beetles. <i>Current Opinion in Insect Science</i> , 2016, 17, 1-9.	2.2	125
15	Multifaceted biological insights from a draft genome sequence of the tobacco hornworm moth, <i>Manduca sexta</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2016, 76, 118-147.	1.2	154
16	Chitin Metabolic Pathways in Insects and Their Regulation. , 2016, , 31-65.		12
17	Arylalkylamine N-acetyltransferase 1 gene (TcAANAT1) is required for cuticle morphology and pigmentation of the adult red flour beetle, <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2016, 79, 119-129.	1.2	39
18	Biosynthesis, Turnover, and Functions of Chitin in Insects. <i>Annual Review of Entomology</i> , 2016, 61, 177-196.	5.7	291

#	ARTICLE	IF	CITATIONS
19	Overview of chitin metabolism enzymes in <i>Manduca sexta</i> : Identification, domain organization, phylogenetic analysis and gene expression. <i>Insect Biochemistry and Molecular Biology</i> , 2015, 62, 114-126.	1.2	95
20	Cuticular protein with a low complexity sequence becomes cross-linked during insect cuticle sclerotization and is required for the adult molt. <i>Scientific Reports</i> , 2015, 5, 10484.	1.6	67
21	<i>Tribolium castaneum</i> RR-1 Cuticular Protein TcCPR4 Is Required for Formation of Pore Canals in Rigid Cuticle. <i>PLoS Genetics</i> , 2015, 11, e1004963.	1.5	69
22	Knickkopf and retroactive proteins are required for formation of laminar serosal procuticle during embryonic development of <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2015, 60, 1-6.	1.2	22
23	Analysis of chitin-binding proteins from <i>Manduca sexta</i> provides new insights into evolution of peritrophin A-type chitin-binding domains in insects. <i>Insect Biochemistry and Molecular Biology</i> , 2015, 62, 127-141.	1.2	88
24	Chitin is a necessary component to maintain the barrier function of the peritrophic matrix in the insect midgut. <i>Insect Biochemistry and Molecular Biology</i> , 2015, 56, 21-28.	1.2	104
25	Functional Specialization Among Members Of Knickkopf Family Of Proteins In Insect Cuticle Organization. <i>PLoS Genetics</i> , 2014, 10, e1004537.	1.5	19
26	A Major Facilitator Superfamily protein encoded by TcMuck gene is not required for cuticle pigmentation, growth and development in <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2014, 49, 43-48.	1.2	2
27	Two major cuticular proteins are required for assembly of horizontal laminae and vertical pore canals in rigid cuticle of <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2014, 53, 22-29.	1.2	76
28	Two essential peritrophic matrix proteins mediate matrix barrier functions in the insect midgut. <i>Insect Biochemistry and Molecular Biology</i> , 2014, 49, 24-34.	1.2	82
29	Gene Families of Cuticular Proteins Analogous to Peritrophins (CPAPs) in <i>Tribolium castaneum</i> Have Diverse Functions. <i>PLoS ONE</i> , 2012, 7, e49844.	1.1	95
30	Knickkopf protein protects and organizes chitin in the newly synthesized insect exoskeleton. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 17028-17033.	3.3	106
31	Cuticle tanning in <i>Tribolium castaneum</i> . <i>Entomological Research</i> , 2011, 41, 293-293.	0.6	1
32	Expression Profiles and Functional Analysis of Genes Encoding Chitin Deacetylases, Extracellular Matrix-Modifying Proteins in <i>Tribolium castaneum</i> . <i>Entomological Research</i> , 2011, 41, 294-294.	0.6	1
33	Functional Analysis of Genes of Chitin Metabolism in <i>Tribolium castaneum</i> by RNA interference. <i>Entomological Research</i> , 2011, 41, 295-295.	0.6	0
34	RNAi-based functional analysis of yellow-e in <i>Tribolium castaneum</i> . <i>Entomological Research</i> , 2011, 41, 296-296.	0.6	0
35	Two Major Structural Proteins Are Required for Rigid Adult Cuticle Formation in the Red Flour Beetle, <i>Tribolium castaneum</i> . <i>Entomological Research</i> , 2011, 41, 297-297.	0.6	0
36	Why a Membrane-Anchored Chitinase (CHT7) in <i>Tribolium</i> ?. <i>Entomological Research</i> , 2011, 41, 298-298.	0.6	1

#	ARTICLE	IF	CITATIONS
37	The effect of lufenuron, a chitin synthesis inhibitor, on oogenesis of <i>Rhodnius prolixus</i> . <i>Pesticide Biochemistry and Physiology</i> , 2010, 98, 59-67.	1.6	33
38	Oxalic acid-induced resistance to <i>Rhizoctonia solani</i> in rice is associated with induction of phenolics, peroxidase and pathogenesis-related proteins. <i>Journal of Plant Interactions</i> , 2010, 5, 147-157.	1.0	41
39	Chymotrypsin-like peptidases from <i>Tribolium castaneum</i> : A role in molting revealed by RNA interference. <i>Insect Biochemistry and Molecular Biology</i> , 2010, 40, 274-283.	1.2	49
40	Genes encoding proteins with peritrophin A-type chitin-binding domains in <i>Tribolium castaneum</i> are grouped into three distinct families based on phylogeny, expression and function. <i>Insect Biochemistry and Molecular Biology</i> , 2010, 40, 214-227.	1.2	141
41	Molecular and Functional Analyses of Amino Acid Decarboxylases Involved in Cuticle Tanning in <i>Tribolium castaneum</i> . <i>Journal of Biological Chemistry</i> , 2009, 284, 16584-16594.	1.6	181
42	Analysis of functions of the chitin deacetylase gene family in <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2009, 39, 355-365.	1.2	145
43	Domain organization and phylogenetic analysis of the chitinase-like family of proteins in three species of insects. <i>Insect Biochemistry and Molecular Biology</i> , 2008, 38, 452-466.	1.2	129
44	Characterization and expression of the β -N-acetylhexosaminidase gene family of <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2008, 38, 478-489.	1.2	84
45	Domain organization and phylogenetic analysis of proteins from the chitin deacetylase gene family of <i>Tribolium castaneum</i> and three other species of insects. <i>Insect Biochemistry and Molecular Biology</i> , 2008, 38, 440-451.	1.2	130
46	Chitin synthases are required for survival, fecundity and egg hatch in the red flour beetle, <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2008, 38, 959-962.	1.2	145
47	Functional specialization among insect chitinase family genes revealed by RNA interference. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 6650-6655.	3.3	221
48	A chitin-like component in <i>Aedes aegypti</i> eggshells, eggs and ovaries. <i>Insect Biochemistry and Molecular Biology</i> , 2007, 37, 1249-1261.	1.2	94
49	Chitin synthase genes in <i>Manduca sexta</i> : characterization of a gut-specific transcript and differential tissue expression of alternately spliced mRNAs during development. <i>Insect Biochemistry and Molecular Biology</i> , 2005, 35, 529-540.	1.2	110
50	Sequences of cDNAs and expression of genes encoding chitin synthase and chitinase in the midgut of <i>Spodoptera frugiperda</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2005, 35, 1249-1259.	1.2	89
51	Characterization of two chitin synthase genes of the red flour beetle, <i>Tribolium castaneum</i> , and alternate exon usage in one of the genes during development. <i>Insect Biochemistry and Molecular Biology</i> , 2004, 34, 291-304.	1.2	167
52	Sequence of a cDNA and expression of the gene encoding a putative epidermal chitin synthase of <i>Manduca sexta</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2002, 32, 1497-1506.	1.2	76
53	Insect resistance of transgenic tobacco expressing an insect chitinase gene. <i>Transgenic Research</i> , 1998, 7, 77-84.	1.3	151
54	Induction of thaumatin-like proteins (TLPs) in <i>Rhizoctonia solani</i> -infected rice and characterization of two new cDNA clones. <i>Physiologia Plantarum</i> , 1998, 102, 21-28.	2.6	61

#	ARTICLE	IF	CITATIONS
55	Induction of chitinases and beta-1,3-glucanases in <i>Rhizoctonia solani</i> -infected rice plants: Isolation of an infection-related chitinase cDNA clone. <i>Physiologia Plantarum</i> , 1996, 97, 39-46.	2.6	43
56	Variability in antifungal proteins in the grains of maize, sorghum and wheat. <i>Physiologia Plantarum</i> , 1993, 88, 339-349.	2.6	30
57	A novel cereal storage protein: molecular genetics of the 19 kDa globulin of rice. <i>Plant Molecular Biology</i> , 1992, 18, 151-154.	2.0	43
58	Nucleotide sequence of a cDNA clone that encodes the maize inhibitor of trypsin and activated Hageman factor. <i>Plant Molecular Biology</i> , 1992, 18, 813-814.	2.0	23
59	Nucleotide sequence of a cloned rice genomic DNA fragment that encodes a 10 kDa prolamin polypeptide. <i>Nucleic Acids Research</i> , 1990, 18, 683-683.	6.5	14
60	Nucleotide and predicted amino acid sequences of two different genes for high-pI α -amylases from barley. <i>Plant Molecular Biology</i> , 1989, 12, 119-121.	2.0	24
61	Identification of an endochitinase cDNA clone from barley aleurone cells. <i>Plant Molecular Biology</i> , 1989, 12, 403-412.	2.0	44
62	Structure and organization of two divergent α -amylase genes from barley. <i>Plant Molecular Biology</i> , 1987, 9, 3-17.	2.0	69