

# William Severi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1856034/publications.pdf>

Version: 2024-02-01

62

papers

856

citations

623734

14

h-index

552781

26

g-index

62

all docs

62

docs citations

62

times ranked

1165

citing authors

#	ARTICLE	IF	CITATIONS
1	Growth, red blood cells, and gill alterations of red pacu <i>(</i><scp><i>Piaractus</i></scp>) Tj ETQq1 1 0.784314 rgBT /Overlock biofloc. Journal of the World Aquaculture Society, 2022, 53, 652-668.	2.4	4
2	Assessment of different ionic adjustment strategies in lowâ€salinity water on the growth of <i>Litopenaeus vannamei</i> and microbial community stoichiometry in a symbiotic nursery system. Aquaculture Research, 2022, 53, 50-62.	1.8	9
3	To what degree do spatial and limnological predictors explain the occurrence of a submerged macrophyte species in lotic and semi-lotic/lentic environments of a dammed river?. Limnology, 2021, 22, 101-110.	1.5	2
4	Limnological layers improve species distribution modeling of aquatic macrophytes at fine-spatial resolution. Acta Botanica Brasiliensis, 2021, 35, 9-16.	0.8	1
5	Reduced genetic diversity and the success of the invasive peacock bass (Cichliformes: Cichlidae). Brazilian Journal of Biology, 2021, 84, e248656.	0.9	0
6	Production of <i>Daphnia similis</i> Claus, 1876 using wastewater from tilapia cultivation in a biofloc system. Aquaculture International, 2020, 28, 403-419.	2.2	4
7	Trophic and limnological changes in highly fragmented rivers predict the decreasing abundance of detritivorous fish. Ecological Indicators, 2020, 110, 105933.	6.3	15
8	Effects of anthropic actions and forest areas on a neotropical aquatic ecosystem. Science of the Total Environment, 2019, 691, 367-377.	8.0	12
9	Phytoplankton communities in aquaculture system (integration of shrimp and seaweed). Chemistry and Ecology, 2019, 35, 903-921.	1.6	5
10	Effects of addition of <i>Navicula</i> sp. (diatom) in different densities to postlarvae of shrimp <i>Litopenaeus vannamei</i> reared in a BFT system: Growth, survival, productivity and fatty acid profile. Aquaculture Research, 2019, 50, 2231-2239.	1.8	19
11	Dynamics of early lifeâ€history stages of fish along an estuarineÂgradient. Fisheries Oceanography, 2019, 28, 402-418.	1.7	10
12	Response of aquatic macrophyte biomass to limnological changes under water level fluctuation in tropical reservoirs. Brazilian Journal of Biology, 2019, 79, 120-126.	0.9	8
13	STRUCTURE OF THE FISH ASSEMBLAGE AND FUNCTIONAL GUILDS IN THE ESTUARY OF MARACAÃPE, NORTHEAST COAST OF BRAZIL. Boletim Do Instituto De Pesca, 2019, 45, .	0.5	7
14	TILAPIA CULTIVATED IN A LOW-SALINITY BIOFLOC SYSTEM SUPPLEMENTED WITH &lt;i&gt;Chlorella vulgaris&lt;/i&gt; AND DIFFERENT MOLASSES APPLICATION RATES. Boletim Do Instituto De Pesca, 2019, 45, .	0.5	9
15	Nile tilapia fingerling cultivated in a low-salinity biofloc system at different stocking densities. Spanish Journal of Agricultural Research, 2019, 16, e0612.	0.6	16
16	HÃBITO ALIMENTAR DO BEIJUPIRÃ•EM PERNAMBUCO, NORDESTE DO BRASIL. Boletim Do Instituto De Pesca, 2019, 45, .	0.5	1
17	AnÃ¡lise quantitativa triannual da riqueza Ãactia em funÃ§Ã£o da lua e perÃodos do dia: estudo de caso na zona de arrebentÃ£o, ItamaracÃ¡, Pernambuco. Revista Ibero-americana De CiÃªncias Ambientais, 2019, 10, 302-315.	0.1	0
18	Similarities in correlates of native and introduced fish species richness distribution in Brazilian reservoirs. Hydrobiologia, 2018, 817, 167-177.	2.0	12

#	ARTICLE	IF	CITATIONS
19	Cumulative ecological effects of a Neotropical reservoir cascade across multiple assemblages. <i>Hydrobiologia</i> , 2018, 819, 77-91.	2.0	47
20	Bioremediation of shrimp biofloc wastewater using clam, seaweed and fish. <i>Chemistry and Ecology</i> , 2018, 34, 901-913.	1.6	17
21	Effects of two commercial feeds with high and low crude protein content on the performance of white shrimp <i>Litopenaeus vannamei</i> raised in an integrated biofloc system with the seaweed <i>Gracilaria birdiae</i> . <i>Spanish Journal of Agricultural Research</i> , 2018, 16, e0603.	0.6	9
22	Microsatellite assessment of the genetic diversity in indigenous populations of curimba ( <i>Prochilodus</i> ) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 13	1.1	11
23	Effect of the addition of <i>Chaetoceros calcitrans</i> , <i>Navicula</i> sp. and <i>Phaeodactylum tricornutum</i> (diatoms) on phytoplankton composition and growth of <i>Litopenaeus vannamei</i> (Boone) postlarvae reared in a biofloc system. <i>Aquaculture Research</i> , 2017, 48, 4155-4164.	1.8	21
24	Environmental filters predict the trait composition of fish communities in reservoir cascades. <i>Hydrobiologia</i> , 2017, 802, 245-253.	2.0	64
25	Morphological development of larvae and juveniles of <i>Prochilodus argenteus</i> . <i>Ciencia Rural</i> , 2017, 47, .	0.5	1
26	Bioremediation and biocontrol of commercial probiotic in marine shrimp culture with biofloc. <i>Latin American Journal of Aquatic Research</i> , 2017, 45, 167-176.	0.6	31
27	Novelty on the market, novelty in the environment: The invasion of non-native fish jaguar guapote (Perciformes) in northeastern Brazil. <i>Neotropical Biology and Conservation</i> , 2017, 12, .	0.9	9
28	Effect of the addition of diatoms ( <i>Navicula</i> spp.) and rotifers ( <i>Brachionus plicatilis</i> ) on water quality and growth of the <i>Litopenaeus vannamei</i> postlarvae reared in a biofloc system. <i>Aquaculture Research</i> , 2016, 47, 3990-3997.	1.8	34
29	Water quality, Vibriodensity and growth of Pacific white shrimp <i>Litopenaeus vannamei</i> (Boone) in an integrated biofloc system with red seaweed <i>Gracilaria birdiae</i> (Greville). <i>Aquaculture Research</i> , 2016, 47, 940-950.	1.8	34
30	Distribution of benthic macroinvertebrates in a tropical reservoir cascade. <i>Hydrobiologia</i> , 2016, 765, 265-275.	2.0	23
31	Integrated approach to the understanding of the degradation of an urban river: local perceptions, environmental parameters and geoprocessing. <i>Journal of Ethnobiology and Ethnomedicine</i> , 2015, 11, 69.	2.6	14
32	Early development of two tropical fishes (Perciformes: Sciaenidae) from the Pantanal of Mato Grosso, Brazil. <i>Revista De Biologia Tropical</i> , 2015, 63, 1105.	0.4	3
33	Estrutura da assembleia ictioplânctonica em dois estuários tropicais de Pernambuco (Brasil), sujeitos a diferentes condições hidrológicas. <i>Revista Brasileira de Ciencias Agrarias</i> , 2015, 10, 304-314.	0.2	3
34	Water quality, phytoplankton composition and growth of <i>Litopenaeus vannamei</i> (Boone) in an integrated biofloc system with <i>Gracilaria birdiae</i> (Greville) and <i>Gracilaria domingensis</i> (Katz). <i>Aquaculture International</i> , 2014, 22, 1649-1664.	2.2	29
35	Influência do nível hidrológico sobre a dieta de <i>Leporinus reinhardtii</i> (Characiformes, Anostomidae) em um reservatório do semiárido brasileiro. <i>Iheringia - Serie Zoologia</i> , 2014, 104, 290-298.	0.5	6
36	Samambaias aquáticas da bacia do rio de Contas, Bahia, Brasil. <i>Neotropical Biology and Conservation</i> , 2014, 9, .	0.9	1

#	ARTICLE	IF	CITATIONS
37	Spatial-temporal variation of <i>Achirus</i> larvae (Actinopterygii: Achiridae) in mangrove, beach and reef habitats in north-eastern Brazil. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2013, 93, 381-388.	0.8	11
38	Genetic Diversity of Captive and Wild Threatened Catfish <i>Pseudoplatystoma corruscans</i> in the São Francisco River. <i>Reviews in Fisheries Science</i> , 2013, 21, 237-246.	2.1	14
39	The influence of seasonality on fish life stages and residence in surf zones: a case of study in a tropical region. <i>Biota Neotropica</i> , 2013, 13, 181-192.	1.0	9
40	Caatinga Revisited: Ecology and Conservation of an Important Seasonal Dry Forest. <i>Scientific World Journal</i> , The, 2012, 2012, 1-18.	2.1	170
41	Composição da ictiofauna em ambientes marginais e tributários do mādio-submādio rio São Francisco. <i>Revista Brasileira de Ciencias Agrarias</i> , 2012, 7, 358-366.	0.2	4
42	New species of <i>Radiospongilla</i> (Porifera: Spongillidae) from Brazilian inland waters. <i>Zootaxa</i> , 2011, 3132, 56.	0.5	11
43	Buccal apparatus and gastrointestinal tract dimensions associated to the diet of early life stages of <i>Centropomus undecimalis</i> (Centropomidae, Actinopterygii). <i>Iheringia - Serie Zoologia</i> , 2011, 101, 85-92.	0.5	5
44	O gradiente rio-barragem do reservatório de Sobradinho afeta a composição florística, riqueza e formas biológicas das macrofitas aquáticas?. <i>Rodriguesia</i> , 2011, 62, 731-742.	0.9	13
45	Diet composition and food overlap of <i>Acestorhynchus britskii</i> and <i>A. lacustris</i> (Characiformes: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 10 Biological Sciences, 2011, 33, .	0.3	3
46	Ictiofauna do reservatório de Duas Unas, bacia do rio Jaboatão, Pernambuco: resultados preliminares da composição e estrutura da assembleia. <i>Revista Brasileira de Ciencias Agrarias</i> , 2011, 6, 351-361.	0.2	4
47	Morphological development of <i>Anchoviella vaillanti</i> (Steindachner, 1908) (Clupeiformes: Engraulidae) larvae and early juveniles. <i>Neotropical Ichthyology</i> , 2010, 8, 805-812.	1.0	5
48	Estrutura da assembleia de peixes de uma lagoa marginal desconectada do rio, no submādio Rio São Francisco, Pernambuco. <i>Biota Neotropica</i> , 2009, 9, 117-129.	1.0	12
49	Desenvolvimento ovariano de <i>Plagioscion squamosissimus</i> (Heckel, 1840) (Actinopterygii, Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 10 Tf 5)	1.0	5
50	Description of <i>Atherinella brasiliensis</i> (Quoy & Gaimard, 1825) (Atheriniformes: Atherinopsidae) larvae from the Jaguaribe River estuary, Itamaracá island, Northeastern Brazil. <i>Neotropical Ichthyology</i> , 2007, 5, 369-374.	1.0	4
51	Dinâmica espacial e temporal de zoeas de <i>Brachyura</i> (Crustacea, Decapoda) no estuário do Rio Jaguaribe, Itamaracá, Pernambuco, Brasil. <i>Iheringia - Serie Zoologia</i> , 2007, 97, 434-440.	0.5	7
52	Morphological development of <i>Pellona flavipinnis</i> post-yolk-sac larvae and juveniles (Clupeiformes: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50)	0.5	5
53	Structural and morphological features of <i>Piaractus mesopotamicus</i> (Holmberg, 1887) gills. <i>Revista Brasileira De Biologia</i> , 2000, 60, 493-501.	0.3	10
54	Ventilatory flow relative to intrabuccal and intraopercular volumes in the serrasalmid fish <i>Piaractus mesopotamicus</i> during normoxia and exposed to graded hypoxia. <i>Revista Brasileira De Biologia</i> , 2000, 60, 249-254.	0.3	8

#	ARTICLE	IF	CITATIONS
55	Desenvolvimento larval inicial de <i>Helostoma temminckii</i> Cuvier & Valenciennes (Helostomatidae), Tj ETQq1 1 0,784314 gBT /Over	0,5	
56	Functional morphology of gills and respiratory area of two active rheophilic fish species, <i>Plagioscion squamosissimus</i> and <i>Prochilodus scrofa</i> . Journal of Fish Biology, 1998, 52, 50-61.	1.6	15
57	The occurrence of aerial respiration in <i>Rhinelepis strigosa</i> during progressive hypoxia. Journal of Fish Biology, 1998, 52, 369-379.	1.6	17
58	Respiratory gill surface of the serrasalmid fish, <i>Piaractus mesopotamicus</i> . Journal of Fish Biology, 1997, 50, 127-136.	1.6	11
59	The gill filament muscles in two loricariid fish (genus <i>Hypostomus</i> and <i>Rhinelepis</i> ). Journal of Fish Biology, 1995, 46, 1082-1085.	1.6	10
60	Respiratory gill surface area of a facultative air-breathing loricariid fish, <i>Rhinelepis strigosa</i> . Canadian Journal of Zoology, 1994, 72, 2009-2015.	1.0	12
61	REPRODUCTION ASPECTS OF COBIA CAUGHT IN PERNAMBUCO COAST, NORTHEASTERN BRAZIL. Boletim Do Instituto De Pesca, 0, 47, .	0,5	1
62	Ecomorphological relations of sympatric juveniles of Clupeiformes from a Brazilian sandy beach. Iheringia - Serie Zoologia, 0, 112, .	0,5	1