
Silvina Stella Maris Villanueva

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1861354/publications.pdf Version: 2024-02-01



SILVINA STELLA MARIS

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Multidrug resistance-associated protein 2 is negatively regulated by oxidative stress in rat intestine via a posttranslational mechanism. Impact on its membrane barrier function. Toxicology, 2021, 460, 152873. | 4.2 | 0 |
| 2 | Role of interleukin 1 beta in the regulation of rat intestinal multidrug resistance-associated protein 2 under conditions of experimental endotoxemia. Toxicology, 2020, 441, 152527. | 4.2 | 3 |
| 3 | Reversion of down-regulation of intestinal multidrug resistance-associated protein 2 in fructose-fed rats by geraniol and vitamin C: Potential role of inflammatory response and oxidative stress. Journal of Nutritional Biochemistry, 2019, 68, 7-15. | 4.2 | 10 |
| 4 | Biphasic modulation of cAMP levels by the contraceptive nomegestrol acetate. Impact on P-glycoprotein expression and activity in hepatic cells. Biochemical Pharmacology, 2018, 154, 118-126. | 4.4 | 3 |
| 5 | Acute regulation of multidrug resistance-associated protein 2 localization and activity by cAMP and estradiol-17β-d-glucuronide in rat intestine and Caco-2 cells. Archives of Toxicology, 2018, 92, 777-788. | 4.2 | 8 |
| 6 | Inhibition of multidrug resistance-associated protein 2 (MRP2) activity by the contraceptive nomegestrol acetate in HepG2 and Caco-2 cells. European Journal of Pharmaceutical Sciences, 2018, 122, 205-213. | 4.0 | 6 |
| 7 | Intestinal multidrug resistance-associated protein 2 is down-regulated in fructose-fed rats. Journal of Nutritional Biochemistry, 2017, 40, 178-186. | 4.2 | 13 |
| 8 | Glucagon-like peptide 2 prevents down-regulation of intestinal multidrug resistance-associated protein 2 and P-glycoprotein in endotoxemic rats. Toxicology, 2017, 390, 22-31. | 4.2 | 9 |
| 9 | The phytoestrogen genistein enhances multidrug resistance in breast cancer cell lines by translational regulation of ABC transporters. Cancer Letters, 2016, 376, 165-172. | 7.2 | 50 |
| 10 | Modulation of expression and activity of intestinal multidrug resistance-associated protein 2 by xenobiotics. Toxicology and Applied Pharmacology, 2016, 303, 45-57. | 2.8 | 22 |
| 11 | Regulation of Multidrug Resistance Proteins by Genistein in a Hepatocarcinoma Cell Line: Impact on Sorafenib Cytotoxicity. PLoS ONE, 2015, 10, e0119502. | 2.5 | 56 |
| 12 | Coordinated induction of GST and MRP2 by cAMP in Caco-2 cells: Role of protein kinase A signaling pathway and toxicological relevance. Toxicology and Applied Pharmacology, 2015, 287, 178-190. | 2.8 | 16 |
| 13 | Estrogen receptor-α mediates human multidrug resistance associated protein 3 induction by 17α-ethynylestradiol. Role of activator protein-1. Biochemical Pharmacology, 2013, 86, 401-409. | 4.4 | 17 |
| 14 | Induction of Hepatic Multidrug Resistance-Associated Protein 3 by Ethynylestradiol Is Independent of Cholestasis and Mediated by Estrogen Receptor. Drug Metabolism and Disposition, 2013, 41, 275-280. | 3.3 | 20 |
| 15 | Acetaminophen-induced stimulation of MDR1 expression and activity in rat intestine and in LS 174T human intestinal cell line. Biochemical Pharmacology, 2011, 81, 244-250. | 4.4 | 20 |
| 16 | Pregnane X receptor mediates the induction of P-glycoprotein by spironolactone in HepG2 cells. Toxicology, 2011, 285, 18-24. | 4.2 | 46 |