

# Mi Jung Lee

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/1862954/mi-jung-lee-publications-by-citations.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

72  
papers

1,483  
citations

22  
h-index

35  
g-index

75  
ext. papers

1,752  
ext. citations

3.9  
avg. IF

3.73  
L-index

#	Paper	IF	Citations
72	Circulating Klotho levels in CKD and relationship to progression. <i>American Journal of Kidney Diseases</i> , <b>2013</b> , 61, 899-909	7.4	116
71	Thyroid hormone replacement therapy attenuates the decline of renal function in chronic kidney disease patients with subclinical hypothyroidism. <i>Thyroid</i> , <b>2013</b> , 23, 654-61	6.2	79
70	Preservation of renal function by thyroid hormone replacement therapy in chronic kidney disease patients with subclinical hypothyroidism. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2012</b> , 97, 2732-40	5.6	79
69	An increase in mean platelet volume from baseline is associated with mortality in patients with severe sepsis or septic shock. <i>PLoS ONE</i> , <b>2015</b> , 10, e0119437	3.7	78
68	Decreased circulating C3 levels and mesangial C3 deposition predict renal outcome in patients with IgA nephropathy. <i>PLoS ONE</i> , <b>2012</b> , 7, e40495	3.7	73
67	High-Dose Versus Conventional-Dose Continuous Venovenous Hemodiafiltration and Patient and Kidney Survival and Cytokine Removal in Sepsis-Associated Acute Kidney Injury: A Randomized Controlled Trial. <i>American Journal of Kidney Diseases</i> , <b>2016</b> , 68, 599-608	7.4	64
66	Irisin, a novel myokine is an independent predictor for sarcopenia and carotid atherosclerosis in dialysis patients. <i>Atherosclerosis</i> , <b>2015</b> , 242, 476-82	3.1	55
65	Plasma levels of soluble receptor for advanced glycation end products (sRAGE) and proinflammatory ligand for RAGE (EN-RAGE) are associated with carotid atherosclerosis in patients with peritoneal dialysis. <i>Atherosclerosis</i> , <b>2012</b> , 220, 208-14	3.1	50
64	Decreased circulating klotho levels in patients undergoing dialysis and relationship to oxidative stress and inflammation. <i>Peritoneal Dialysis International</i> , <b>2015</b> , 35, 43-51	2.8	40
63	Interdialytic weight gain and cardiovascular outcome in incident hemodialysis patients. <i>American Journal of Nephrology</i> , <b>2014</b> , 39, 427-35	4.6	40
62	Clinical outcomes, when matched at presentation, do not vary between adult-onset Henrich-Schleim purpura nephritis and IgA nephropathy. <i>Kidney International</i> , <b>2012</b> , 82, 1304-12	9.9	36
61	Prognostic Value of Residual Urine Volume, GFR by 24-hour Urine Collection, and eGFR in Patients Receiving Dialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , <b>2017</b> , 12, 426-434	6.9	30
60	Diastolic dysfunction is an independent predictor of cardiovascular events in incident dialysis patients with preserved systolic function. <i>PLoS ONE</i> , <b>2015</b> , 10, e0118694	3.7	29
59	Vitamin D and Hypertension. <i>Electrolyte and Blood Pressure</i> , <b>2017</b> , 15, 1-11	1.1	28
58	Early initiation of continuous renal replacement therapy improves patient survival in severe progressive septic acute kidney injury. <i>Journal of Critical Care</i> , <b>2012</b> , 27, 743.e9-18	4	27
57	Comparison of the Haas and the Oxford classifications for prediction of renal outcome in patients with IgA nephropathy. <i>Human Pathology</i> , <b>2014</b> , 45, 236-43	3.7	26
56	Progression of aortic arch calcification over 1 year is an independent predictor of mortality in incident peritoneal dialysis patients. <i>PLoS ONE</i> , <b>2012</b> , 7, e48793	3.7	26

55	Glomerular IgG deposition predicts renal outcome in patients with IgA nephropathy. <i>Modern Pathology</i> , <b>2016</b> , 29, 743-52	9.8	25
54	Good glycemic control is associated with better survival in diabetic patients on peritoneal dialysis: a prospective observational study. <i>PLoS ONE</i> , <b>2012</b> , 7, e30072	3.7	24
53	Change of Nutritional Status Assessed Using Subjective Global Assessment Is Associated With All-Cause Mortality in Incident Dialysis Patients. <i>Medicine (United States)</i> , <b>2016</b> , 95, e2714	1.8	24
52	Clinical implication of crescentic lesions in immunoglobulin A nephropathy. <i>Nephrology Dialysis Transplantation</i> , <b>2014</b> , 29, 356-64	4.3	23
51	Visceral fat thickness is associated with carotid atherosclerosis in peritoneal dialysis patients. <i>Obesity</i> , <b>2012</b> , 20, 1301-7	8	22
50	Leptin/adiponectin ratio is an independent predictor of mortality in nondiabetic peritoneal dialysis patients. <i>Peritoneal Dialysis International</i> , <b>2013</b> , 33, 67-74	2.8	22
49	Diastolic dysfunction is associated with an increased risk of contrast-induced nephropathy: a retrospective cohort study. <i>BMC Nephrology</i> , <b>2013</b> , 14, 146	2.7	20
48	Urine output is associated with prognosis in patients with acute kidney injury requiring continuous renal replacement therapy. <i>Journal of Critical Care</i> , <b>2013</b> , 28, 379-88	4	20
47	Optimal proteinuria target for renoprotection in patients with IgA nephropathy. <i>PLoS ONE</i> , <b>2014</b> , 9, e101375	3.7	19
46	Podocyte hypertrophy precedes apoptosis under experimental diabetic conditions. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , <b>2015</b> , 20, 1056-71	5.4	18
45	Endothelial dysfunction is associated with major adverse cardiovascular events in peritoneal dialysis patients. <i>Medicine (United States)</i> , <b>2014</b> , 93, e73	1.8	18
44	Low serum intact parathyroid hormone level is an independent risk factor for overall mortality and major adverse cardiac and cerebrovascular events in incident dialysis patients. <i>Osteoporosis International</i> , <b>2016</b> , 27, 2717-2726	5.3	18
43	The benefit of specialized team approaches in patients with acute kidney injury undergoing continuous renal replacement therapy: propensity score matched analysis. <i>Critical Care</i> , <b>2014</b> , 18, 454	10.8	17
42	Elevated osteoprotegerin is associated with inflammation, malnutrition and new onset cardiovascular events in peritoneal dialysis patients. <i>Atherosclerosis</i> , <b>2011</b> , 219, 925-30	3.1	17
41	The atherogenic index of plasma and the risk of mortality in incident dialysis patients: Results from a nationwide prospective cohort in Korea. <i>PLoS ONE</i> , <b>2017</b> , 12, e0177499	3.7	17
40	Glycemic Control Modifies Difference in Mortality Risk Between Hemodialysis and Peritoneal Dialysis in Incident Dialysis Patients With Diabetes: Results From a Nationwide Prospective Cohort in Korea. <i>Medicine (United States)</i> , <b>2016</b> , 95, e3118	1.8	17
39	Electrocardiographic left ventricular hypertrophy and outcome in hemodialysis patients. <i>PLoS ONE</i> , <b>2012</b> , 7, e35534	3.7	16
38	Mean platelet volume is a prognostic factor in patients with acute kidney injury requiring continuous renal replacement therapy. <i>Journal of Critical Care</i> , <b>2014</b> , 29, 1016-21	4	15

37	The impact of low triiodothyronine levels on mortality is mediated by malnutrition and cardiac dysfunction in incident hemodialysis patients. <i>European Journal of Endocrinology</i> , <b>2013</b> , 169, 409-19	6.5	14
36	The effect of specialized continuous renal replacement therapy team in acute kidney injury patients treatment. <i>Yonsei Medical Journal</i> , <b>2015</b> , 56, 658-65	3	14
35	Which Biomarker is the Best for Predicting Mortality in Incident Peritoneal Dialysis Patients: NT-ProBNP, Cardiac TnT, or hsCRP?: A Prospective Observational Study. <i>Medicine (United States)</i> , <b>2015</b> , 94, e1636	1.8	13
34	Normal body mass index with central obesity has increased risk of coronary artery calcification in Korean patients with chronic kidney disease. <i>Kidney International</i> , <b>2016</b> , 90, 1368-1376	9.9	12
33	Low circulating adiponectin levels are associated with insulin resistance in non-obese peritoneal dialysis patients. <i>Endocrine Journal</i> , <b>2012</b> , 59, 685-95	2.9	12
32	The relationship of initial transferrin saturation to cardiovascular parameters and outcomes in patients initiating dialysis. <i>PLoS ONE</i> , <b>2014</b> , 9, e87231	3.7	11
31	Removal of kidney stones by extracorporeal shock wave lithotripsy is associated with delayed progression of chronic kidney disease. <i>Yonsei Medical Journal</i> , <b>2012</b> , 53, 708-14	3	10
30	The effect of renin-angiotensin system blockade on renal protection in chronic kidney disease patients with hyperkalemia. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , <b>2014</b> , 15, 491-7		9
29	Left atrial enlargement is associated with a rapid decline in residual renal function in ESRD patients on peritoneal dialysis. <i>Journal of the American Society of Echocardiography</i> , <b>2012</b> , 25, 421-7	5.8	9
28	Changes in geriatric nutritional risk index and risk of major adverse cardiac and cerebrovascular events in incident peritoneal dialysis patients. <i>Kidney Research and Clinical Practice</i> , <b>2017</b> , 36, 377-386	3.6	8
27	Serum Ferritin Predicts Mortality Regardless of Inflammatory and Nutritional Status in Patients Starting Dialysis: A Prospective Cohort Study. <i>Blood Purification</i> , <b>2015</b> , 40, 209-17	3.1	8
26	The effect of HMG-CoA reductase inhibitor on insulin resistance in patients undergoing peritoneal dialysis. <i>Cardiovascular Drugs and Therapy</i> , <b>2012</b> , 26, 501-9	3.9	8
25	Insulin resistance is associated with new-onset cardiovascular events in nondiabetic patients undergoing peritoneal dialysis. <i>Kidney Research and Clinical Practice</i> , <b>2014</b> , 33, 192-8	3.6	7
24	Stepwise treatment using corticosteroids alone and in combination with cyclosporine in Korean patients with idiopathic membranous nephropathy. <i>Yonsei Medical Journal</i> , <b>2013</b> , 54, 973-82	3	7
23	Sagittal abdominal diameter is an independent predictor of all-cause and cardiovascular mortality in incident peritoneal dialysis patients. <i>PLoS ONE</i> , <b>2013</b> , 8, e77082	3.7	7
22	Albuminuria as a Risk Factor for Anemia in Chronic Kidney Disease: Result from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>PLoS ONE</i> , <b>2015</b> , 10, e0139747	3.7	7
21	Association between Vitamin D Deficiency and Anemia in Patients with End-Stage Renal Disease: A Cross-Sectional Study. <i>Yonsei Medical Journal</i> , <b>2016</b> , 57, 1159-64	3	7
20	Delta neutrophil index is an independent predictor of mortality in septic acute kidney injury patients treated with continuous renal replacement therapy. <i>BMC Nephrology</i> , <b>2017</b> , 18, 94	2.7	6

19	Changes in echocardiographic parameters according to the rate of residual renal function decline in incident peritoneal dialysis patients. <i>Medicine (United States)</i> , <b>2015</b> , 94, e427	1.8	6
18	Decrease in waist-to-hip ratio reduced the development of chronic kidney disease in non-obese non-alcoholic fatty liver disease. <i>Scientific Reports</i> , <b>2020</b> , 10, 8996	4.9	6
17	NT-proBNP: is it a more significant risk factor for mortality than troponin T in incident hemodialysis patients?. <i>Medicine (United States)</i> , <b>2014</b> , 93, e241	1.8	6
16	Association between the ratio of insulin-like growth factor-I to insulin-like growth factor binding protein-3 and inflammation in incident automated peritoneal dialysis patients. <i>Growth Hormone and IGF Research</i> , <b>2013</b> , 23, 170-4	2	6
15	Increased dialysate MCP-1 is associated with cardiovascular mortality in peritoneal dialysis patients: a prospective observational study. <i>American Journal of Nephrology</i> , <b>2014</b> , 40, 291-9	4.6	6
14	Low Mitochondrial DNA Copy Number is Associated With Adverse Clinical Outcomes in Peritoneal Dialysis Patients. <i>Medicine (United States)</i> , <b>2016</b> , 95, e2717	1.8	6
13	Urine Osmolality and Renal Outcome in Patients with Chronic Kidney Disease: Results from the KNOW-CKD. <i>Kidney and Blood Pressure Research</i> , <b>2019</b> , 44, 1089-1100	3.1	5
12	Membranous glomerulonephritis in a patient with myelodysplastic syndrome-refractory cytopenia with multilineage dysplasia. <i>Kidney Research and Clinical Practice</i> , <b>2013</b> , 32, 134-7	3.6	5
11	LDL cholesterol affects clinical outcomes in incident hemodialysis patients during the early stages of dialysis. <i>Blood Purification</i> , <b>2014</b> , 38, 131-9	3.1	5
10	A novel multiplex real-time PCR assay for the concurrent detection of hepatitis A, B and C viruses in patients with acute hepatitis. <i>PLoS ONE</i> , <b>2012</b> , 7, e49106	3.7	5
9	Association of serum uric acid level with coronary artery stenosis severity in Korean end-stage renal disease patients. <i>Kidney Research and Clinical Practice</i> , <b>2017</b> , 36, 282-289	3.6	5
8	Predictive value of serum albumin-to-globulin ratio for incident chronic kidney disease: A 12-year community-based prospective study. <i>PLoS ONE</i> , <b>2020</b> , 15, e0238421	3.7	5
7	Is It Beneficial to Convert to a Neutral-pH Bicarbonate/Lactate-Buffered PD Solution in Long-Term CAPD Patients? A Single-Center Prospective Study. <i>Peritoneal Dialysis International</i> , <b>2015</b> , 35, 366-9	2.8	4
6	Effect of peritoneal dialysis modality on the 1-year rate of decline of residual renal function. <i>Yonsei Medical Journal</i> , <b>2014</b> , 55, 141-8	3	4
5	A case of Tsutsugamushi disease presenting with nephrotic syndrome. <i>Korean Journal of Internal Medicine</i> , <b>2013</b> , 28, 728-31	2.5	4
4	Syndrome of inappropriate antidiuretic hormone secretion in a patient with large cell neuroendocrine carcinoma. <i>Yonsei Medical Journal</i> , <b>2012</b> , 53, 667-9	3	3
3	Prognostic value of soluble ST2 and soluble LR11 on mortality and cardiovascular events in peritoneal dialysis patients. <i>BMC Nephrology</i> , <b>2020</b> , 21, 228	2.7	2
2	Comparison of dietary intake patterns in hemodialysis patients by nutritional status: A cross-sectional analysis. <i>Kidney Research and Clinical Practice</i> , <b>2020</b> , 39, 202-212	3.6	2

1

Smoking Cessation and Coronary Artery Calcification in CKD. *Clinical Journal of the American Society of Nephrology: CJASN*, **2021**, 16, 870-879

6.9 ○