## Annick Ortalo-Magné

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1868395/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Impact of phosphate concentration on the metabolome of biofilms of the marine bacterium Pseudoalteromonas lipolytica. Metabolomics, 2022, 18, 18.	3.0	2
2	Disrupting quorum sensing alters social interactions in Chromobacterium violaceum. Npj Biofilms and Microbiomes, 2021, 7, 40.	6.4	30
3	Metabolomic and proteomic changes induced by growth inhibitory concentrations of copper in the biofilm-forming marine bacterium <i>Pseudoalteromonas lipolytica</i> . Metallomics, 2019, 11, 1887-1899.	2.4	12
4	Sterols from the brown alga Cystoseira foeniculacea: Degradation of fucosterol into saringosterol epimers. Arabian Journal of Chemistry, 2019, 12, 1474-1478.	4.9	18
5	Metabolome and proteome changes between biofilm and planktonic phenotypes of the marine bacterium <i>Pseudoalteromonas lipolytica</i> TC8. Biofouling, 2018, 34, 132-148.	2.2	31
6	Discrimination of Four Marine Biofilm-Forming Bacteria by LC–MS Metabolomics and Influence of Culture Parameters. Journal of Proteome Research, 2017, 16, 1962-1975.	3.7	43
7	Characterization and anti-biofilm activity of extracellular polymeric substances produced by the marine biofilm-forming bacterium <i>Pseudoalteromonas ulvae</i> strain TC14. Biofouling, 2016, 32, 547-560.	2.2	39
8	Extraction, Purification, and NMR Analysis of Terpenes from Brown Algae. Methods in Molecular Biology, 2015, 1308, 207-223.	0.9	13
9	Cystophloroketals A–E, Unusual Phloroglucinol–Meroterpenoid Hybrids from the Brown Alga <i>Cystoseira tamariscifolia</i> . Journal of Natural Products, 2015, 78, 1663-1670.	3.0	27
10	Modulation of violacein production and phenotypes associated with biofilm by exogenous quorum sensing N-acylhomoserine lactones in the marine bacterium Pseudoalteromonas ulvae TC14. Microbiology (United Kingdom), 2015, 161, 2039-2051.	1.8	38
11	Identification of Bacterial Strains Isolated from the Mediterranean Sea Exhibiting Different Abilities of Biofilm Formation. Microbial Ecology, 2014, 68, 94-110.	2.8	46
12	Correlation between synthesis variation of 2-alkylquinolones and the antifungal activity of a Burkholderia cepacia strain collection. World Journal of Microbiology and Biotechnology, 2012, 28, 275-281.	3.6	16
13	Environmental Burkholderia cepacia Strain Cs5 Acting by Two Analogous Alkyl-Quinolones and a Didecyl-Phthalate Against a Broad Spectrum of Phytopathogens Fungi. Current Microbiology, 2011, 62, 1490-1495.	2.2	39
14	Dictyotadimer A, a new dissymmetric bis-diterpene from a brown alga of the genus Dictyota. Tetrahedron Letters, 2011, 52, 1031-1035.	1.4	14
15	Antifungal activities of an endophytic Pseudomonas fluorescens strain Pf1TZ harbouring genes from pyoluteorin and phenazine clusters. Biotechnology Letters, 2010, 32, 1279-1285.	2.2	15
16	Eicosapentaenoic acid: Possible precursor of the phloroglucinol derivatives isolated from the brown alga Zonaria tournefortii (J.V. Lamouroux) Montagne. Biochemical Systematics and Ecology, 2009, 37, 55-58.	1.3	16
17	Diterpenoids from the Mediterranean Brown Alga <i>Dictyota</i> sp. Evaluated as Antifouling Substances against a Marine Bacterial Biofilm. Journal of Natural Products, 2009, 72, 1299-1304.	3.0	63
18	Trihydroxylated linear diterpenes from the brown alga Bifurcaria bifurcata (Fucales, Phaeophyta). Biochemical Systematics and Ecology, 2008, 36, 484-489.	1.3	15

#	Article	IF	CITATIONS
19	Antifouling Activity of Meroditerpenoids from the Marine Brown Alga <i>Halidrys siliquosa</i> . Journal of Natural Products, 2008, 71, 1121-1126.	3.0	57
20	Meroditerpenoids and Derivatives from the Brown Alga <i>Cystoseira baccata</i> and Their Antifouling Properties. Journal of Natural Products, 2008, 71, 1806-1811.	3.0	60
21	An extract from the brown alga Bifurcaria bifurcata induces irreversible arrest of cell proliferation in a non-small-cell bronchopulmonary carcinoma line. Journal of Applied Phycology, 2006, 18, 87-93.	2.8	27
22	Polar acyclic diterpenoids from Bifurcaria bifurcata (Fucales, Phaeophyta). Phytochemistry, 2005, 66, 2316-2323.	2.9	42
23	Seasonal variation in antifouling activity of crude extracts of the brown alga Bifurcaria bifurcata (Cystoseiraceae) against cyprids of Balanus amphitrite and the marine bacteria Cobetia marina and Pseudoalteromonas haloplanktis. Journal of Experimental Marine Biology and Ecology, 2004, 313, 47-62.	1.5	113
24	Trihydroxylated linear diterpenes from the brown alga Bifurcaria bifurcata. Phytochemistry, 2004, 65, 2063-2069.	2.9	46
25	Isolation of the Volatile Compounds from the Brown AlgaDictyopteris membranaceaby Focused Microwave-Assisted Hydrodistillation. Journal of Essential Oil Research, 2002, 14, 422-424.	2.7	8
26	Seasonal variations in the chemical composition of Bifurcaria bifurcata (Cystoseiraceae). Biochemical Systematics and Ecology, 2002, 30, 61-64.	1.3	16
27	(S)-12-Hydroxygeranylgeraniol-derived diterpenes from the brown alga Bifurcaria bifurcata. Phytochemistry, 2001, 57, 529-535.	2.9	35
28	Acyclic diterpenes and sterols from the genera Bifurcaria and Bifurcariopsis (Cystoseiraceae,) Tj ETQq0 0 0 rgBT /	Overlock 1	0 Tf 50 382

29	Cytometric detection of mycobacterial surface antigens: exposure of mannosyl epitopes and of the arabinan segment of arabinomannans. Journal of Bacteriology, 1996, 178, 7254-7259.	2.2	21
30	Extracellular and surface-exposed polysaccharides of non-tuberculous mycobacteria. Microbiology (United Kingdom), 1996, 142, 1513-1520.	1.8	88
31	The outermost capsular arabinomannans and other mannoconjugates of virulent and avirulent tubercle bacilli. Microbiology (United Kingdom), 1996, 142, 927-935.	1.8	58
32	Identification of the surface-exposed lipids on the cell envelopes of Mycobacterium tuberculosis and other mycobacterial species, Journal of Bacteriology, 1996, 178, 456-461	2.2	235