

# Paul J Fletcher

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1871693/publications.pdf>

Version: 2024-02-01

187  
papers

9,641  
citations

27035

58  
h-index

56606

87  
g-index

188  
all docs

188  
docs citations

188  
times ranked

8628  
citing authors

#	ARTICLE	IF	CITATIONS
1	Effects of pimavanserin and lorcaserin on alcohol self-administration and reinstatement in male and female rats. <i>Neuropharmacology</i> , 2022, , 109150.	2.0	3
2	Prenatal disruption of D1R-SynGAP complex causes cognitive deficits in adulthood. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 105, 110122.	2.5	3
3	The serotonin 2C receptor agonist lorcaserin, alone and in combination with the opioid receptor antagonist naltrexone, attenuates binge-like ethanol drinking. <i>Addiction Biology</i> , 2021, 26, e13040.	1.4	9
4	Lorcaserin: A review of its preclinical and clinical pharmacology and therapeutic potential. , 2020, 205, 107417.		52
5	Clozapine reliably increases the motivation for food: parsing the role of the 5-HT <sub>2c</sub> and H1 receptors. <i>Psychopharmacology</i> , 2020, 237, 957-966.	1.5	9
6	The pharmacological stressor yohimbine, but not U50,488, increases responding for conditioned reinforcers paired with ethanol or sucrose. <i>Psychopharmacology</i> , 2020, 237, 3689-3702.	1.5	7
7	Serotonin transporter protein in autopsied brain of chronic users of cocaine. <i>Psychopharmacology</i> , 2020, 237, 2661-2671.	1.5	7
8	Effects of exposure to chronic uncertainty and a sensitizing regimen of amphetamine injections on locomotion, decision-making, and dopamine receptors in rats. <i>Neuropsychopharmacology</i> , 2020, 45, 811-822.	2.8	13
9	Median raphe serotonin neurons promote anxiety-like behavior via inputs to the dorsal hippocampus. <i>Neuropharmacology</i> , 2020, 168, 107985.	2.0	42
10	Disruption of SynGAP-dopamine D1 receptor complexes alters actin and microtubule dynamics and impairs GABAergic interneuron migration. <i>Science Signaling</i> , 2019, 12, .	1.6	11
11	Nicotine enhances responding for conditioned reinforcement via $\alpha 4\beta 2$ nicotinic acetylcholine receptors in the ventral tegmental area, but not the nucleus accumbens or the prefrontal cortex. <i>Neuropharmacology</i> , 2019, 148, 68-76.	2.0	3
12	Kappa opioid receptors mediate yohimbine-induced increases in impulsivity in the 5-choice serial reaction time task. <i>Behavioural Brain Research</i> , 2019, 359, 258-265.	1.2	11
13	Dorsal raphe serotonin neurons inhibit operant responding for reward via inputs to the ventral tegmental area but not the nucleus accumbens: evidence from studies combining optogenetic stimulation and serotonin reuptake inhibition. <i>Neuropsychopharmacology</i> , 2019, 44, 793-804.	2.8	39
14	Preclinical evidence for combining the 5-HT <sub>2C</sub> receptor agonist lorcaserin and varenicline as a treatment for nicotine dependence. <i>Addiction Biology</i> , 2019, 24, 376-387.	1.4	9
15	Clozapine reduces nicotine self-administration, blunts reinstatement of nicotine-seeking but increases responding for food. <i>Addiction Biology</i> , 2019, 24, 565-576.	1.4	10
16	Adolescent exposure to $\Delta^9$ -tetrahydrocannabinol delays acquisition of paired-associates learning in adulthood. <i>Psychopharmacology</i> , 2019, 236, 1875-1886.	1.5	23
17	Deep brain stimulation and fluoxetine exert different long-term changes in the serotonergic system. <i>Neuropharmacology</i> , 2018, 135, 63-72.	2.0	22
18	Acquisition of nicotine self-administration in amphetamine and phencyclidine models of schizophrenia: A role for stress?. <i>Schizophrenia Research</i> , 2018, 194, 98-106.	1.1	6

#	ARTICLE	IF	CITATIONS
19	Objective assessment of exploratory behaviour in schizophrenia using wireless motion capture. <i>Schizophrenia Research</i> , 2018, 195, 122-129.	1.1	3
20	Deep brain stimulation induces antidepressant-like effects in serotonin transporter knockout mice. <i>Brain Stimulation</i> , 2018, 11, 423-425.	0.7	17
21	Objective investigation of activity preference in schizophrenia: A pilot study. <i>Psychiatry Research</i> , 2018, 267, 551-559.	1.7	1
22	Effects of 5-HT <sub>1A</sub> , 5-HT <sub>2A</sub> and 5-HT <sub>2C</sub> receptor agonists and antagonists on responding for a conditioned reinforcer and its enhancement by methylphenidate. <i>Psychopharmacology</i> , 2017, 234, 889-902.	1.5	8
23	Pdxdc1 modulates prepulse inhibition of acoustic startle in the mouse. <i>Translational Psychiatry</i> , 2017, 7, e1125-e1125.	2.4	12
24	Studies To Examine Potential Tolerability Differences between the 5-HT <sub>2C</sub> Receptor Selective Agonists Lorcaserin and CP-809101. <i>ACS Chemical Neuroscience</i> , 2017, 8, 1074-1084.	1.7	8
25	Role of impulsivity and reward in the anti-obesity actions of 5-HT <sub>2C</sub> receptor agonists. <i>Journal of Psychopharmacology</i> , 2017, 31, 1403-1418.	2.0	30
26	Pharmacological Modulation of 5-HT <sub>2C</sub> Receptor Activity Produces Bidirectional Changes in Locomotor Activity, Responding for a Conditioned Reinforcer, and Mesolimbic DA Release in C57BL/6 Mice. <i>Neuropsychopharmacology</i> , 2017, 42, 2178-2187.	2.8	24
27	Lipoic acid and haloperidol-induced vacuous chewing movements: Implications for prophylactic antioxidant use in tardive dyskinesia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2017, 72, 23-29.	2.5	10
28	Activation of Dopamine D1-D2 Receptor Complex Attenuates Cocaine Reward and Reinstatement of Cocaine-Seeking through Inhibition of DARPP-32, ERK, and $\hat{I}^{\beta}$ FosB. <i>Frontiers in Pharmacology</i> , 2017, 8, 924.	1.6	55
29	Uncertainty exposure causes behavioural sensitization and increases risky decision-making in male rats: toward modelling gambling disorder. <i>Journal of Psychiatry and Neuroscience</i> , 2017, 42, 404-413.	1.4	72
30	Decreased Incentive Motivation Following Knockout or Acute Blockade of the Serotonin Transporter: Role of the 5-HT <sub>2C</sub> Receptor. <i>Neuropsychopharmacology</i> , 2016, 41, 2566-2576.	2.8	22
31	Disruption of a dopamine receptor complex amplifies the actions of cocaine. <i>European Neuropsychopharmacology</i> , 2016, 26, 1366-1377.	0.3	36
32	Mgat5 modulates the effect of early life stress on adult behavior and physical health in mice. <i>Behavioural Brain Research</i> , 2016, 312, 253-264.	1.2	14
33	Atypical antipsychotics and effects on feeding: from mice to men. <i>Psychopharmacology</i> , 2016, 233, 2629-2653.	1.5	38
34	Responding for a conditioned reinforcer or unconditioned sensory reinforcer in mice: interactions with environmental enrichment, social isolation, and monoamine reuptake inhibitors. <i>Psychopharmacology</i> , 2016, 233, 983-993.	1.5	6
35	Lorcaserin and CP-809101 reduce motor impulsivity and reinstatement of food seeking behavior in male rats: Implications for understanding the anti-obesity property of 5-HT <sub>2C</sub> receptor agonists. <i>Psychopharmacology</i> , 2016, 233, 2841-2856.	1.5	35
36	Low Impulsive Action, but not Impulsive Choice, Predicts Greater Conditioned Reinforcer Salience and Augmented Nucleus Accumbens Dopamine Release. <i>Neuropsychopharmacology</i> , 2016, 41, 2091-2100.	2.8	25

#	ARTICLE	IF	CITATIONS
37	The 5-HT <sub>2C</sub> receptor agonist lorcaserin reduces cocaine self-administration, reinstatement of cocaine-seeking and cocaine induced locomotor activity. <i>Neuropharmacology</i> , 2016, 101, 237-245.	2.0	59
38	The adrenergic $\alpha_2$ antagonist atipamezole alters the behavioural effects of pramipexole and increases pramipexole concentration in blood plasma. <i>Life Sciences</i> , 2016, 151, 300-304.	2.0	1
39	Characterization of the 5-HT <sub>2C</sub> receptor agonist lorcaserin on efficacy and safety measures in a rat model of diet-induced obesity. <i>Pharmacology Research and Perspectives</i> , 2015, 3, e00084.	1.1	25
40	Antipsychotics and Amotivation. <i>Neuropsychopharmacology</i> , 2015, 40, 1539-1548.	2.8	45
41	The Serotonin 2C Receptor Agonist Lorcaserin Attenuates Intracranial Self-Stimulation and Blocks the Reward-Enhancing Effects of Nicotine. <i>ACS Chemical Neuroscience</i> , 2015, 6, 1231-1240.	1.7	30
42	Therapeutic Potential of 5-HT <sub>2C</sub> Receptor Agonists for Addictive Disorders. <i>ACS Chemical Neuroscience</i> , 2015, 6, 1071-1088.	1.7	75
43	Behavioral effects of food-derived opioid-like peptides in rodents: Implications for schizophrenia?. <i>Pharmacology Biochemistry and Behavior</i> , 2015, 134, 70-78.	1.3	22
44	Low dose pramipexole causes D3 receptor-independent reduction of $\Delta$ locomotion and responding for a conditioned reinforcer. <i>Neuropharmacology</i> , 2015, 89, 225-231.	2.0	6
45	Chronic exposure to a gambling-like schedule of reward predictive stimuli can promote sensitization to amphetamine in rats. <i>Frontiers in Behavioral Neuroscience</i> , 2014, 8, 36.	1.0	93
46	Responding for a conditioned reinforcer, and its enhancement by nicotine, is blocked by dopamine receptor antagonists and a 5-HT <sub>2C</sub> receptor agonist but not by a 5-HT <sub>2A</sub> receptor antagonist. <i>Pharmacology Biochemistry and Behavior</i> , 2014, 125, 40-47.	1.3	18
47	A Dopamine D2 Receptor-DISC1 Protein Complex may Contribute to Antipsychotic-Like Effects. <i>Neuron</i> , 2014, 84, 1302-1316.	3.8	91
48	Examination of the effects of varenicline, bupropion, lorcaserin, or naltrexone on responding for conditioned reinforcement in nicotine-exposed rats. <i>Behavioural Pharmacology</i> , 2014, 25, 775-783.	0.8	20
49	The effects of nicotine exposure during Pavlovian conditioning in rats on several measures of incentive motivation for a conditioned stimulus paired with water. <i>Psychopharmacology</i> , 2014, 231, 2261-2271.	1.5	20
50	Oxidative stress and the antipsychotic-induced vacuous chewing movement model of tardive dyskinesia: evidence for antioxidant-based prevention strategies. <i>Psychopharmacology</i> , 2014, 231, 2237-2249.	1.5	28
51	Responding for conditioned reinforcement in C57BL/6 and CD-1 mice, and Sprague-Dawley rats: Effects of methylphenidate and amphetamine. <i>Psychopharmacology</i> , 2014, 231, 4503-4516.	1.5	8
52	Effects of intracerebroventricular (ICV) olanzapine on insulin sensitivity and secretion in vivo: An animal model. <i>European Neuropsychopharmacology</i> , 2014, 24, 448-458.	0.3	18
53	Nicotine-induced enhancement of responding for conditioned reinforcement in rats: Role of prior nicotine exposure and $\alpha_4\beta_2$ nicotinic receptors. <i>Psychopharmacology</i> , 2013, 225, 429-440.	1.5	26
54	Impulsive action in the 5-choice serial reaction time test in 5-HT <sub>2C</sub> receptor null mutant mice. <i>Psychopharmacology</i> , 2013, 226, 561-570.	1.5	35

#	ARTICLE	IF	CITATIONS
55	From obesity to substance abuse: therapeutic opportunities for 5-HT <sub>2C</sub> receptor agonists. <i>Trends in Pharmacological Sciences</i> , 2013, 34, 560-570.	4.0	90
56	Antagonizing 5-HT <sub>2A</sub> receptors with M100907 and stimulating 5-HT <sub>2C</sub> receptors with Ro60-0175 blocks cocaine-induced locomotion and zif268 mRNA expression in Sprague-Dawley rats. <i>Behavioural Brain Research</i> , 2013, 240, 171-181.	1.2	20
57	Deep brain stimulation of the subthalamic nucleus increases premature responding in a rat gambling task. <i>Behavioural Brain Research</i> , 2013, 245, 76-82.	1.2	25
58	Chronic olanzapine administration in rats: Effect of route of administration on weight, food intake and body composition. <i>Pharmacology Biochemistry and Behavior</i> , 2013, 103, 717-722.	1.3	19
59	Evaluation of chemically diverse 5-HT <sub>2C</sub> receptor agonists on behaviours motivated by food and nicotine and on side effect profiles. <i>Psychopharmacology</i> , 2013, 226, 475-490.	1.5	51
60	Disrupted-In-Schizophrenia-1 Gln31Leu Polymorphism Results in Social Anhedonia Associated with Monoaminergic Imbalance and Reduction of CREB and $\beta$ -arrestin-1,2 in the Nucleus Accumbens in a Mouse Model of Depression. <i>Neuropsychopharmacology</i> , 2013, 38, 423-436.	2.8	51
61	Acute Effects of Single-Dose Olanzapine on Metabolic, Endocrine, and Inflammatory Markers in Healthy Controls. <i>Journal of Clinical Psychopharmacology</i> , 2013, 33, 740-746.	0.7	67
62	Adolescent Cocaine Exposure Causes Enduring Macroscale Changes in Mouse Brain Structure. <i>Journal of Neuroscience</i> , 2013, 33, 1797-1803.	1.7	38
63	Double Dissociation between Regulation of Conditioned Disgust and Taste Avoidance by Serotonin Availability at the 5-HT <sub>3</sub> Receptor in the Posterior and Anterior Insular Cortex. <i>Journal of Neuroscience</i> , 2012, 32, 13709-13717.	1.7	60
64	Contribution of Decreased Serotonin Release to the Antidyskinetic Effects of Deep Brain Stimulation in a Rodent Model of Tardive Dyskinesia: Comparison of the Subthalamic and Entopeduncular Nuclei. <i>Journal of Neuroscience</i> , 2012, 32, 9574-9581.	1.7	56
65	Effects of the 5-HT <sub>2C</sub> receptor agonist Ro60-0175 and the 5-HT <sub>2A</sub> receptor antagonist M100907 on nicotine self-administration and reinstatement. <i>Neuropharmacology</i> , 2012, 62, 2288-2298.	2.0	65
66	Cannabidiol, a non- $\psi$ psychotropic component of cannabis, attenuates vomiting and nausea-like behaviour via indirect agonism of 5-HT <sub>1A</sub> somatodendritic autoreceptors in the dorsal raphe nucleus. <i>British Journal of Pharmacology</i> , 2012, 165, 2620-2634.	2.7	202
67	Age and sex differences in impulsive action in rats: The role of dopamine and glutamate. <i>Behavioural Brain Research</i> , 2012, 230, 21-33.	1.2	68
68	The 5-HT <sub>2C</sub> Receptor Agonist Lorcaserin Reduces Nicotine Self-Administration, Discrimination, and Reinstatement: Relationship to Feeding Behavior and Impulse Control. <i>Neuropsychopharmacology</i> , 2012, 37, 1177-1191.	2.8	122
69	Reply to: Electrical Brain Stimulation in Depression: Which Target(s)? <i>Biological Psychiatry</i> , 2011, 69, e7-e8.	0.7	4
70	Atypical antipsychotics and effects of muscarinic, serotonergic, dopaminergic and histaminergic receptor binding on insulin secretion in vivo: An animal model. <i>Schizophrenia Research</i> , 2011, 131, 90-95.	1.1	67
71	Serotonin and Reward-Related Behavior: Focus on 5-HT <sub>2C</sub> Receptors. <i>Receptors</i> , 2011, , 293-324.	0.2	6
72	Impulsive action induced by amphetamine, cocaine and MK801 is reduced by 5-HT <sub>2C</sub> receptor stimulation and 5-HT <sub>2A</sub> receptor blockade. <i>Neuropharmacology</i> , 2011, 61, 468-477.	2.0	90

#	ARTICLE	IF	CITATIONS
73	Enhanced Incentive Motivation for Sucrose-Paired Cues in Adolescent Rats: Possible Roles for Dopamine and Opioid Systems. <i>Neuropsychopharmacology</i> , 2011, 36, 1631-1643.	2.8	22
74	Early-life maternal separation and social isolation produce an increase in impulsive action but not impulsive choice.. <i>Behavioral Neuroscience</i> , 2011, 125, 481-491.	0.6	62
75	The effects of adolescent methylphenidate self-administration on responding for a conditioned reward, amphetamine-induced locomotor activity, and neuronal activation. <i>Psychopharmacology</i> , 2010, 208, 455-468.	1.5	19
76	Genetic and pharmacological evidence that 5-HT <sub>2C</sub> receptor activation, but not inhibition, affects motivation to feed under a progressive ratio schedule of reinforcement. <i>Pharmacology Biochemistry and Behavior</i> , 2010, 97, 170-178.	1.3	26
77	Enhanced dopamine function in DISC1 $\Delta$ 100P mutant mice: implications for schizophrenia. <i>Genes, Brain and Behavior</i> , 2010, 9, 777-789.	1.1	89
78	Uncoupling the dopamine D1-D2 receptor complex exerts antidepressant-like effects. <i>Nature Medicine</i> , 2010, 16, 1393-1395.	15.2	158
79	The Dopamine D1-D2 Receptor Heteromer Localizes in Dynorphin/Enkephalin Neurons. <i>Journal of Biological Chemistry</i> , 2010, 285, 36625-36634.	1.6	162
80	Antidepressant-Like Effects of Medial Prefrontal Cortex Deep Brain Stimulation in Rats. <i>Biological Psychiatry</i> , 2010, 67, 117-124.	0.7	284
81	Schizophrenia, amphetamine-induced sensitized state and acute amphetamine exposure all show a common alteration: increased dopamine D2 receptor dimerization. <i>Molecular Brain</i> , 2010, 3, 25.	1.3	79
82	The Nicotinic Acetylcholine Receptor $\alpha$ 5 Subunit Plays a Key Role in Attention Circuitry and Accuracy. <i>Journal of Neuroscience</i> , 2010, 30, 9241-9252.	1.7	132
83	Gestational treatment with methylazoxymethanol (MAM) that disrupts hippocampal-dependent memory does not alter behavioural response to cocaine. <i>Pharmacology Biochemistry and Behavior</i> , 2009, 93, 382-390.	1.3	13
84	The role of noradrenaline and 5-hydroxytryptamine in yohimbine-induced increases in alcohol-seeking in rats. <i>Psychopharmacology</i> , 2009, 204, 477-488.	1.5	48
85	Effects of 5-HT depletion in the frontal cortex or nucleus accumbens on response inhibition measured in the 5-choice serial reaction time test and on a DRL schedule. <i>Behavioural Brain Research</i> , 2009, 201, 88-98.	1.2	32
86	Characterizing the effects of 5-HT <sub>2C</sub> receptor ligands on motor activity and feeding behaviour in 5-HT <sub>2C</sub> receptor knockout mice. <i>Neuropharmacology</i> , 2009, 57, 259-267.	2.0	71
87	Insulin resistance and secretion in vivo: Effects of different antipsychotics in an animal model. <i>Schizophrenia Research</i> , 2009, 108, 127-133.	1.1	106
88	Abnormalities in brain structure and behavior in GSK-3 $\alpha$ mutant mice. <i>Molecular Brain</i> , 2009, 2, 35.	1.3	162
89	Intra-median raphe nucleus (MRN) infusions of muscimol, a GABA-A receptor agonist, reinstate alcohol seeking in rats: role of impulsivity and reward. <i>Psychopharmacology</i> , 2008, 195, 605-615.	1.5	37
90	Characterization of methylphenidate self-administration and reinstatement in the rat. <i>Psychopharmacology</i> , 2008, 199, 55-66.	1.5	39

#	ARTICLE	IF	CITATIONS
91	Reduced fear and aggression and altered serotonin metabolism in <i>Gtf2ird1</i> -targeted mice. <i>Genes, Brain and Behavior</i> , 2008, 7, 224-234.	1.1	125
92	A sensitizing regimen of amphetamine that disrupts attentional set-shifting does not disrupt working or long-term memory. <i>Behavioural Brain Research</i> , 2008, 189, 170-179.	1.2	55
93	Serotonin receptors as potential targets for modulation of nicotine use and dependence. <i>Progress in Brain Research</i> , 2008, 172, 361-383.	0.9	50
94	The 5-HT <sub>2C</sub> Receptor Agonist Ro60-0175 Reduces Cocaine Self-Administration and Reinstatement Induced by the Stressor Yohimbine, and Contextual Cues. <i>Neuropsychopharmacology</i> , 2008, 33, 1402-1412.	2.8	107
95	Insulin Resistance and Decreased Glucose-Stimulated Insulin Secretion After Acute Olanzapine Administration. <i>Journal of Clinical Psychopharmacology</i> , 2008, 28, 494-499.	0.7	86
96	Time Course of the Antipsychotic Effect and the Underlying Behavioral Mechanisms. <i>Neuropsychopharmacology</i> , 2007, 32, 263-272.	2.8	65
97	A Sensitizing Regimen of Amphetamine Impairs Visual Attention in the 5-Choice Serial Reaction Time Test: Reversal by a D1 Receptor Agonist Injected into the Medial Prefrontal Cortex. <i>Neuropsychopharmacology</i> , 2007, 32, 1122-1132.	2.8	69
98	Gestational Methylazoxymethanol Acetate Treatment Impairs Select Cognitive Functions: Parallels to Schizophrenia. <i>Neuropsychopharmacology</i> , 2007, 32, 483-492.	2.8	104
99	The amphetamine-induced sensitized state as a model of schizophrenia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2007, 31, 1556-1571.	2.5	186
100	Dopamine transporter cell surface localization facilitated by a direct interaction with the dopamine D2 receptor. <i>EMBO Journal</i> , 2007, 26, 2127-2136.	3.5	182
101	Opposing effects of 5-HT <sub>2A</sub> and 5-HT <sub>2C</sub> receptor antagonists in the rat and mouse on premature responding in the five-choice serial reaction time test. <i>Psychopharmacology</i> , 2007, 195, 223-234.	1.5	185
102	Effects of dexfenfluramine and 5-HT <sub>3</sub> receptor antagonists on stress-induced reinstatement of alcohol seeking in rats. <i>Psychopharmacology</i> , 2006, 186, 82-92.	1.5	47
103	The effects of the 5-HT <sub>2C</sub> receptor antagonist SB242084 on locomotor activity induced by selective, or mixed, indirect serotonergic and dopaminergic agonists. <i>Psychopharmacology</i> , 2006, 187, 515-525.	1.5	82
104	Effects of central neurokinin-1 receptor antagonism on cocaine- and opiate-induced locomotor activity and self-administration behaviour in rats. <i>Pharmacology Biochemistry and Behavior</i> , 2006, 84, 94-101.	1.3	21
105	Early life tactile stimulation changes adult rat responsiveness to amphetamine. <i>Pharmacology Biochemistry and Behavior</i> , 2006, 84, 497-503.	1.3	40
106	Dissociation between In Vivo Occupancy and Functional Antagonism of Dopamine D2 Receptors: Comparing Aripiprazole to Other Antipsychotics in Animal Models. <i>Neuropsychopharmacology</i> , 2006, 31, 1854-1863.	2.8	194
107	Sensitization to amphetamine, but not PCP, impairs attentional set shifting: reversal by a D1 receptor agonist injected into the medial prefrontal cortex. <i>Psychopharmacology</i> , 2005, 183, 190-200.	1.5	113
108	Sensitization to amphetamine, but not phencyclidine, disrupts prepulse inhibition and latent inhibition. <i>Psychopharmacology</i> , 2005, 180, 366-376.	1.5	74

#	ARTICLE	IF	CITATIONS
109	Activation of central neurokinin-1 receptors induces reinstatement of cocaine-seeking behavior. <i>Neuroscience Letters</i> , 2005, 390, 42-47.	1.0	20
110	Effects of injections of 8-hydroxy-2-(di-n-propylamino)tetralin or muscimol in the median raphe nucleus on c-fos mRNA in the rat brain. <i>Neuroscience</i> , 2005, 131, 475-479.	1.1	11
111	A putative animal model of the "prodromal" state of schizophrenia. <i>Biological Psychiatry</i> , 2005, 57, 586-593.	0.7	63
112	Injection of the 5-HT <sub>2C</sub> Receptor Agonist Ro60-0175 into the Ventral Tegmental Area Reduces Cocaine-Induced Locomotor Activity and Cocaine Self-Administration. <i>Neuropsychopharmacology</i> , 2004, 29, 308-318.	2.8	122
113	Evaluation of the motor initiation hypothesis of APD-induced conditioned avoidance decreases. <i>Pharmacology Biochemistry and Behavior</i> , 2004, 78, 811-819.	1.3	40
114	Fluoxetine, but not sertraline or citalopram, potentiates the locomotor stimulant effect of cocaine: possible pharmacokinetic effects. <i>Psychopharmacology</i> , 2004, 174, 406-13.	1.5	20
115	Infusion of the substance P analogue, DiMe-C7, into the ventral tegmental area induces reinstatement of cocaine-seeking behaviour in rats. <i>Psychopharmacology</i> , 2004, 177, 111-120.	1.5	38
116	5,7-Dihydroxytryptamine Lesions of the Dorsal and Median Raphe Nuclei Interfere With Lithium-Induced Conditioned Gaping, but Not Conditioned Taste Avoidance, in Rats.. <i>Behavioral Neuroscience</i> , 2004, 118, 1391-1399.	0.6	35
117	The 5-HT <sub>2A</sub> receptor antagonist M100,907 attenuates motor and 'impulsive-type' behaviours produced by NMDA receptor antagonism. <i>Psychopharmacology</i> , 2003, 170, 309-319.	1.5	162
118	Serotonin and drug reward: focus on 5-HT <sub>2C</sub> receptors. <i>European Journal of Pharmacology</i> , 2003, 480, 151-162.	1.7	147
119	Up-regulated dopamine D1 receptor binding can be detected in vivo following repeated SCH 23390, but not SKF 81297 or 6-hydroxydopamine, treatments. <i>European Journal of Pharmacology</i> , 2003, 459, 195-201.	1.7	6
120	Amphetamine-sensitized animals show a sensorimotor gating and neurochemical abnormality similar to that of schizophrenia. <i>Schizophrenia Research</i> , 2003, 64, 103-114.	1.1	86
121	Neonatal Ablation of the Nigrostriatal Dopamine Pathway Does Not Influence Limb Development in Rats. <i>Experimental Neurology</i> , 2002, 177, 547-556.	2.0	3
122	The Role of Corticotropin-Releasing Factor in the Median Raphe Nucleus in Relapse to Alcohol. <i>Journal of Neuroscience</i> , 2002, 22, 7844-7849.	1.7	127
123	Cyclooxygenase inhibitor modulation of dopamine-related behaviours. <i>European Journal of Pharmacology</i> , 2002, 450, 141-151.	1.7	29
124	Activation of 5-HT <sub>1B</sub> receptors in the nucleus accumbens reduces self-administration of amphetamine on a progressive ratio schedule. <i>Pharmacology Biochemistry and Behavior</i> , 2002, 71, 717-725.	1.3	49
125	Editorial. <i>Pharmacology Biochemistry and Behavior</i> , 2002, 71, 531-532.	1.3	0
126	Differential Effects of the 5-HT <sub>2A</sub> Receptor Antagonist M100,907 and the 5-HT <sub>2C</sub> Receptor Antagonist SB242,084 on Cocaine-induced Locomotor Activity, Cocaine Self-administration and Cocaine-induced Reinstatement of Responding. <i>Neuropsychopharmacology</i> , 2002, 27, 576-86.	2.8	210



#	ARTICLE	IF	CITATIONS
127	Multiple 5-HT receptors are involved in the effects of acute MDMA treatment: studies on locomotor activity and responding for conditioned reinforcement. <i>Psychopharmacology</i> , 2002, 162, 282-291.	1.5	76
128	The adenosine A2A agonist CGS 21680 reverses the reduction in prepulse inhibition of the acoustic startle response induced by phencyclidine, but not by apomorphine and amphetamine. <i>Psychopharmacology</i> , 2001, 156, 187-193.	1.5	20
129	Extracellular amino acid profiles in the paraventricular nucleus of the rat hypothalamus are influenced by diet composition. <i>Brain Research</i> , 2001, 892, 320-328.	1.1	34
130	Reduced Brain Serotonin Activity Disrupts Prepulse Inhibition of the Acoustic Startle Reflex Effects of 5,7-dihydroxytryptamine and p-chlorophenylalanine. <i>Neuropsychopharmacology</i> , 2001, 24, 399-409.	2.8	61
131	Pre-Exposure to (±)3,4-methylenedioxy-methamphetamine (MDMA) Facilitates Acquisition of Intravenous Cocaine Self-Administration in Rats. <i>Neuropsychopharmacology</i> , 2001, 25, 195-203.	2.8	41
132	Subchronic fluoxetine treatment induces a transient potentiation of amphetamine-induced hyperlocomotion: possible pharmacokinetic interaction. <i>Behavioural Pharmacology</i> , 2000, 11, 109-116.	0.8	15
133	Modification of dopamine D1 receptor knockout phenotype in mice lacking both dopamine D1 and D3 receptors. <i>European Journal of Pharmacology</i> , 2000, 399, 171-181.	1.7	48
134	Antipsychoticlike effects of amoxapine, without catalepsy, using the prepulse inhibition of the acoustic startle reflex test in rats. <i>Biological Psychiatry</i> , 2000, 47, 670-676.	0.7	11
135	The rewarding properties of neuropeptide Y in perifornical hypothalamus vs. nucleus accumbens. <i>Peptides</i> , 2000, 21, 1279-1287.	1.2	65
136	Dietary protein content affects the profiles of extracellular amino acids in the medial preoptic area of freely moving rats. <i>Life Sciences</i> , 2000, 66, 1105-1118.	2.0	24
137	Acute fluoxetine treatment potentiates amphetamine hyperactivity and amphetamine-induced nucleus accumbens dopamine release: possible pharmacokinetic interaction. <i>Psychopharmacology</i> , 1999, 141, 421-427.	1.5	31
138	Activation of 5-HT 1B receptors in the nucleus accumbens reduces amphetamine-induced enhancement of responding for conditioned reward. <i>Psychopharmacology</i> , 1999, 142, 165-174.	1.5	63
139	The potentiating effect of sertraline and fluoxetine on amphetamine-induced locomotor activity is not mediated by serotonin. <i>Psychopharmacology</i> , 1999, 143, 426-432.	1.5	26
140	Depletion of brain serotonin following intra-raphe injections. <i>Psychopharmacology</i> , 1999, 146, 185-193.	1.5	28
141	Selective destruction of brain serotonin neurons by 5,7-dihydroxytryptamine increases responding for a conditioned reward. <i>Psychopharmacology</i> , 1999, 147, 291-299.	1.5	75
142	The adenosine A1 receptor agonist N6-cyclopentyladenosine blocks the disruptive effect of phencyclidine on prepulse inhibition of the acoustic startle response in the rat. <i>European Journal of Pharmacology</i> , 1999, 369, 325-329.	1.7	30
143	Spatial learning deficit in dopamine D1 receptor knockout mice. <i>European Journal of Pharmacology</i> , 1999, 383, 95-106.	1.7	153
144	Measurement of Blood-Brain Barrier Permeability of Rats with Alpha-Aminoisobutyric Acid During Microdialysis. <i>Physiology and Behavior</i> , 1999, 67, 587-598.	1.0	12

#	ARTICLE	IF	CITATIONS
145	RU-24969 disrupts d-amphetamine self-administration and responding for conditioned reward via stimulation of 5-HT1B receptors. <i>Behavioural Pharmacology</i> , 1999, 10, 183-193.	0.8	30
146	Reversal of fenfluramine and fluoxetine anorexia by 8-OH-DPAT is attenuated following raphe injection of 5,7-dihydroxytryptamine. <i>Brain Research</i> , 1998, 800, 62-68.	1.1	21
147	Injections of d-amphetamine into the ventral pallidum increase locomotor activity and responding for conditioned reward: a comparison with injections into the nucleus accumbens. <i>Brain Research</i> , 1998, 805, 29-40.	1.1	35
148	Disruption of dopamine D1 receptor gene expression attenuates alcohol-seeking behavior. <i>European Journal of Pharmacology</i> , 1998, 353, 149-158.	1.7	154
149	Acute Amino Acid Loads That Deplete Brain Serotonin Fail to Alter Behavior. <i>Pharmacology Biochemistry and Behavior</i> , 1998, 59, 115-121.	1.3	26
150	A Comparison of the Effects of Risperidone, Raclopride, and Ritanserin on Intravenous Self-Administration of d-Amphetamine. <i>Pharmacology Biochemistry and Behavior</i> , 1998, 60, 55-60.	1.3	19
151	Reinstatement of alcohol-seeking by priming injections of alcohol and exposure to stress in rats. <i>Psychopharmacology</i> , 1998, 135, 169-174.	1.5	288
152	Neuropeptide Y-induced operant responding for sucrose is not mediated by dopamine. <i>Peptides</i> , 1998, 19, 1667-1673.	1.2	38
153	Fluoxetine attenuates morphine-induced locomotion and blocks morphine-sensitization. <i>European Journal of Pharmacology</i> , 1997, 337, 161-164.	1.7	24
154	Differential effects of ondansetron and $\pm$ -flupenthixol on responding for conditioned reward. <i>Psychopharmacology</i> , 1997, 134, 64-72.	1.5	22
155	Injection of 5-HT into the nucleus accumbens reduces the effects of d-amphetamine on responding for conditioned reward. <i>Psychopharmacology</i> , 1996, 126, 62-69.	1.5	39
156	Effects of d-fenfluramine and metergoline on responding for conditioned reward and the response potentiating effect of nucleus accumbens-d-amphetamine. <i>Psychopharmacology</i> , 1995, 118, 155-163.	1.5	46
157	Further studies to examine the nature of dexfenfluramine-induced suppression of heroin self-administration. <i>Psychopharmacology</i> , 1995, 120, 134-141.	1.5	26
158	Median raphe injections of 8-OH-DPAT lower frequency thresholds for lateral hypothalamic self-stimulation. <i>Pharmacology Biochemistry and Behavior</i> , 1995, 52, 65-71.	1.3	54
159	Effects of combined or separate 5,7-dihydroxytryptamine lesions of the dorsal and median raphe nuclei on responding maintained by a DRL 20s schedule of food reinforcement. <i>Brain Research</i> , 1995, 675, 45-54.	1.1	76
160	Median and dorsal raphe injections of the 5-HT1A agonist, 8-OH-DPAT, and the GABAA agonist, muscimol, increase voluntary ethanol intake in wistar rats. <i>Neuropharmacology</i> , 1994, 33, 349-358.	2.0	79
161	Injecting 5-HT into the PVN does not prevent feeding induced by injecting 8-OH-DPAT into the raphe. <i>Pharmacology Biochemistry and Behavior</i> , 1993, 46, 487-491.	1.3	39
162	Conditioned place preference induced by microinjection of 8-OH-DPAT into the dorsal or median raphe nucleus. <i>Psychopharmacology</i> , 1993, 113, 31-36.	1.5	64

#	ARTICLE	IF	CITATIONS
163	Radiofrequency lesions of the PVN fail to modify the effects of serotonergic drugs on food intake. <i>Brain Research</i> , 1993, 630, 1-9.	1.1	29
164	A comparison of the effects of dorsal or median raphe injections of 8-OH-DPAT in three operant tasks measuring response inhibition. <i>Behavioural Brain Research</i> , 1993, 54, 187-197.	1.2	59
165	A comparison of the effects of the 5-HT <sub>1</sub> agonists TFMPP and RU 24969 on feeding following peripheral or medial hypothalamic injection. <i>Brain Research</i> , 1992, 580, 265-272.	1.1	27
166	Effect of drugs influencing 5-HT function on ethanol drinking and feeding behaviour in rats: Studies using a drinkometer system. <i>Neuroscience and Biobehavioral Reviews</i> , 1992, 16, 535-552.	2.9	88
167	The polyamine synthesis inhibitor $\hat{\pm}$ -difluoromethylornithine is neuroprotective against N-methyl-D-aspartate-induced brain damage in vivo. <i>European Journal of Pharmacology</i> , 1991, 209, 101-103.	1.7	24
168	Dopamine receptor blockade in nucleus accumbens or caudate nucleus differentially affects feeding induced by 8-OH-DPAT injected into dorsal or median raphe. <i>Brain Research</i> , 1991, 552, 181-189.	1.1	64
169	Opiate antagonists inhibit feeding induced by 8-OH-DPAT: possible mediation in the nucleus accumbens. <i>Brain Research</i> , 1991, 560, 260-267.	1.1	21
170	Influence of taste and food texture on the feeding responses induced by 8-OH-DPAT and gepirone. <i>Psychopharmacology</i> , 1991, 104, 302-306.	1.5	14
171	Effects of 8-OH-DPAT, buspirone and ICS 205-930 on feeding in a novel environment: comparisons with chlordiazepoxide and FG 7142. <i>Psychopharmacology</i> , 1990, 102, 301-308.	1.5	28
172	A pharmacological analysis of the eating response induced by 8-OH-DPAT injected into the dorsal raphe nucleus reveals the involvement of a dopaminergic mechanism. <i>Psychopharmacology</i> , 1990, 100, 188-194.	1.5	33
173	Dorsal raphe microinjection of 5-HT and indirect 5-HT agonists induces feeding in rats. <i>European Journal of Pharmacology</i> , 1990, 184, 265-271.	1.7	33
174	The involvement of 5-hydroxytryptaminergic and dopaminergic mechanisms in the eating induced by buspirone, gepirone and ipsapirone. <i>British Journal of Pharmacology</i> , 1990, 99, 519-525.	2.7	36
175	A comparison of the effects of tryptamine and 5-hydroxytryptamine on feeding following injection into the paraventricular nucleus of the hypothalamus. <i>Pharmacology Biochemistry and Behavior</i> , 1989, 32, 907-911.	1.3	28
176	Enhancement of 5-HT-induced anorexia: a test of the reversibility of monoamine oxidase inhibitors. <i>Psychopharmacology</i> , 1989, 98, 265-268.	1.5	3
177	Tryptamine impairs the acquisition of a one-way active avoidance task. <i>Pharmacology Biochemistry and Behavior</i> , 1989, 32, 317-321.	1.3	7
178	Octopamine injected into the paraventricular nucleus induces eating in rats: a comparison with noradrenaline-induced eating. <i>British Journal of Pharmacology</i> , 1989, 97, 483-489.	2.7	13
179	Increased food intake in satiated rats induced by the 5-HT antagonists methysergide, metergoline and ritanserin. <i>Psychopharmacology</i> , 1988, 96, 237-42.	1.5	90
180	The effects of d-fenfluramine on saccharin intake and preference, and on food and water intake. <i>Pharmacology Biochemistry and Behavior</i> , 1988, 29, 687-691.	1.3	12

#	ARTICLE	IF	CITATIONS
181	8-OH-DPAT elicits gnawing, and eating of solid but not liquid foods. <i>Psychopharmacology</i> , 1987, 92, 192-5.	1.5	35
182	The anorectic action of peripheral 5-HT examined in the runway: evidence for an action on satiation. <i>Psychopharmacology</i> , 1987, 93, 498-501.	1.5	6
183	Dissociation of the anorectic actions of 5-HTP and fenfluramine. <i>Psychopharmacology</i> , 1986, 89, 216-220.	1.5	35
184	Behavioural and pharmacological investigations of 5-HT hypophagia and hyperdipsia. <i>Pharmacology Biochemistry and Behavior</i> , 1986, 25, 23-28.	1.3	20
185	Microstructural analysis of the anorectic action of peripherally administered 5-HT. <i>Pharmacology Biochemistry and Behavior</i> , 1986, 24, 1133-1136.	1.3	32
186	The anorectic action of peripherally administered 5-HT is enhanced by vagotomy. <i>Physiology and Behavior</i> , 1985, 34, 861-866.	1.0	29
187	Effects of manipulations of peripheral serotonin on feeding and drinking in the rat. <i>Pharmacology Biochemistry and Behavior</i> , 1984, 20, 835-840.	1.3	72