Paul J Fletcher

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1871693/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Effects of pimavanserin and lorcaserin on alcohol self-administration and reinstatement in male and female rats. Neuropharmacology, 2022, , 109150.	4.1	3
2	Prenatal disruption of D1R-SynGAP complex causes cognitive deficits in adulthood. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2021, 105, 110122.	4.8	3
3	The serotonin 2C receptor agonist lorcaserin, alone and in combination with the opioid receptor antagonist naltrexone, attenuates bingeâ€like ethanol drinking. Addiction Biology, 2021, 26, e13040.	2.6	9
4	Lorcaserin: A review of its preclinical and clinical pharmacology and therapeutic potential. , 2020, 205, 107417.		52
5	Clozapine reliably increases the motivation for food: parsing the role of the 5-HT2c and H1 receptors. Psychopharmacology, 2020, 237, 957-966.	3.1	9
6	The pharmacological stressor yohimbine, but not U50,488, increases responding for conditioned reinforcers paired with ethanol or sucrose. Psychopharmacology, 2020, 237, 3689-3702.	3.1	7
7	Serotonin transporter protein in autopsied brain of chronic users of cocaine. Psychopharmacology, 2020, 237, 2661-2671.	3.1	7
8	Effects of exposure to chronic uncertainty and a sensitizing regimen of amphetamine injections on locomotion, decision-making, and dopamine receptors in rats. Neuropsychopharmacology, 2020, 45, 811-822.	5.4	13
9	Median raphe serotonin neurons promote anxiety-like behavior via inputs to the dorsal hippocampus. Neuropharmacology, 2020, 168, 107985.	4.1	42
10	Disruption of SynGAP–dopamine D1 receptor complexes alters actin and microtubule dynamics and impairs GABAergic interneuron migration. Science Signaling, 2019, 12, .	3.6	11
11	Nicotine enhances responding for conditioned reinforcement via α4β2 nicotinic acetylcholine receptors in the ventral tegmental area, but not the nucleus accumbens or the prefrontal cortex. Neuropharmacology, 2019, 148, 68-76.	4.1	3
12	Kappa opioid receptors mediate yohimbine-induced increases in impulsivity in the 5-choice serial reaction time task. Behavioural Brain Research, 2019, 359, 258-265.	2.2	11
13	Dorsal raphe serotonin neurons inhibit operant responding for reward via inputs to the ventral tegmental area but not the nucleus accumbens: evidence from studies combining optogenetic stimulation and serotonin reuptake inhibition. Neuropsychopharmacology, 2019, 44, 793-804.	5.4	39
14	Preclinical evidence for combining the 5â€ <scp>HT_{2C}</scp> receptor agonist lorcaserin and varenicline as a treatment for nicotine dependence. Addiction Biology, 2019, 24, 376-387.	2.6	9
15	Clozapine reduces nicotine selfâ€administration, blunts reinstatement of nicotineâ€seeking but increases responding for food. Addiction Biology, 2019, 24, 565-576.	2.6	10
16	Adolescent exposure to Δ9-tetrahydrocannabinol delays acquisition of paired-associates learning in adulthood. Psychopharmacology, 2019, 236, 1875-1886.	3.1	23
17	Deep brain stimulation and fluoxetine exert different long-term changes in the serotonergic system. Neuropharmacology, 2018, 135, 63-72.	4.1	22
18	Acquisition of nicotine self-administration in amphetamine and phencyclidine models of schizophrenia: A role for stress?. Schizophrenia Research, 2018, 194, 98-106.	2.0	6

#	Article	IF	CITATIONS
19	Objective assessment of exploratory behaviour in schizophrenia using wireless motion capture. Schizophrenia Research, 2018, 195, 122-129.	2.0	3
20	Deep brain stimulation induces antidepressant-like effects in serotonin transporter knockout mice. Brain Stimulation, 2018, 11, 423-425.	1.6	17
21	Objective investigation of activity preference in schizophrenia: A pilot study. Psychiatry Research, 2018, 267, 551-559.	3.3	1
22	Effects of 5-HT1A, 5-HT2A and 5-HT2C receptor agonists and antagonists on responding for a conditioned reinforcer and its enhancement by methylphenidate. Psychopharmacology, 2017, 234, 889-902.	3.1	8
23	Pdxdc1 modulates prepulse inhibition of acoustic startle in the mouse. Translational Psychiatry, 2017, 7, e1125-e1125.	4.8	12
24	Studies To Examine Potential Tolerability Differences between the 5-HT _{2C} Receptor Selective Agonists Lorcaserin and CP-809101. ACS Chemical Neuroscience, 2017, 8, 1074-1084.	3.5	8
25	Role of impulsivity and reward in the anti-obesity actions of 5-HT _{2C} receptor agonists. Journal of Psychopharmacology, 2017, 31, 1403-1418.	4.0	30
26	Pharmacological Modulation of 5-HT2C Receptor Activity Produces Bidirectional Changes in Locomotor Activity, Responding for a Conditioned Reinforcer, and Mesolimbic DA Release in C57BL/6 Mice. Neuropsychopharmacology, 2017, 42, 2178-2187.	5.4	24
27	Lipoic acid and haloperidol-induced vacuous chewing movements: Implications for prophylactic antioxidant use in tardive dyskinesia. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2017, 72, 23-29.	4.8	10
28	Activation of Dopamine D1-D2 Receptor Complex Attenuates Cocaine Reward and Reinstatement of Cocaine-Seeking through Inhibition of DARPP-32, ERK, and ΔFosB. Frontiers in Pharmacology, 2017, 8, 924.	3.5	55
29	Uncertainty exposure causes behavioural sensitization and increases risky decision-making in male rats: toward modelling gambling disorder. Journal of Psychiatry and Neuroscience, 2017, 42, 404-413.	2.4	72
30	Decreased Incentive Motivation Following Knockout or Acute Blockade of the Serotonin Transporter: Role of the 5-HT2C Receptor. Neuropsychopharmacology, 2016, 41, 2566-2576.	5.4	22
31	Disruption of a dopamine receptor complex amplifies the actions of cocaine. European Neuropsychopharmacology, 2016, 26, 1366-1377.	0.7	36
32	Mgat5 modulates the effect of early life stress on adult behavior and physical health in mice. Behavioural Brain Research, 2016, 312, 253-264.	2.2	14
33	Atypical antipsychotics and effects on feeding: from mice to men. Psychopharmacology, 2016, 233, 2629-2653.	3.1	38
34	Responding for a conditioned reinforcer or unconditioned sensory reinforcer in mice: interactions with environmental enrichment, social isolation, and monoamine reuptake inhibitors. Psychopharmacology, 2016, 233, 983-993.	3.1	6
35	Lorcaserin and CP-809101 reduce motor impulsivity and reinstatement of food seeking behavior in male rats: Implications for understanding the anti-obesity property of 5-HT2C receptor agonists. Psychopharmacology, 2016, 233, 2841-2856.	3.1	35
36	Low Impulsive Action, but not Impulsive Choice, Predicts Greater Conditioned Reinforcer Salience and Augmented Nucleus Accumbens Dopamine Release. Neuropsychopharmacology, 2016, 41, 2091-2100.	5.4	25

#	Article	IF	CITATIONS
37	The 5-HT 2C receptor agonist lorcaserin reduces cocaine self-administration, reinstatement of cocaine-seeking and cocaine induced locomotor activity. Neuropharmacology, 2016, 101, 237-245.	4.1	59
38	The adrenergic $\hat{l}\pm 2$ antagonist atipamezole alters the behavioural effects of pramipexole and increases pramipexole concentration in blood plasma. Life Sciences, 2016, 151, 300-304.	4.3	1
39	Characterization of the 5â€HT _{2C} receptor agonist lorcaserin on efficacy and safety measures in a rat model of dietâ€induced obesity. Pharmacology Research and Perspectives, 2015, 3, e00084.	2.4	25
40	Antipsychotics and Amotivation. Neuropsychopharmacology, 2015, 40, 1539-1548.	5.4	45
41	The Serotonin 2C Receptor Agonist Lorcaserin Attenuates Intracranial Self-Stimulation and Blocks the Reward-Enhancing Effects of Nicotine. ACS Chemical Neuroscience, 2015, 6, 1231-1240.	3.5	30
42	Therapeutic Potential of 5-HT _{2C} Receptor Agonists for Addictive Disorders. ACS Chemical Neuroscience, 2015, 6, 1071-1088.	3.5	75
43	Behavioral effects of food-derived opioid-like peptides in rodents: Implications for schizophrenia?. Pharmacology Biochemistry and Behavior, 2015, 134, 70-78.	2.9	22
44	Low dose pramipexole causes D3 receptor-independent reduction ofÂlocomotion and responding for a conditioned reinforcer. Neuropharmacology, 2015, 89, 225-231.	4.1	6
45	Chronic exposure to a gambling-like schedule of reward predictive stimuli can promote sensitization to amphetamine in rats. Frontiers in Behavioral Neuroscience, 2014, 8, 36.	2.0	93
46	Responding for a conditioned reinforcer, and its enhancement by nicotine, is blocked by dopamine receptor antagonists and a 5-HT 2C receptor agonist but not by a 5-HT 2A receptor antagonist. Pharmacology Biochemistry and Behavior, 2014, 125, 40-47.	2.9	18
47	A Dopamine D2 Receptor-DISC1 Protein Complex may Contribute to Antipsychotic-Like Effects. Neuron, 2014, 84, 1302-1316.	8.1	91
48	Examination of the effects of varenicline, bupropion, lorcaserin, or naltrexone on responding for conditioned reinforcement in nicotine-exposed rats. Behavioural Pharmacology, 2014, 25, 775-783.	1.7	20
49	The effects of nicotine exposure during Pavlovian conditioning in rats on several measures of incentive motivation for a conditioned stimulus paired with water. Psychopharmacology, 2014, 231, 2261-2271.	3.1	20
50	Oxidative stress and the antipsychotic-induced vacuous chewing movement model of tardive dyskinesia: evidence for antioxidant-based prevention strategies. Psychopharmacology, 2014, 231, 2237-2249.	3.1	28
51	Responding for conditioned reinforcement in C57BL/6 and CD-1 mice, and Sprague-Dawley rats: Effects of methylphenidate and amphetamine. Psychopharmacology, 2014, 231, 4503-4516.	3.1	8
52	Effects of intracerebroventricular (ICV) olanzapine on insulin sensitivity and secretion in vivo: An animal model. European Neuropsychopharmacology, 2014, 24, 448-458.	0.7	18
53	Nicotine-induced enhancement of responding for conditioned reinforcement in rats: Role of prior nicotine exposure and α4β2 nicotinic receptors. Psychopharmacology, 2013, 225, 429-440.	3.1	26
54	Impulsive action in the 5-choice serial reaction time test in 5-HT2C receptor null mutant mice. Psychopharmacology, 2013, 226, 561-570.	3.1	35

#	Article	IF	CITATIONS
55	From obesity to substance abuse: therapeutic opportunities for 5-HT2C receptor agonists. Trends in Pharmacological Sciences, 2013, 34, 560-570.	8.7	90
56	Antagonizing 5-HT2A receptors with M100907 and stimulating 5-HT2C receptors with Ro60-0175 blocks cocaine-induced locomotion and zif268 mRNA expression in Sprague-Dawley rats. Behavioural Brain Research, 2013, 240, 171-181.	2.2	20
57	Deep brain stimulation of the subthalamic nucleus increases premature responding in a rat gambling task. Behavioural Brain Research, 2013, 245, 76-82.	2.2	25
58	Chronic olanzapine administration in rats: Effect of route of administration on weight, food intake and body composition. Pharmacology Biochemistry and Behavior, 2013, 103, 717-722.	2.9	19
59	Evaluation of chemically diverse 5-HT2C receptor agonists on behaviours motivated by food and nicotine and on side effect profiles. Psychopharmacology, 2013, 226, 475-490.	3.1	51
60	Disrupted-In-Schizophrenia-1 Gln31Leu Polymorphism Results in Social Anhedonia Associated with Monoaminergic Imbalance and Reduction of CREB and β-arrestin-1,2 in the Nucleus Accumbens in a Mouse Model of Depression. Neuropsychopharmacology, 2013, 38, 423-436.	5.4	51
61	Acute Effects of Single-Dose Olanzapine on Metabolic, Endocrine, and Inflammatory Markers in Healthy Controls. Journal of Clinical Psychopharmacology, 2013, 33, 740-746.	1.4	67
62	Adolescent Cocaine Exposure Causes Enduring Macroscale Changes in Mouse Brain Structure. Journal of Neuroscience, 2013, 33, 1797-1803.	3.6	38
63	Double Dissociation between Regulation of Conditioned Disgust and Taste Avoidance by Serotonin Availability at the 5-HT ₃ Receptor in the Posterior and Anterior Insular Cortex. Journal of Neuroscience, 2012, 32, 13709-13717.	3.6	60
64	Contribution of Decreased Serotonin Release to the Antidyskinetic Effects of Deep Brain Stimulation in a Rodent Model of Tardive Dyskinesia: Comparison of the Subthalamic and Entopeduncular Nuclei. Journal of Neuroscience, 2012, 32, 9574-9581.	3.6	56
65	Effects of the 5-HT2C receptor agonist Ro60-0175 and the 5-HT2A receptor antagonist M100907 on nicotine self-administration and reinstatement. Neuropharmacology, 2012, 62, 2288-2298.	4.1	65
66	Cannabidiol, a nonâ€psychotropic component of cannabis, attenuates vomiting and nauseaâ€like behaviour via indirect agonism of 5â€HT _{1A} somatodendritic autoreceptors in the dorsal raphe nucleus. British Journal of Pharmacology, 2012, 165, 2620-2634.	5.4	202
67	Age and sex differences in impulsive action in rats: The role of dopamine and glutamate. Behavioural Brain Research, 2012, 230, 21-33.	2.2	68
68	The 5-HT2C Receptor Agonist Lorcaserin Reduces Nicotine Self-Administration, Discrimination, and Reinstatement: Relationship to Feeding Behavior and Impulse Control. Neuropsychopharmacology, 2012, 37, 1177-1191.	5.4	122
69	Reply to: Electrical Brain Stimulation in Depression: Which Target(s)?. Biological Psychiatry, 2011, 69, e7-e8.	1.3	4
70	Atypical antipsychotics and effects of muscarinic, serotonergic, dopaminergic and histaminergic receptor binding on insulin secretion in vivo: An animal model. Schizophrenia Research, 2011, 131, 90-95.	2.0	67
71	Serotonin and Reward-Related Behavior: Focus on 5-HT2C Receptors. Receptors, 2011, , 293-324.	0.2	6
72	Impulsive action induced by amphetamine, cocaine and MK801 is reduced by 5-HT2C receptor stimulation and 5-HT2A receptor blockade. Neuropharmacology, 2011, 61, 468-477.	4.1	90

#	Article	IF	CITATIONS
73	Enhanced Incentive Motivation for Sucrose-Paired Cues in Adolescent Rats: Possible Roles for Dopamine and Opioid Systems. Neuropsychopharmacology, 2011, 36, 1631-1643.	5.4	22
74	Early-life maternal separation and social isolation produce an increase in impulsive action but not impulsive choice Behavioral Neuroscience, 2011, 125, 481-491.	1.2	62
75	The effects of adolescent methylphenidate self-administration on responding for a conditioned reward, amphetamine-induced locomotor activity, and neuronal activation. Psychopharmacology, 2010, 208, 455-468.	3.1	19
76	Genetic and pharmacological evidence that 5-HT2C receptor activation, but not inhibition, affects motivation to feed under a progressive ratio schedule of reinforcement. Pharmacology Biochemistry and Behavior, 2010, 97, 170-178.	2.9	26
77	Enhanced dopamine function in DISC1â€L100P mutant mice: implications for schizophrenia. Genes, Brain and Behavior, 2010, 9, 777-789.	2.2	89
78	Uncoupling the dopamine D1-D2 receptor complex exerts antidepressant-like effects. Nature Medicine, 2010, 16, 1393-1395.	30.7	158
79	The Dopamine D1-D2 Receptor Heteromer Localizes in Dynorphin/Enkephalin Neurons. Journal of Biological Chemistry, 2010, 285, 36625-36634.	3.4	162
80	Antidepressant-Like Effects of Medial Prefrontal Cortex Deep Brain Stimulation in Rats. Biological Psychiatry, 2010, 67, 117-124.	1.3	284
81	Schizophrenia, amphetamine-induced sensitized state and acute amphetamine exposure all show a common alteration: increased dopamine D2 receptor dimerization. Molecular Brain, 2010, 3, 25.	2.6	79
82	The Nicotinic Acetylcholine Receptor Â5 Subunit Plays a Key Role in Attention Circuitry and Accuracy. Journal of Neuroscience, 2010, 30, 9241-9252.	3.6	132
83	Gestational treatment with methylazoxymethanol (MAM) that disrupts hippocampal-dependent memory does not alter behavioural response to cocaine. Pharmacology Biochemistry and Behavior, 2009, 93, 382-390.	2.9	13
84	The role of noradrenaline and 5-hydroxytryptamine in yohimbine-induced increases in alcohol-seeking in rats. Psychopharmacology, 2009, 204, 477-488.	3.1	48
85	Effects of 5-HT depletion in the frontal cortex or nucleus accumbens on response inhibition measured in the 5-choice serial reaction time test and on a DRL schedule. Behavioural Brain Research, 2009, 201, 88-98.	2.2	32
86	Characterizing the effects of 5-HT2C receptor ligands on motor activity and feeding behaviour in 5-HT2C receptor knockout mice. Neuropharmacology, 2009, 57, 259-267.	4.1	71
87	Insulin resistance and secretion in vivo: Effects of different antipsychotics in an animal model. Schizophrenia Research, 2009, 108, 127-133.	2.0	106
88	Abnormalities in brain structure and behavior in GSK-3alpha mutant mice. Molecular Brain, 2009, 2, 35.	2.6	162
89	Intra-median raphe nucleus (MRN) infusions of muscimol, a GABA-A receptor agonist, reinstate alcohol seeking in rats: role of impulsivity and reward. Psychopharmacology, 2008, 195, 605-615.	3.1	37
90	Characterization of methylphenidate self-administration and reinstatement in the rat. Psychopharmacology, 2008, 199, 55-66.	3.1	39

#	Article	IF	CITATIONS
91	Reduced fear and aggression and altered serotonin metabolism in <i>Gtf2ird1</i> â€ŧargeted mice. Genes, Brain and Behavior, 2008, 7, 224-234.	2.2	125
92	A sensitizing regimen of amphetamine that disrupts attentional set-shifting does not disrupt working or long-term memory. Behavioural Brain Research, 2008, 189, 170-179.	2.2	55
93	Serotonin receptors as potential targets for modulation of nicotine use and dependence. Progress in Brain Research, 2008, 172, 361-383.	1.4	50
94	The 5-HT2C Receptor Agonist Ro60-0175 Reduces Cocaine Self-Administration and Reinstatement Induced by the Stressor Yohimbine, and Contextual Cues. Neuropsychopharmacology, 2008, 33, 1402-1412.	5.4	107
95	Insulin Resistance and Decreased Glucose-Stimulated Insulin Secretion After Acute Olanzapine Administration. Journal of Clinical Psychopharmacology, 2008, 28, 494-499.	1.4	86
96	Time Course of the Antipsychotic Effect and the Underlying Behavioral Mechanisms. Neuropsychopharmacology, 2007, 32, 263-272.	5.4	65
97	A Sensitizing Regimen of Amphetamine Impairs Visual Attention in the 5-Choice Serial Reaction Time Test: Reversal by a D1 Receptor Agonist Injected into the Medial Prefrontal Cortex. Neuropsychopharmacology, 2007, 32, 1122-1132.	5.4	69
98	Gestational Methylazoxymethanol Acetate Treatment Impairs Select Cognitive Functions: Parallels to Schizophrenia. Neuropsychopharmacology, 2007, 32, 483-492.	5.4	104
99	The amphetamine-induced sensitized state as a model of schizophrenia. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2007, 31, 1556-1571.	4.8	186
100	Dopamine transporter cell surface localization facilitated by a direct interaction with the dopamine D2 receptor. EMBO Journal, 2007, 26, 2127-2136.	7.8	182
101	Opposing effects of 5-HT2A and 5-HT2C receptor antagonists in the rat and mouse on premature responding in the five-choice serial reaction time test. Psychopharmacology, 2007, 195, 223-234.	3.1	185
102	Effects of dexfenfluramine and 5-HT3 receptor antagonists on stress-induced reinstatement of alcohol seeking in rats. Psychopharmacology, 2006, 186, 82-92.	3.1	47
103	The effects of the 5-HT2C receptor antagonist SB242084 on locomotor activity induced by selective, or mixed, indirect serotonergic and dopaminergic agonists. Psychopharmacology, 2006, 187, 515-525.	3.1	82
104	Effects of central neurokinin-1 receptor antagonism on cocaine- and opiate-induced locomotor activity and self-administration behaviour in rats. Pharmacology Biochemistry and Behavior, 2006, 84, 94-101.	2.9	21
105	Early life tactile stimulation changes adult rat responsiveness to amphetamine. Pharmacology Biochemistry and Behavior, 2006, 84, 497-503.	2.9	40
106	Dissociation between In Vivo Occupancy and Functional Antagonism of Dopamine D2 Receptors: Comparing Aripiprazole to Other Antipsychotics in Animal Models. Neuropsychopharmacology, 2006, 31, 1854-1863.	5.4	194
107	Sensitization to amphetamine, but not PCP, impairs attentional set shifting: reversal by a D1 receptor agonist injected into the medial prefrontal cortex. Psychopharmacology, 2005, 183, 190-200.	3.1	113
108	Sensitization to amphetamine, but not phencyclidine, disrupts prepulse inhibition and latent inhibition. Psychopharmacology, 2005, 180, 366-376.	3.1	74

#	Article	IF	CITATIONS
109	Activation of central neurokinin-1 receptors induces reinstatement of cocaine-seeking behavior. Neuroscience Letters, 2005, 390, 42-47.	2.1	20
110	Effects of injections of 8-hydroxy-2-(di-n-propylamino)tetralin or muscimol in the median raphe nucleus on c-fos mRNA in the rat brain. Neuroscience, 2005, 131, 475-479.	2.3	11
111	A putative animal model of the "prodromal―state of schizophrenia. Biological Psychiatry, 2005, 57, 586-593.	1.3	63
112	Injection of the 5-HT2C Receptor Agonist Ro60-0175 into the Ventral Tegmental Area Reduces Cocaine-Induced Locomotor Activity and Cocaine Self-Administration. Neuropsychopharmacology, 2004, 29, 308-318.	5.4	122
113	Evaluation of the motor initiation hypothesis of APD-induced conditioned avoidance decreases. Pharmacology Biochemistry and Behavior, 2004, 78, 811-819.	2.9	40
114	Fluoxetine, but not sertraline or citalopram, potentiates the locomotor stimulant effect of cocaine: possible pharmacokinetic effects. Psychopharmacology, 2004, 174, 406-13.	3.1	20
115	Infusion of the substance P analogue, DiMe-C7, into the ventral tegmental area induces reinstatement of cocaine-seeking behaviour in rats. Psychopharmacology, 2004, 177, 111-120.	3.1	38
116	5,7-Dihydroxytryptamine Lesions of the Dorsal and Median Raphe Nuclei Interfere With Lithium-Induced Conditioned Gaping, but Not Conditioned Taste Avoidance, in Rats Behavioral Neuroscience, 2004, 118, 1391-1399.	1.2	35
117	The 5-HT2A receptor antagonist M100,907 attenuates motor and 'impulsive-type' behaviours produced by NMDA receptor antagonism. Psychopharmacology, 2003, 170, 309-319.	3.1	162
118	Serotonin and drug reward: focus on 5-HT2C receptors. European Journal of Pharmacology, 2003, 480, 151-162.	3.5	147
119	Up-regulated dopamine D1 receptor binding can be detected in vivo following repeated SCH 23390, but not SKF 81297 or 6-hydroxydopamine, treatments. European Journal of Pharmacology, 2003, 459, 195-201.	3.5	6
120	Amphetamine-sensitized animals show a sensorimotor gating and neurochemical abnormality similar to that of schizophrenia. Schizophrenia Research, 2003, 64, 103-114.	2.0	86
121	Neonatal Ablation of the Nigrostriatal Dopamine Pathway Does Not Influence Limb Development in Rats. Experimental Neurology, 2002, 177, 547-556.	4.1	3
122	The Role of Corticotropin-Releasing Factor in the Median Raphe Nucleus in Relapse to Alcohol. Journal of Neuroscience, 2002, 22, 7844-7849.	3.6	127
123	Cyclooxygenase inhibitor modulation of dopamine-related behaviours. European Journal of Pharmacology, 2002, 450, 141-151.	3.5	29
124	Activation of 5-HT1B receptors in the nucleus accumbens reduces self-administration of amphetamine on a progressive ratio schedule. Pharmacology Biochemistry and Behavior, 2002, 71, 717-725.	2.9	49
125	Editorial. Pharmacology Biochemistry and Behavior, 2002, 71, 531-532.	2.9	0
126	Differential Effects of the 5-HT2A Receptor Antagonist M100,907 and the 5-HT2C Receptor Antagonist SB242,084 on Cocaine-induced Locomotor Activity, Cocaine Self-administration and Cocaine-induced Reinstatement of Responding. Neuropsychopharmacology, 2002, 27, 576-86.	5.4	210

#	Article	IF	CITATIONS
127	Multiple 5-HT receptors are involved in the effects of acute MDMA treatment: studies on locomotor activity and responding for conditioned reinforcement. Psychopharmacology, 2002, 162, 282-291.	3.1	76
128	The adenosine A2A agonist CGS 21680 reverses the reduction in prepulse inhibition of the acoustic startle response induced by phencyclidine, but not by apomorphine and amphetamine. Psychopharmacology, 2001, 156, 187-193.	3.1	20
129	Extracellular amino acid profiles in the paraventricular nucleus of the rat hypothalamus are influenced by diet composition. Brain Research, 2001, 892, 320-328.	2.2	34
130	Reduced Brain Serotonin Activity Disrupts Prepulse Inhibition of the Acoustic Startle Reflex Effects of 5,7-dihydroxytryptamine and p-chlorophenylalanine. Neuropsychopharmacology, 2001, 24, 399-409.	5.4	61
131	Pre-Exposure to (\hat{A}_{\pm}) 3,4-methylenedioxy-methamphetamine (MDMA) Facilitates Acquisition of Intravenous Cocaine Self-Administration in Rats. Neuropsychopharmacology, 2001, 25, 195-203.	5.4	41
132	Subchronic fluoxetine treatment induces a transient potentiation of amphetamine-induced hyperlocomotion: possible pharmacokinetic interaction. Behavioural Pharmacology, 2000, 11, 109-116.	1.7	15
133	Modification of dopamine D1 receptor knockout phenotype in mice lacking both dopamine D1 and D3 receptors. European Journal of Pharmacology, 2000, 399, 171-181.	3.5	48
134	Antipsychoticlike effects of amoxapine, without catalepsy, using the prepulse inhibition of the acoustic startle reflex test in rats. Biological Psychiatry, 2000, 47, 670-676.	1.3	11
135	The rewarding properties of neuropeptide Y in perifornical hypothalamus vs. nucleus accumbens. Peptides, 2000, 21, 1279-1287.	2.4	65
136	Dietary protein content affects the profiles of extracellular amino acids in the medial preoptic area of freely moving rats. Life Sciences, 2000, 66, 1105-1118.	4.3	24
137	Acute fluoxetine treatment potentiates amphetamine hyperactivity and amphetamine-induced nucleus accumbens dopamine release: possible pharmacokinetic interaction. Psychopharmacology, 1999, 141, 421-427.	3.1	31
138	Activation of 5-HT 1B receptors in the nucleus accumbens reduces amphetamine-induced enhancement of responding for conditioned reward. Psychopharmacology, 1999, 142, 165-174.	3.1	63
139	The potentiating effect of sertraline and fluoxetine on amphetamine-induced locomotor activity is not mediated by serotonin. Psychopharmacology, 1999, 143, 426-432.	3.1	26
140	Depletion of brain serotonin following intra-raphe injections. Psychopharmacology, 1999, 146, 185-193.	3.1	28
141	Selective destruction of brain serotonin neurons by 5,7-dihydroxytryptamine increases responding for a conditioned reward. Psychopharmacology, 1999, 147, 291-299.	3.1	75
142	The adenosine A1 receptor agonist N6-cyclopentyladenosine blocks the disruptive effect of phencyclidine on prepulse inhibition of the acoustic startle response in the rat. European Journal of Pharmacology, 1999, 369, 325-329.	3.5	30
143	Spatial learning deficit in dopamine D1 receptor knockout mice. European Journal of Pharmacology, 1999, 383, 95-106.	3.5	153
144	Measurement of Blood–Brain Barrier Permeability of Rats with Alpha-Aminoisobutyric Acid During Microdialysis. Physiology and Behavior, 1999, 67, 587-598.	2.1	12

#	Article	IF	CITATIONS
145	RU-24969 disrupts d-amphetamine self-administration and responding for conditioned reward via stimulation of 5-HT1B receptors. Behavioural Pharmacology, 1999, 10, 183-193.	1.7	30
146	Reversal of fenfluramine and fluoxetine anorexia by 8-OH-DPAT is attenuated following raphe injection of 5,7-dihydroxytryptamine. Brain Research, 1998, 800, 62-68.	2.2	21
147	Injections of d-amphetamine into the ventral pallidum increase locomotor activity and responding for conditioned reward: a comparison with injections into the nucleus accumbens. Brain Research, 1998, 805, 29-40.	2.2	35
148	Disruption of dopamine D1 receptor gene expression attenuates alcohol-seeking behavior. European Journal of Pharmacology, 1998, 353, 149-158.	3.5	154
149	Acute Amino Acid Loads That Deplete Brain Serotonin Fail to Alter Behavior. Pharmacology Biochemistry and Behavior, 1998, 59, 115-121.	2.9	26
150	A Comparison of the Effects of Risperidone, Raclopride, and Ritanserin on Intravenous Self-Administration of d-Amphetamine. Pharmacology Biochemistry and Behavior, 1998, 60, 55-60.	2.9	19
151	Reinstatement of alcohol-seeking by priming injections of alcohol and exposure to stress in rats. Psychopharmacology, 1998, 135, 169-174.	3.1	288
152	Neuropeptide Y-induced operant responding for sucrose is not mediated by dopamine. Peptides, 1998, 19, 1667-1673.	2.4	38
153	Fluoxetine attenuates morphine-induced locomotion and blocks morphine-sensitization. European Journal of Pharmacology, 1997, 337, 161-164.	3.5	24
154	Differential effects of ondansetron and α-flupenthixol on responding for conditioned reward. Psychopharmacology, 1997, 134, 64-72.	3.1	22
155	Injection of 5-HT into the nucleus accumbens reduces the effects ofd-amphetamine on responding for conditioned reward. Psychopharmacology, 1996, 126, 62-69.	3.1	39
156	Effects ofd-fenfluramine and metergoline on responding for conditioned reward and the response potentiating effect of nucleus accumbensd-amphetamine. Psychopharmacology, 1995, 118, 155-163.	3.1	46
157	Further studies to examine the nature of dexfenfluramine-induced suppression of heroin self-administration. Psychopharmacology, 1995, 120, 134-141.	3.1	26
158	Median raphe injections of 8-OH-DPAT lower frequency thresholds for lateral hypothalamic self-stimulation. Pharmacology Biochemistry and Behavior, 1995, 52, 65-71.	2.9	54
159	Effects of combined or separate 5,7-dihydroxytryptamine lesions of the dorsal and median raphe nuclei on responding maintained by a DRL 20s schedule of food reinforcement. Brain Research, 1995, 675, 45-54.	2.2	76
160	Median and dorsal raphe injections of the 5-HT1A agonist, 8-OH-DPAT, and the GABAA agonist, muscimol, increase voluntary ethanol intake in wistar rats. Neuropharmacology, 1994, 33, 349-358.	4.1	79
161	Injecting 5-HT into the PVN does not prevent feeding induced by injecting 8-OH-DPAT into the raphe. Pharmacology Biochemistry and Behavior, 1993, 46, 487-491.	2.9	39
162	Conditioned place preference induced by microinjection of 8-OH-DPAT into the dorsal or median raphe nucleus. Psychopharmacology, 1993, 113, 31-36.	3.1	64

#	Article	IF	CITATIONS
163	Radiofrequency lesions of the PVN fail to modify the effects of serotonergic drugs on food intake. Brain Research, 1993, 630, 1-9.	2.2	29
164	A comparison of the effects of dorsal or median raphe injections of 8-OH-DPAT in three operant tasks measuring response inhibition. Behavioural Brain Research, 1993, 54, 187-197.	2.2	59
165	A comparison of the effects of the 5-HT1 agonists TFMPP and RU 24969 on feeding following peripheral or medial hypothalamic injection. Brain Research, 1992, 580, 265-272.	2.2	27
166	Effect of drugs influencing 5-HT function on ethanol drinking and feeding behaviour in rats: Studies using a drinkometer system. Neuroscience and Biobehavioral Reviews, 1992, 16, 535-552.	6.1	88
167	The polyamine synthesis inhibitor α-difluoromethylornithine is neuroprotective against N-methyl-D-aspartate-induced brain damage in vivo. European Journal of Pharmacology, 1991, 209, 101-103.	3.5	24
168	Dopamine receptor blockade in nucleus accumbens or caudate nucleus differentially affects feeding induced by 8-OH-DPAT injected into dorsal or median raphe. Brain Research, 1991, 552, 181-189.	2.2	64
169	Opiate antagonists inhibit feeding induced by 8-OH-DPAT: possible mediation in the nucleus accumbens. Brain Research, 1991, 560, 260-267.	2.2	21
170	Influence of taste and food texture on the feeding responses induced by 8-OH-DPAT and gepirone. Psychopharmacology, 1991, 104, 302-306.	3.1	14
171	Effects of 8-OH-DPAT, buspirone and ICS 205-930 on feeding in a novel environment: comparisons with chlordiazepoxide and FG 7142. Psychopharmacology, 1990, 102, 301-308.	3.1	28
172	A pharmacological analysis of the eating response induced by 8-OH-DPAT injected into the dorsal raphé nucleus reveals the involvement of a dopaminergic mechanism. Psychopharmacology, 1990, 100, 188-194.	3.1	33
173	Dorsal raphe microinjection of 5-HT and indirect 5-HT agonists induces feeding in rats. European Journal of Pharmacology, 1990, 184, 265-271.	3.5	33
174	The involvement of 5â€hydroxytryptaminergic and dopaminergic mechanisms in the eating induced by buspirone, gepirone and ipsapirone. British Journal of Pharmacology, 1990, 99, 519-525.	5.4	36
175	A comparison of the effects of tryptamine and 5-hydroxytryptamine on feeding following injection into the paraventricular nucleus of the hypothalamus. Pharmacology Biochemistry and Behavior, 1989, 32, 907-911.	2.9	28
176	Enhancement of 5-HT-induced anorexia: a test of the reversibility of monoamine oxidase inhibitors. Psychopharmacology, 1989, 98, 265-268.	3.1	3
177	Tryptamine impairs the acquisition of a one-way active avoidance task. Pharmacology Biochemistry and Behavior, 1989, 32, 317-321.	2.9	7
178	<i>m</i> â€Octopamine injected into the paraventricular nucleus induces eating in rats: a comparison with noradrenalineâ€induced eating. British Journal of Pharmacology, 1989, 97, 483-489.	5.4	13
179	Increased food intake in satiated rats induced by the 5-HT antagonists methysergide, metergoline and ritanserin. Psychopharmacology, 1988, 96, 237-42.	3.1	90
180	The effects of d-fenfluramine on saccharin intake and preference, and on food and water intake. Pharmacology Biochemistry and Behavior, 1988, 29, 687-691.	2.9	12

#	Article	IF	CITATIONS
181	8-OH-DPAT elicits gnawing, and eating of solid but not liquid foods. Psychopharmacology, 1987, 92, 192-5.	3.1	35
182	The anorectic action of peripheral 5-HT examined in the runway: evidence for an action on satiation. Psychopharmacology, 1987, 93, 498-501.	3.1	6
183	Dissociation of the anorectic actions of 5-HTP and fenfluramine. Psychopharmacology, 1986, 89, 216-220.	3.1	35
184	Behavioural and pharmacological investigations of 5-HT hypophagia and hyperdipsia. Pharmacology Biochemistry and Behavior, 1986, 25, 23-28.	2.9	20
185	Microstructural analysis of the anorectic action of peripherally administered 5-HT. Pharmacology Biochemistry and Behavior, 1986, 24, 1133-1136.	2.9	32
186	The anorectic action of peripherally administered 5-HT is enhanced by vagotomy. Physiology and Behavior, 1985, 34, 861-866.	2.1	29
187	Effects of manipulations of peripheral serotonin on feeding and drinking in the rat. Pharmacology Biochemistry and Behavior, 1984, 20, 835-840.	2.9	72