

Clia Duarte Cruz

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1879513/celia-duarte-cruz-publications-by-year.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

61 papers	1,870 citations	26 h-index	42 g-index
85 ext. papers	2,086 ext. citations	3.6 avg, IF	4.72 L-index

#	Paper	IF	Citations
61	Rewired glycosylation activity promotes scarless regeneration and functional recovery in spiny mice after complete spinal cord transection.. <i>Developmental Cell</i> , 2021 ,	10.2	3
60	Biomarkers for Bladder Pain Syndrome/Interstitial Cystitis. <i>Current Bladder Dysfunction Reports</i> , 2021 , 16, 12-18	0.4	1
59	The urethra in continence and sensation: Neural aspects of urethral function. <i>Neurourology and Urodynamics</i> , 2021 , 40, 744-752	2.3	1
58	Chronic Pain After Spinal Cord Injury: Is There a Role for Neuron-Immune Dysregulation?. <i>Frontiers in Physiology</i> , 2020 , 11, 748	4.6	6
57	TASCI-transcutaneous tibial nerve stimulation in patients with acute spinal cord injury to prevent neurogenic detrusor overactivity: protocol for a nationwide, randomised, sham-controlled, double-blind clinical trial. <i>BMJ Open</i> , 2020 , 10, e039164	3	6
56	Effects of early intravesical administration of resiniferatoxin to spinal cord-injured rats in neurogenic detrusor overactivity. <i>Neurourology and Urodynamics</i> , 2019 , 38, 1540-1550	2.3	5
55	Peripherally administered melanocortins induce mice fat browning and prevent obesity. <i>International Journal of Obesity</i> , 2019 , 43, 1058-1069	5.5	5
54	Underactive bladder in aging rats is associated with a reduced number of serotonin-expressing cells in the urethra and is improved by serotonin application to the urethra. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2019 , 11, 248-254	1.9	6
53	Partners in Crime: NGF and BDNF in Visceral Dysfunction. <i>Current Neuropharmacology</i> , 2019 , 17, 1021-1038	3.8	16
52	Evidence for an urethro-vesical crosstalk mediated by serotonin. <i>Neurourology and Urodynamics</i> , 2018 , 37, 2389-2397	2.3	11
51	Author Reply. <i>Urology</i> , 2017 , 99, 55-56	1.6	
50	Urinary Neurotrophin Levels Increase in Women With Stress Urinary Incontinence After a Midurethral Sling Procedure. <i>Urology</i> , 2017 , 99, 49-56	1.6	5
49	Impairment of sensory afferents by intrathecal administration of botulinum toxin A improves neurogenic detrusor overactivity in chronic spinal cord injured rats. <i>Experimental Neurology</i> , 2016 , 285, 159-166	5.7	17
48	Can the adrenergic system be implicated in the pathophysiology of bladder pain syndrome/interstitial cystitis? A clinical and experimental study. <i>Neurourology and Urodynamics</i> , 2015 , 34, 489-96	2.3	21
47	Urinary bladder inflammation induces changes in urothelial nerve growth factor and TRPV1 channels. <i>British Journal of Pharmacology</i> , 2015 , 172, 1691-9	8.6	27
46	Biomarkers of spinal cord injury and ensuing bladder dysfunction. <i>Advanced Drug Delivery Reviews</i> , 2015 , 82-83, 153-9	18.5	26
45	The role of brain-derived neurotrophic factor (BDNF) in the development of neurogenic detrusor overactivity (NDO). <i>Journal of Neuroscience</i> , 2015 , 35, 2146-60	6.6	30

44	Co-administration of transient receptor potential vanilloid 4 (TRPV4) and TRPV1 antagonists potentiate the effect of each drug in a rat model of cystitis. <i>BJU International</i> , 2015 , 115, 452-60	5.6	16
43	Neurotrophins in bladder function: what do we know and where do we go from here?. <i>Neurourology and Urodynamics</i> , 2014 , 33, 39-45	2.3	48
42	Ulcerative and nonulcerative forms of bladder pain syndrome/interstitial cystitis do not differ in symptom intensity or response to onabotulinum toxin A. <i>Urology</i> , 2014 , 83, 1030-4	1.6	36
41	Intrathecal administration of botulinum toxin type A improves urinary bladder function and reduces pain in rats with cystitis. <i>European Journal of Pain</i> , 2014 , 18, 1480-9	3.7	29
40	Biomarkers in lower urinary tract symptoms/overactive bladder: a critical overview. <i>Current Opinion in Urology</i> , 2014 , 24, 352-7	2.8	19
39	MP76-06 URINARY NEUROTROPHINS AND QMAX VARIATION MAY PREDICT DE NOVO URGENCY IN SUI PATIENTS AFTER A MIDURETHRAL SLING (MUS) SURGERY. <i>Journal of Urology</i> , 2014 , 191,	2.5	1
38	Brain-derived neurotrophic factor, acting at the spinal cord level, participates in bladder hyperactivity and referred pain during chronic bladder inflammation. <i>Neuroscience</i> , 2013 , 234, 88-102	3.9	20
37	Transient receptor potential channels in bladder function. <i>Acta Physiologica</i> , 2013 , 207, 110-22	5.6	34
36	Urinary neurotrophic factors in healthy individuals and patients with overactive bladder. <i>Journal of Urology</i> , 2013 , 189, 359-65	2.5	55
35	Spread of onabotulinumtoxinA after bladder injection. Experimental study using the distribution of cleaved SNAP-25 as the marker of the toxin action. <i>European Urology</i> , 2012 , 61, 1178-84	10.2	57
34	Transient receptor potential vanilloid 1 mediates nerve growth factor-induced bladder hyperactivity and noxious input. <i>BJU International</i> , 2012 , 110, E422-8	5.6	20
33	Animal Models of Cystitis. <i>Methods in Pharmacology and Toxicology</i> , 2012 , 397-409	1.1	
32	Effect of onabotulinumtoxinA on intramural parasympathetic ganglia: an experimental study in the guinea pig bladder. <i>Journal of Urology</i> , 2012 , 187, 1121-6	2.5	24
31	817 URINARY NEUROTROPHIC FACTORS IN BLADDER PAIN SYNDROME/INTERSTITIAL CYSTITIS. <i>Journal of Urology</i> , 2012 , 187,	2.5	1
30	Rat detrusor overactivity induced by chronic spinalization can be abolished by a transient receptor potential vanilloid 1 (TRPV1) antagonist. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2012 , 166, 35-8	2.4	26
29	Neurotrophins as regulators of urinary bladder function. <i>Nature Reviews Urology</i> , 2012 , 9, 628-37	5.5	64
28	365 TRPV1 and TRPV4 antagonists have synergistic effect for treating bladder overactivity in rats. <i>European Urology Supplements</i> , 2012 , 11, e365	0.9	4
27	Severe burn injury induces a characteristic activation of extracellular signal-regulated kinase 1/2 in spinal dorsal horn neurons. <i>European Journal of Pain</i> , 2011 , 15, 683-90	3.7	6

26	Spinal cord injury and bladder dysfunction: new ideas about an old problem. <i>Scientific World Journal, The</i> , 2011 , 11, 214-34	2.2	48
25	Minocycline completely reverses mechanical hyperalgesia in diabetic rats through microglia-induced changes in the expression of the potassium chloride co-transporter 2 (KCC2) at the spinal cord. <i>Diabetes, Obesity and Metabolism</i> , 2011 , 13, 150-9	6.7	61
24	Nerve growth factor in bladder dysfunction: contributing factor, biomarker, and therapeutic target. <i>Neurourology and Urodynamics</i> , 2011 , 30, 1227-41	2.3	96
23	Neurotrophins in the lower urinary tract: becoming of age. <i>Current Neuropharmacology</i> , 2011 , 9, 553-8	7.6	12
22	Biomarkers in overactive bladder: a new objective and noninvasive tool?. <i>Advances in Urology</i> , 2011 , 2011, 382431	1.6	45
21	191 EFFECT OF BOTULINUM TOXIN TYPE A ON INTRAMURAL PARASYMPATHETIC GANGLIA OF THE GUINEA-PIG BLADDER. <i>Journal of Urology</i> , 2010 , 183,	2.5	1
20	Sequestration of brain derived nerve factor by intravenous delivery of TrkB-Ig2 reduces bladder overactivity and noxious input in animals with chronic cystitis. <i>Neuroscience</i> , 2010 , 166, 907-16	3.9	36
19	Distribution of the high-affinity binding site and intracellular target of botulinum toxin type A in the human bladder. <i>European Urology</i> , 2010 , 57, 884-90	10.2	70
18	Trigonal injection of botulinum toxin A in patients with refractory bladder pain syndrome/interstitial cystitis. <i>European Urology</i> , 2010 , 58, 360-5	10.2	141
17	258 DISTRIBUTION AND NEUROCHEMISTRY OF BOTULINUM TOXIN TYPE A RECEPTORS IN THE URINARY BLADDER. <i>European Journal of Pain</i> , 2009 , 13, S81b	3.7	
16	652 CHEMICAL NEUROMODULATION IN PATIENTS WITH BLADDER PAIN SYNDROME (BPS). <i>European Journal of Pain</i> , 2009 , 13, S189	3.7	
15	653 INTRATHECAL BLOCKADE OF NGF DECREASES REFERRED PAIN IN RATS MODEL OF CHRONIC BLADDER INFLAMMATION. <i>European Journal of Pain</i> , 2009 , 13, S189a	3.7	
14	654 INTRATHECAL BDNF SEQUESTRATION REDUCES REFERRED PAIN AND BLADDER OVERACTIVITY IN AN ANIMAL MODEL OF CHRONIC BLADDER INFLAMMATION. <i>European Journal of Pain</i> , 2009 , 13, S189b	3.7	
13	GRC-6211, a new oral specific TRPV1 antagonist, decreases bladder overactivity and noxious bladder input in cystitis animal models. <i>Journal of Urology</i> , 2009 , 181, 379-86	2.5	78
12	223 DISTRIBUTION AND NEUROCHEMISTRY OF HIGH AFFINITY BINDING SITES FOR BOTULINUM TOXIN TYPE A IN THE URINARY BLADDER. <i>European Urology Supplements</i> , 2009 , 8, 176	0.9	4
11	Intrathecal delivery of resiniferatoxin (RTX) reduces detrusor overactivity and spinal expression of TRPV1 in spinal cord injured animals. <i>Experimental Neurology</i> , 2008 , 214, 301-8	5.7	26
10	Transient receptor potential vanilloid subfamily 1 is essential for the generation of noxious bladder input and bladder overactivity in cystitis. <i>Journal of Urology</i> , 2007 , 177, 1537-41	2.5	95
9	The activation of the ERK pathway contributes to the spinal c-fos expression observed after noxious bladder stimulation. <i>Somatosensory & Motor Research</i> , 2007 , 24, 15-20	1.2	26

8	The ERK 1 and 2 pathway in the nervous system: from basic aspects to possible clinical applications in pain and visceral dysfunction. <i>Current Neuropharmacology</i> , 2007 , 5, 244-52	7.6	61
7	Increased extracellular signal regulated kinases phosphorylation in the adrenal gland in response to chronic ACTH treatment. <i>Journal of Endocrinology</i> , 2007 , 192, 647-58	4.7	22
6	Spinal ERK activation contributes to the regulation of bladder function in spinal cord injured rats. <i>Experimental Neurology</i> , 2006 , 200, 66-73	5.7	22
5	Inhibition of ERK phosphorylation decreases nociceptive behaviour in monoarthritic rats. <i>Pain</i> , 2005 , 116, 411-419	8	69
4	Increased spinal cord phosphorylation of extracellular signal-regulated kinases mediates micturition overactivity in rats with chronic bladder inflammation. <i>European Journal of Neuroscience</i> , 2005 , 21, 773-81	3.5	53
3	ACTH modulates ERK phosphorylation in the adrenal gland in a time-dependent manner. <i>Endocrine Research</i> , 2004 , 30, 661-6	1.9	12
2	Nerve growth factor regulates galanin and c-jun overexpression occurring in dorsal root ganglion cells after intravesical resiniferatoxin application. <i>Brain Research</i> , 2002 , 951, 264-9	3.7	22
1	Vanilloid receptor 1 expression in the rat urinary tract. <i>Neuroscience</i> , 2002 , 109, 787-98	3.9	201