

MÃ©lanie Tremblay

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/189105/publications.pdf>

Version: 2024-02-01

18
papers

506
citations

840776

11
h-index

888059

17
g-index

19
all docs

19
docs citations

19
times ranked

659
citing authors

#	ARTICLE	IF	CITATIONS
1	Elevated Serum Liver-Type Fatty Acid Binding Protein Levels in Non-acetaminophen Acute Liver Failure Patients with Organ Dysfunction. <i>Digestive Diseases and Sciences</i> , 2021, 66, 273-283.	2.3	8
2	Bile duct ligation renders the brain susceptible to hypotension-induced neuronal degeneration: Implications of ammonia. <i>Journal of Neurochemistry</i> , 2021, 157, 561-573.	3.9	10
3	Genetically engineered <i>E. coli</i> Nissle attenuates hyperammonemia and prevents memory impairment in bile duct ligated rats. <i>Liver International</i> , 2021, 41, 1020-1032.	3.9	10
4	An Investigation of PS- <i>b</i> -PEO Polymersomes for the Oral Treatment and Diagnosis of Hyperammonemia. <i>Small</i> , 2019, 15, e1902347.	10.0	22
5	Progressive resistance training prevents loss of muscle mass and strength in bile duct ligated rats. <i>Liver International</i> , 2019, 39, 676-683.	3.9	10
6	REPLY:. <i>Hepatology</i> , 2017, 66, 670-671.	7.3	1
7	The bile duct ligated rat: A relevant model to study muscle mass loss in cirrhosis. <i>Metabolic Brain Disease</i> , 2017, 32, 513-518.	2.9	30
8	Elevated FABP1 serum levels are associated with poorer survival in acetaminophen-induced acute liver failure. <i>Hepatology</i> , 2017, 65, 938-949.	7.3	49
9	The association between FABP7 serum levels with survival and neurological complications in acetaminophen-induced acute liver failure: a nested case-control study. <i>Annals of Intensive Care</i> , 2017, 7, 99.	4.6	4
10	Liposome-Supported Peritoneal Dialysis for the Treatment of Hyperammonemia-Associated Encephalopathy. <i>Advanced Functional Materials</i> , 2016, 26, 8382-8389.	14.9	24
11	Induction of systemic oxidative stress leads to brain oedema in portacaval shunted rats. <i>Liver International</i> , 2014, 34, 1322-1329.	3.9	16
12	Increased brain lactate is central to the development of brain edema in rats with chronic liver disease. <i>Journal of Hepatology</i> , 2014, 60, 554-560.	3.7	65
13	Systemic oxidative stress is implicated in the pathogenesis of brain edema in rats with chronic liver failure. <i>Free Radical Biology and Medicine</i> , 2012, 52, 1228-1235.	2.9	78
14	AST-120 (spherical carbon adsorbent) lowers ammonia levels and attenuates brain edema in bile duct ligated rats. <i>Hepatology</i> , 2011, 53, 1995-2002.	7.3	74
15	Portacaval anastomosis-induced hyperammonemia does not lead to oxidative stress. <i>Metabolic Brain Disease</i> , 2010, 25, 11-15.	2.9	11
16	Direct binding of anti-DNA topoisomerase I autoantibodies to the cell surface of fibroblasts in patients with systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2004, 50, 3265-3274.	6.7	76
17	Mechanism of Kex2p inhibition by its proregion. <i>FEBS Letters</i> , 2001, 508, 332-336.	2.8	17
18	Sex is associated with differences in oxidative stress and susceptibility to severe hepatic encephalopathy in bile duct ligated rats. <i>Journal of Neurochemistry</i> , 0, , .	3.9	0