

Jan Suda

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1899702/publications.pdf>

Version: 2024-02-01

97
papers

7,351
citations

71102

41
h-index

64796

79
g-index

117
all docs

117
docs citations

117
times ranked

6472
citing authors

#	ARTICLE	IF	CITATIONS
1	Morphological and environmental differentiation as prezygotic reproductive barriers between parapatric and allopatric <i>Campanula rotundifolia</i> agg. cytotypes. <i>Annals of Botany</i> , 2023, 131, 71-86.	2.9	2
2	Diverse fungal communities associated with the roots of isoetid plants are structured by host plant identity. <i>Fungal Ecology</i> , 2020, 45, 100914.	1.6	10
3	Associations between genomic ancestry, genome size and capitula morphology in the invasive meadow knapweed hybrid complex (<i>Centaurea A-moncktonii</i>) in eastern North America. <i>AoB PLANTS</i> , 2019, 11, plz055.	2.3	5
4	Diversity in genome size and GC content shows adaptive potential in orchids and is closely linked to partial endoreplication, plant life history traits and climatic conditions. <i>New Phytologist</i> , 2019, 224, 1642-1656.	7.3	63
5	Climatic conditions and human activities shape diploid-tetraploid coexistence at different spatial scales in the common weed <i>Tripleurospermum inodorum</i> (Asteraceae). <i>Journal of Biogeography</i> , 2019, 46, 1355-1366.	3.0	23
6	Nuclear Genome Size in Contrast to Sex Chromosome Number Variability in the Human Bed Bug, <i>Cimex lectularius</i> (Heteroptera: Cimicidae). <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2019, 95, 746-756.	1.5	8
7	Ploidy-altered phenotype interacts with local environment and may enhance polyploid establishment in <i>Knautia serpentinicola</i> (Caprifoliaceae). <i>New Phytologist</i> , 2019, 221, 1117-1127.	7.3	24
8	Small genome separates native and invasive populations in an ecologically important cosmopolitan grass. <i>Ecology</i> , 2018, 99, 79-90.	3.2	54
9	Sympatric diploid and tetraploid cytotypes of <i>Centaurea stoebe</i> s.l. do not differ in arbuscular mycorrhizal communities and mycorrhizal growth response. <i>American Journal of Botany</i> , 2018, 105, 1995-2007.	1.7	9
10	Does higher ploidy level increase the risk of invasion? A case study with two geo-cytotypes of <i>Solidago gigantea</i> Aiton (Asteraceae). <i>Journal of Plant Ecology</i> , 2018, 11, 317-327.	2.3	25
11	Polyploid species rely on vegetative reproduction more than diploids: a re-examination of the old hypothesis. <i>Annals of Botany</i> , 2017, 120, 341-349.	2.9	67
12	Evolutionary dynamics of mixed-ploidy populations in an annual herb: dispersal, local persistence and recurrent origins of polyploids. <i>Annals of Botany</i> , 2017, 120, 303-315.	2.9	59
13	Patterns, causes and consequences of genome size variation in Restionaceae of the Cape flora. <i>Botanical Journal of the Linnean Society</i> , 2017, 183, 515-531.	1.6	5
14	Mixed-Ploidy Species: Progress and Opportunities in Polyploid Research. <i>Trends in Plant Science</i> , 2017, 22, 1041-1055.	8.8	165
15	A New Species of <i>Cleisostoma</i> (Orchidaceae) from the Hon Ba Nature Reserve in Vietnam: A Multidisciplinary Assessment. <i>PLoS ONE</i> , 2016, 11, e0150631.	2.5	8
16	Phylogenetic marker development for target enrichment from transcriptome and genome skim data: the pipeline and its application in southern African <i>Oxalis</i> (Oxalidaceae). <i>Molecular Ecology Resources</i> , 2016, 16, 1124-1135.	4.8	101
17	Do ploidy level and nuclear genome size and latitude of origin modify the expression of <i>Phragmites australis</i> traits and interactions with herbivores?. <i>Biological Invasions</i> , 2016, 18, 2531-2549.	2.4	44
18	The Enigma of Progressively Partial Endoreplication: New Insights Provided by Flow Cytometry and Next-Generation Sequencing. <i>Genome Biology and Evolution</i> , 2016, 8, 1996-2005.	2.5	19

#	ARTICLE	IF	CITATIONS
19	Speciesâ€rich and polyploidâ€poor: Insights into the evolutionary role of wholeâ€genome duplication from the Cape flora biodiversity hotspot. <i>American Journal of Botany</i> , 2016, 103, 1336-1347.	1.7	28
20	Challenges of flowâ€cytometric estimation of nuclear genome size in orchids, a plant group with both wholeâ€genome and progressively partial endoreplication. <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2015, 87, 958-966.	1.5	51
21	Evolutionary and Taxonomic Implications of Variation in Nuclear Genome Size: Lesson from the Grass Genus <i>Anthoxanthum</i> (Poaceae). <i>PLoS ONE</i> , 2015, 10, e0133748.	2.5	24
22	Ecological differentiation, lack of hybrids involving diploids, and asymmetric gene flow between polyploids in narrow contact zones of <i>Senecio carniolicus</i> (syn. <i>Jacobaea carniolica</i>), <i>Tj ETQq0 0 0 rgBTgOverlook 10 Tf 50</i>		
23	Ecological differentiation of diploid and polyploid cytotypes of <i>Senecio carniolicus sensu lato</i> (Asteraceae) is stronger in areas of sympatry. <i>Annals of Botany</i> , 2015, 117, mcv176.	2.9	26
24	Evaluating the relationship between diploid and tetraploid <i>Vaccinium oxycoccos</i> (Ericaceae) in eastern Canada. <i>Botany</i> , 2015, 93, 623-636.	1.0	16
25	The origin of unique diversity in deglaciated areas: traces of <i>Pleistocene</i> processes in northâ€European endemics from the <i>Galium pusillum</i> polyploid complex (<i>Rubiaceae</i>). <i>Molecular Ecology</i> , 2015, 24, 1311-1334.	3.9	13
26	Niche partitioning in arbuscular mycorrhizal communities in temperate grasslands: a lesson from adjacent serpentine and nonserpentine habitats. <i>Molecular Ecology</i> , 2015, 24, 1831-1843.	3.9	31
27	The hidden side of plant invasions: the role of genome size. <i>New Phytologist</i> , 2015, 205, 994-1007.	7.3	99
28	Naturalization of central European plants in North America: species traits, habitats, propagule pressure, residence time. <i>Ecology</i> , 2015, 96, 762-774.	3.2	166
29	Genome size and phenotypic variation of <i>Nymphaea</i> (Nymphaeaceae) species from Eastern Europe and temperate Asia. <i>Acta Societatis Botanicorum Poloniae</i> , 2015, 84, 277-286.	0.8	3
30	Phylogenetic signal in growth and reproductive traits and in their plasticity: the <i>Descurainia</i> radiation in the Canary Islands. <i>Botanical Journal of the Linnean Society</i> , 2014, 174, 384-398.	1.6	6
31	Morphological <i>versus</i> genetic diversity of <i>Viola reichenbachiana</i> and <i>V. Arviniana</i> (sect. <i>Viola</i> , <i>Violaceae</i>) from soils differing in heavy metal content. <i>Plant Biology</i> , 2014, 16, 924-934.	3.8	20
32	Intraspecific ploidy variation: A hidden, minor player in plantâ€soilâ€mycorrhizal fungi interactions. <i>American Journal of Botany</i> , 2014, 101, 26-33.	1.7	17
33	Nonadaptive processes governing early stages of polyploid evolution: Insights from a primary contact zone of relict serpentine <i>Knautia arvensis</i> (Caprifoliaceae). <i>American Journal of Botany</i> , 2014, 101, 935-945.	1.7	32
34	Ploidyâ€specific symbiotic interactions: divergence of mycorrhizal fungi between cytotypes of the <i>Gymnadenia conopsea</i> group (<i>Orchidaceae</i>). <i>New Phytologist</i> , 2013, 199, 1022-1033.	7.3	92
35	Diversity and endemism in deglaciated areas: ploidy, relative genome size and niche differentiation in the <i>Galium pusillum</i> complex (<i>Rubiaceae</i>) in Northern and Central Europe. <i>Annals of Botany</i> , 2013, 111, 1095-1108.	2.9	30
36	The spatio-ecological segregation of different cytotypes of <i>Oxalis obtusa</i> (<i>Oxalidaceae</i>) in contact zones. <i>South African Journal of Botany</i> , 2013, 88, 62-68.	2.5	8

#	ARTICLE	IF	CITATIONS
37	The Incidence of Polyploidy in Natural Plant Populations: Major Patterns and Evolutionary Processes. , 2013, , 255-276.		191
38	High ploidy diversity and distinct patterns of cytotype distribution in a widespread species of <i>Oxalis</i> in the Greater Cape Floristic Region. <i>Annals of Botany</i> , 2013, 111, 641-649.	2.9	51
39	Genome size variation in Orchidaceae subfamily Apostasioideae: filling the phylogenetic gap. <i>Botanical Journal of the Linnean Society</i> , 2013, 172, 95-105.	1.6	27
40	Cytogeography of <i>Oxalis pes-caprae</i> in its native range: where are the pentaploids?. <i>Biological Invasions</i> , 2013, 15, 1189-1194.	2.4	14
41	Ploidy frequencies in plants with ploidy heterogeneity: fitting a general gametic model to empirical population data. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2013, 280, 20122387.	2.6	30
42	Hitting the right target: taxonomic challenges for, and of, plant invasions. <i>AoB PLANTS</i> , 2013, 5, plt042-plt042.	2.3	87
43	Two new species of <i>Oxalis</i> (Oxalidaceae) from the Greater Cape Floristic Region. <i>Phytotaxa</i> , 2013, 124, 13.	0.3	6
44	Parental Ploidy Strongly Affects Offspring Fitness in Heteroploid Crosses among Three Cytotypes of Autopolyploid <i>Jacobaea carniolica</i> (Asteraceae). <i>PLoS ONE</i> , 2013, 8, e78959.	2.5	42
45	Ecological effects of cell-level processes: genome size, functional traits and regional abundance of herbaceous plant species. <i>Annals of Botany</i> , 2012, 110, 1357-1367.	2.9	37
46	The more the better? The role of polyploidy in facilitating plant invasions. <i>Annals of Botany</i> , 2012, 109, 19-45.	2.9	707
47	Minority cytotypes in European populations of the <i>Gymnadenia conopsea</i> complex (Orchidaceae) greatly increase intraspecific and intrapopulation diversity. <i>Annals of Botany</i> , 2012, 110, 977-986.	2.9	39
48	Bringing Together Evolution on Serpentine and Polyploidy: Spatiotemporal History of the Diploid-Tetraploid Complex of <i>Knautia arvensis</i> (Dipsacaceae). <i>PLoS ONE</i> , 2012, 7, e39988.	2.5	52
49	Glycerol-treated nuclear suspensionsâ€”an efficient preservation method for flow cytometric analysis of plant samples. <i>Chromosome Research</i> , 2012, 20, 303-315.	2.2	21
50	Surprising spectra of root-associated fungi in submerged aquatic plants. <i>FEMS Microbiology Ecology</i> , 2012, 80, 216-235.	2.7	119
51	The symbiosis with arbuscular mycorrhizal fungi contributes to plant tolerance to serpentine edaphic stress. <i>Soil Biology and Biochemistry</i> , 2012, 44, 56-64.	8.8	64
52	Extensive range persistence in peripheral and interior refugia characterizes Pleistocene range dynamics in a widespread Alpine plant species (<i>Senecio carniolicus</i> , Asteraceae). <i>Molecular Ecology</i> , 2012, 21, 1255-1270.	3.9	44
53	Invasiveness in introduced Australian acacias: the role of species traits and genome size. <i>Diversity and Distributions</i> , 2011, 17, 884-897.	4.1	64
54	Arbuscular mycorrhizal symbiosis on serpentine soils: the effect of native fungal communities on different <i>Knautia arvensis</i> ecotypes. <i>Plant and Soil</i> , 2011, 345, 325-338.	3.7	21

#	ARTICLE	IF	CITATIONS
55	Remarkable coexistence of multiple cytotypes of the <i>Gymnadenia conopsea</i> aggregate (the fragrant) Tj ETQq1 1 0.784314 rgBT /Overlo	2.9	87
56	Apomixis is not prevalent in subnival to nival plants of the European Alps. <i>Annals of Botany</i> , 2011, 108, 381-390.	2.9	32
57	Molecular systematics and ecology of invasive Kangaroo Paws in South Africa: management implications for a horticulturally important genus. <i>Biological Invasions</i> , 2010, 12, 3989-4002.	2.4	22
58	Identification of candidates for cyclotide biosynthesis and cyclisation by expressed sequence tag analysis of <i>Oldenlandia affinis</i> . <i>BMC Genomics</i> , 2010, 11, 111.	2.8	30
59	The quest for suitable reference standards in genome size research. <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2010, 77A, 717-720.	1.5	40
60	Ploidy-specific interactions of three host plants with arbuscular mycorrhizal fungi: Does genome copy number matter?. <i>American Journal of Botany</i> , 2010, 97, 1798-1807.	1.7	20
61	Species boundaries and frequency of hybridization in the <i>Dryopteris carthusiana</i> (<i>Dryopteridaceae</i>) complex: A taxonomic puzzle resolved using genome size data. <i>American Journal of Botany</i> , 2010, 97, 1208-1219.	1.7	31
62	Distribution and habitat segregation on different spatial scales among diploid, tetraploid and hexaploid cytotypes of <i>Senecio carniolicus</i> (<i>Asteraceae</i>) in the Eastern Alps. <i>Annals of Botany</i> , 2010, 106, 967-977.	2.9	109
63	Genome Size. <i>Journal of Botany</i> , 2010, 2010, 1-4.	1.2	14
64	Karyological features of wild and cultivated forms of myrtle (<i>Myrtus communis</i> , <i>Myrtaceae</i>). <i>Genetics and Molecular Research</i> , 2010, 9, 429-433.	0.2	3
65	Genome size diversity in orchids: consequences and evolution. <i>Annals of Botany</i> , 2009, 104, 469-481.	2.9	156
66	Complex pattern of genome size variation in a polymorphic member of the <i>Asteraceae</i> . <i>Journal of Biogeography</i> , 2009, 36, 372-384.	3.0	39
67	Reciprocal Pleistocene origin and postglacial range formation of an allopolyploid and its sympatric ancestors (<i>Androsace adfinis</i> group, <i>Primulaceae</i>). <i>Molecular Phylogenetics and Evolution</i> , 2009, 50, 74-83.	2.7	45
68	Towards resolving the <i>Knautia arvensis</i> agg. (<i>Dipsacaceae</i>) puzzle: primary and secondary contact zones and ploidy segregation at landscape and microgeographic scales. <i>Annals of Botany</i> , 2009, 103, 963-974.	2.9	125
69	Ecological segregation drives fine-scale cytotype distribution of <i>Senecio carniolicus</i> in the Eastern Alps. <i>Preslia</i> , 2009, 81, 309-319.	2.8	39
70	ORIGINAL ARTICLE: Colonization and diversification in the African "sky islands"™ by Eurasian <i>Lychnis</i> L. (<i>Caryophyllaceae</i>). <i>Journal of Biogeography</i> , 2008, 35, 1016-1029.	3.0	55
71	Plant flow cytometry "Far beyond the stone age. <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2008, 73A, 579-580.	1.5	10
72	Complex distribution patterns of di-, tetra-, and hexaploid cytotypes in the European high mountain plant <i>Senecio carniolicus</i> (<i>Asteraceae</i>). <i>American Journal of Botany</i> , 2007, 94, 1391-1401.	1.7	111

#	ARTICLE	IF	CITATIONS
73	Chromosome Numbers and Genome Size Variation in Indian Species of <i>Curcuma</i> (Zingiberaceae). <i>Annals of Botany</i> , 2007, 100, 505-526.	2.9	135
74	Genome Size Variation and Species Relationships in <i>Hieracium</i> Sub-genus <i>Pilosella</i> (Asteraceae) as Inferred by Flow Cytometry. <i>Annals of Botany</i> , 2007, 100, 1323-1335.	2.9	98
75	Applications of Flow Cytometry to Evolutionary and Population Biology. <i>Annual Review of Ecology, Evolution, and Systematics</i> , 2007, 38, 847-876.	8.3	164
76	Circumpolar phylogeography of <i>Juncus biglumis</i> (Juncaceae) inferred from AFLP fingerprints, cpDNA sequences, nuclear DNA content and chromosome numbers. <i>Molecular Phylogenetics and Evolution</i> , 2007, 42, 92-103.	2.7	174
77	Estimation of nuclear DNA content in plants using flow cytometry. <i>Nature Protocols</i> , 2007, 2, 2233-2244.	12.0	1,219
78	Nuclear vs. plastid data: complex Pleistocene history of a circumpolar key species. <i>Molecular Ecology</i> , 2007, 16, 3902-3925.	3.9	243
79	DNA ploidy-level variation in native and invasive populations of <i>Lythrum salicaria</i> at a large geographical scale. <i>Journal of Biogeography</i> , 2007, 35, 070901070439003-???	3.0	14
80	Sympatric diploid and hexaploid cytotypes of <i>Senecio carniolicus</i> (Asteraceae) in the Eastern Alps are separated along an altitudinal gradient. <i>Journal of Plant Research</i> , 2007, 120, 721-725.	2.4	69
81	Estimation of Relative Nuclear DNA Content in Dehydrated Plant Tissues by Flow Cytometry. <i>Current Protocols in Cytometry</i> , 2006, 38, Unit7.30.	3.7	47
82	Ploidy level versus DNA ploidy level: an appeal for consistent terminology. <i>Taxon</i> , 2006, 55, 447-450.	0.7	166
83	Population dynamics and clonal growth of <i>Spartocytisus supranubius</i> (Fabaceae), a dominant shrub in the alpine zone of Tenerife, Canary Islands. <i>Plant Ecology</i> , 2006, 186, 97-108.	1.6	22
84	A modified method of flow cytometric seed screen simplifies the quantification of progeny classes with different ploidy levels. <i>Biologia Plantarum</i> , 2006, 50, 457-460.	1.9	22
85	Reliable DNA ploidy determination in dehydrated tissues of vascular plants by DAPI flow cytometry – new prospects for plant research. <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2006, 69A, 273-280.	1.5	135
86	The ghost of hybridization past: niche pre-emption is not the only explanation of apparent monophyly in island endemics. <i>Journal of Ecology</i> , 2005, 93, 572-575.	4.0	40
87	Genome size variation in Macaronesian angiosperms: forty percent of the Canarian endemic flora completed. <i>Plant Systematics and Evolution</i> , 2005, 252, 215-238.	0.9	67
88	Genome size discriminates between closely related taxa <i>Elytrigia repens</i> and <i>E. intermedia</i> (Poaceae). <i>Trends in Ecology and Evolution</i> , 2005, 20, 45-46.	0.9	45
89	Variation in <i>Lamium</i> subg. <i>Galeobdolon</i> (Lamiaceae) ? insights from ploidy levels, morphology and isozymes. <i>Plant Systematics and Evolution</i> , 2004, 244, 219-244.	0.9	17
90	Cytotype distribution in <i>Empetrum</i> (Ericaceae) at various spatial scales in the Czech Republic. <i>Folia Geobotanica</i> , 2004, 39, 161-171.	0.9	17

#	ARTICLE	IF	CITATIONS
91	Genome Size Variation in Central European Species of <i>Cirsium</i> (Compositae) and their Natural Hybrids. <i>Annals of Botany</i> , 2004, 94, 353-363.	2.9	92
92	(1582) Proposal to conserve the name <i>Inula verbascifolia</i> (Willd.) Hausskn. against <i>I. verbascifolia</i> Poir. (Asteraceae) and with a conserved type. <i>Taxon</i> , 2003, 52, 358-359.	0.7	2
93	Genetic variation within the endangered species <i>Aldrovanda vesiculosa</i> (Droseraceae) as revealed by RAPD analysis. <i>Aquatic Botany</i> , 2003, 75, 159-172.	1.6	23
94	Variation in DNA ploidy Levels of <i>Reynoutria</i> Taxa in the Czech Republic. <i>Annals of Botany</i> , 2003, 92, 265-272.	2.9	63
95	Nuclear DNA Amounts in Macaronesian Angiosperms. <i>Annals of Botany</i> , 2003, 92, 153-164.	2.9	78
96	Sympatric occurrences of various cytotypes of <i>Vaccinium</i> sect. <i>Oxycoccus</i> (Ericaceae). <i>Nordic Journal of Botany</i> , 2002, 22, 593-601.	0.5	17
97	A taxonomic study of the <i>Vaccinium</i> sect. <i>Oxycoccus</i> (Hill) W.D.J. Kock (Ericaceae) in the Czech Republic and adjacent territories. <i>Folia Geobotanica</i> , 2001, 36, 303-320.	0.9	30