

# Penelope Hopwood

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1905912/publications.pdf>

Version: 2024-02-01

11  
papers

2,841  
citations

933264

10  
h-index

1281743

11  
g-index

11  
all docs

11  
docs citations

11  
times ranked

2721  
citing authors

#	ARTICLE	IF	CITATIONS
1	The UK Standardisation of Breast Radiotherapy (START) trials of radiotherapy hypofractionation for treatment of early breast cancer: 10-year follow-up results of two randomised controlled trials. <i>Lancet Oncology, The, 2013, 14, 1086-1094.</i>	5.1	1,141
2	Hypofractionated breast radiotherapy for 1 week versus 3 weeks (FAST-Forward): 5-year efficacy and late normal tissue effects results from a multicentre, non-inferiority, randomised, phase 3 trial. <i>Lancet, The, 2020, 395, 1613-1626.</i>	6.3	603
3	Partial-breast radiotherapy after breast conservation surgery for patients with early breast cancer (UK IMPORT LOW trial): 5-year results from a multicentre, randomised, controlled, phase 3, non-inferiority trial. <i>Lancet, The, 2017, 390, 1048-1060.</i>	6.3	448
4	Comparison of patient-reported breast, arm, and shoulder symptoms and body image after radiotherapy for early breast cancer: 5-year follow-up in the randomised Standardisation of Breast Radiotherapy (START) trials. <i>Lancet Oncology, The, 2010, 11, 231-240.</i>	5.1	201
5	The impact of age and clinical factors on quality of life in early breast cancer: An analysis of 2208 women recruited to the UK START Trial (Standardisation of Breast Radiotherapy Trial). <i>Breast, 2007, 16, 241-251.</i>	0.9	172
6	The assessment of body image in cancer patients. <i>European Journal of Cancer, 1993, 29, 276-281.</i>	1.3	94
7	The course of anxiety and depression over 5 years of follow-up and risk factors in women with early breast cancer: Results from the UK Standardisation of Radiotherapy Trials (START). <i>Breast, 2010, 19, 84-91.</i>	0.9	86
8	Patient-Reported Outcomes Over 5 Years After Whole- or Partial-Breast Radiotherapy: Longitudinal Analysis of the IMPORT LOW (CRUK/06/003) Phase III Randomized Controlled Trial. <i>Journal of Clinical Oncology, 2019, 37, 305-317.</i>	0.8	58
9	Can patient-reported outcomes be used instead of clinician-reported outcomes and photographs as primary endpoints of late normal tissue effects in breast radiotherapy trials? Results from the IMPORT LOW trial. <i>Radiotherapy and Oncology, 2019, 134, 220-230.</i>	0.3	17
10	New challenges in psycho-oncology: An embodied approach to body image. <i>Psycho-Oncology, 2019, 28, 211-218.</i>	1.0	16
11	Quality of life after breast radiotherapy – Authors' reply. <i>Lancet Oncology, The, 2010, 11, 612-613.</i>	5.1	5