

Nader G Abraham

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1908433/nader-g-abraham-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

124
papers

5,958
citations

44
h-index

74
g-index

138
ext. papers

6,560
ext. citations

5.3
avg, IF

5.77
L-index

| # | Paper | IF | Citations |
|-----|---|------|-----------|
| 124 | Pharmacological and clinical aspects of heme oxygenase. <i>Pharmacological Reviews</i> , 2008 , 60, 79-127 | 22.5 | 903 |
| 123 | Heme oxygenase and the cardiovascular-renal system. <i>Free Radical Biology and Medicine</i> , 2005 , 39, 1-25 | 7.8 | 280 |
| 122 | Treatment of obese diabetic mice with a heme oxygenase inducer reduces visceral and subcutaneous adiposity, increases adiponectin levels, and improves insulin sensitivity and glucose tolerance. <i>Diabetes</i> , 2008 , 57, 1526-35 | 0.9 | 268 |
| 121 | Gene transfer of human heme oxygenase into coronary endothelial cells potentially promotes angiogenesis. <i>Journal of Cellular Biochemistry</i> , 1998 , 68, 121-7 | 4.7 | 194 |
| 120 | A heme oxygenase product, presumably carbon monoxide, mediates a vasodepressor function in rats. <i>Hypertension</i> , 1995 , 25, 166-9 | 8.5 | 145 |
| 119 | Heme oxygenase-1 induction remodels adipose tissue and improves insulin sensitivity in obesity-induced diabetic rats. <i>Hypertension</i> , 2009 , 53, 508-15 | 8.5 | 144 |
| 118 | Heme oxygenase-1 attenuates glucose-mediated cell growth arrest and apoptosis in human microvessel endothelial cells. <i>Circulation Research</i> , 2003 , 93, 507-14 | 15.7 | 130 |
| 117 | L-4F treatment reduces adiposity, increases adiponectin levels, and improves insulin sensitivity in obese mice. <i>Journal of Lipid Research</i> , 2008 , 49, 1658-69 | 6.3 | 129 |
| 116 | Heme oxygenase-mediated increases in adiponectin decrease fat content and inflammatory cytokines tumor necrosis factor-alpha and interleukin-6 in Zucker rats and reduce adipogenesis in human mesenchymal stem cells. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2008 , 325, 833-40 | 4.7 | 114 |
| 115 | Up-regulation of heme oxygenase provides vascular protection in an animal model of diabetes through its antioxidant and antiapoptotic effects. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2006 , 319, 1144-52 | 4.7 | 100 |
| 114 | HO-1 expression increases mesenchymal stem cell-derived osteoblasts but decreases adipocyte lineage. <i>Bone</i> , 2010 , 46, 236-43 | 4.7 | 97 |
| 113 | Adipocyte heme oxygenase-1 induction attenuates metabolic syndrome in both male and female obese mice. <i>Hypertension</i> , 2010 , 56, 1124-30 | 8.5 | 96 |
| 112 | Heme oxygenase-1 enhances renal mitochondrial transport carriers and cytochrome C oxidase activity in experimental diabetes. <i>Journal of Biological Chemistry</i> , 2006 , 281, 15687-93 | 5.4 | 93 |
| 111 | Translational Significance of Heme Oxygenase in Obesity and Metabolic Syndrome. <i>Trends in Pharmacological Sciences</i> , 2016 , 37, 17-36 | 13.2 | 91 |
| 110 | Overexpression of human heme oxygenase-1 attenuates endothelial cell sloughing in experimental diabetes. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2004 , 287, H2468-77 | 5.2 | 91 |
| 109 | Epoxyeicosatrienoic acid agonist regulates human mesenchymal stem cell-derived adipocytes through activation of HO-1-pAKT signaling and a decrease in PPAR γ . <i>Stem Cells and Development</i> , 2010 , 19, 1863-73 | 4.4 | 86 |
| 108 | Heme oxygenase-1 gene expression modulates angiotensin II-induced increase in blood pressure. <i>Hypertension</i> , 2004 , 43, 1221-6 | 8.5 | 84 |

| | | | |
|-----|---|------|----|
| 107 | Regulation of diabetic cardiomyopathy by caloric restriction is mediated by intracellular signaling pathways involving SIRT1 and PGC-1 β . <i>Cardiovascular Diabetology</i> , 2018 , 17, 111 | 8.7 | 82 |
| 106 | Heme oxygenase -1 gene therapy: recent advances and therapeutic applications. <i>Current Gene Therapy</i> , 2007 , 7, 89-108 | 4.3 | 80 |
| 105 | Heme oxygenase gene targeting to adipocytes attenuates adiposity and vascular dysfunction in mice fed a high-fat diet. <i>Hypertension</i> , 2012 , 60, 467-75 | 8.5 | 79 |
| 104 | Long-term treatment with the apolipoprotein A1 mimetic peptide increases antioxidants and vascular repair in type I diabetic rats. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2007 , 322, 514-20 | 4.7 | 79 |
| 103 | Increased HO-1 levels ameliorate fatty liver development through a reduction of heme and recruitment of FGF21. <i>Obesity</i> , 2014 , 22, 705-12 | 8 | 78 |
| 102 | Heme oxygenase: the key to renal function regulation. <i>American Journal of Physiology - Renal Physiology</i> , 2009 , 297, F1137-52 | 4.3 | 78 |
| 101 | Increased heme-oxygenase 1 expression in mesenchymal stem cell-derived adipocytes decreases differentiation and lipid accumulation via upregulation of the canonical Wnt signaling cascade. <i>Stem Cell Research and Therapy</i> , 2013 , 4, 28 | 8.3 | 76 |
| 100 | PGC-1 alpha regulates HO-1 expression, mitochondrial dynamics and biogenesis: Role of epoxyeicosatrienoic acid. <i>Prostaglandins and Other Lipid Mediators</i> , 2016 , 125, 8-18 | 3.7 | 70 |
| 99 | Carbon monoxide produced by isolated arterioles attenuates pressure-induced vasoconstriction. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2001 , 281, H350-8 | 5.2 | 69 |
| 98 | Crosstalk between EET and HO-1 downregulates Bach1 and adipogenic marker expression in mesenchymal stem cell derived adipocytes. <i>Prostaglandins and Other Lipid Mediators</i> , 2011 , 96, 54-62 | 3.7 | 66 |
| 97 | Involvement of reactive oxygen species in a feed-forward mechanism of Na/K-ATPase-mediated signaling transduction. <i>Journal of Biological Chemistry</i> , 2013 , 288, 34249-34258 | 5.4 | 65 |
| 96 | Cytochrome P450, drug metabolizing enzymes and arachidonic acid metabolism in bovine ocular tissues. <i>Current Eye Research</i> , 1987 , 6, 623-30 | 2.9 | 65 |
| 95 | HO-1 overexpression and underexpression: Clinical implications. <i>Archives of Biochemistry and Biophysics</i> , 2019 , 673, 108073 | 4.1 | 63 |
| 94 | Sensitivity of human tissue heme oxygenase to a new synthetic metalloporphyrin. <i>Hepatology</i> , 1989 , 10, 365-9 | 11.2 | 63 |
| 93 | pNaKtide inhibits Na/K-ATPase reactive oxygen species amplification and attenuates adipogenesis. <i>Science Advances</i> , 2015 , 1, e1500781 | 14.3 | 59 |
| 92 | CYP2J2 targeting to endothelial cells attenuates adiposity and vascular dysfunction in mice fed a high-fat diet by reprogramming adipocyte phenotype. <i>Hypertension</i> , 2014 , 64, 1352-61 | 8.5 | 56 |
| 91 | Fructose Mediated Non-Alcoholic Fatty Liver Is Attenuated by HO-1-SIRT1 Module in Murine Hepatocytes and Mice Fed a High Fructose Diet. <i>PLoS ONE</i> , 2015 , 10, e0128648 | 3.7 | 55 |
| 90 | Protective effect of HO-1 against oxidative stress in human hepatoma cell line (HepG2) is independent of telomerase enzyme activity. <i>International Journal of Biochemistry and Cell Biology</i> , 2002 , 34, 1619-28 | 5.6 | 54 |

| | | | |
|----|---|------|----|
| 89 | Vasoregulatory function of the heme-heme oxygenase-carbon monoxide system. <i>American Journal of Hypertension</i> , 2001 , 14, 62S-67S | 2.3 | 53 |
| 88 | Epoxyeicosatrienoic Acids Regulate Adipocyte Differentiation of Mouse 3T3 Cells, Via PGC-1 α Activation, Which Is Required for HO-1 Expression and Increased Mitochondrial Function. <i>Stem Cells and Development</i> , 2016 , 25, 1084-94 | 4.4 | 53 |
| 87 | EET agonist prevents adiposity and vascular dysfunction in rats fed a high fat diet via a decrease in Bach 1 and an increase in HO-1 levels. <i>Prostaglandins and Other Lipid Mediators</i> , 2012 , 98, 133-42 | 3.7 | 51 |
| 86 | Regulation of heme oxygenase gene expression by cobalt in rat liver and kidney. <i>FEBS Journal</i> , 1990 , 192, 577-82 | | 50 |
| 85 | Heme induced oxidative stress attenuates sirtuin1 and enhances adipogenesis in mesenchymal stem cells and mouse pre-adipocytes. <i>Journal of Cellular Biochemistry</i> , 2012 , 113, 1926-35 | 4.7 | 49 |
| 84 | EET intervention on Wnt1, NOV, and HO-1 signaling prevents obesity-induced cardiomyopathy in obese mice. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2017 , 313, H368-H380 | 5.2 | 48 |
| 83 | Rat mesenteric arterial dilator response to 11,12-epoxyeicosatrienoic acid is mediated by activating heme oxygenase. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2006 , 291, H1999-2002 | 5.2 | 48 |
| 82 | Bone marrow stem cell transplant into intra-bone cavity prevents type 2 diabetes: role of heme oxygenase-adiponectin. <i>Journal of Autoimmunity</i> , 2008 , 30, 128-35 | 15.5 | 46 |
| 81 | Lentiviral-human heme oxygenase targeting endothelium improved vascular function in angiotensin II animal model of hypertension. <i>Human Gene Therapy</i> , 2011 , 22, 271-82 | 4.8 | 44 |
| 80 | Attenuation of Na/K-ATPase Mediated Oxidant Amplification with pNaKtide Ameliorates Experimental Uremic Cardiomyopathy. <i>Scientific Reports</i> , 2016 , 6, 34592 | 4.9 | 43 |
| 79 | Cyclooxygenase-2 dependent metabolism of 20-HETE increases adiposity and adipocyte enlargement in mesenchymal stem cell-derived adipocytes. <i>Journal of Lipid Research</i> , 2013 , 54, 786-793 | 6.3 | 42 |
| 78 | 11,12-epoxyeicosatrienoic acid stimulates heme-oxygenase-1 in endothelial cells. <i>Prostaglandins and Other Lipid Mediators</i> , 2007 , 82, 155-61 | 3.7 | 40 |
| 77 | Targeting the Heme-Heme Oxygenase System to Prevent Severe Complications Following COVID-19 Infections. <i>Antioxidants</i> , 2020 , 9, | 7.1 | 37 |
| 76 | Therapeutic applications of human heme oxygenase gene transfer and gene therapy. <i>Current Pharmaceutical Design</i> , 2003 , 9, 2513-24 | 3.3 | 37 |
| 75 | Heme oxygenase (HO-1) rescue of adipocyte dysfunction in HO-2 deficient mice via recruitment of epoxyeicosatrienoic acids (EETs) and adiponectin. <i>Cellular Physiology and Biochemistry</i> , 2012 , 29, 99-110 | 3.9 | 36 |
| 74 | Retrovirus-mediated HO gene transfer into endothelial cells protects against oxidant-induced injury. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 1999 , 277, L127-33 | 5.8 | 35 |
| 73 | pNaKtide Attenuates Steatohepatitis and Atherosclerosis by Blocking Na/K-ATPase/ROS Amplification in C57Bl6 and ApoE Knockout Mice Fed a Western Diet. <i>Scientific Reports</i> , 2017 , 7, 193 | 4.9 | 34 |
| 72 | Agonists of epoxyeicosatrienoic acids reduce infarct size and ameliorate cardiac dysfunction via activation of HO-1 and Wnt1 canonical pathway. <i>Prostaglandins and Other Lipid Mediators</i> , 2015 , 116-117, 76-86 | 3.7 | 34 |

| | | | |
|----|---|------|----|
| 71 | PARP-1 inhibition protects the diabetic heart through activation of SIRT1-PGC-1 β axis. <i>Experimental Cell Research</i> , 2018 , 373, 112-118 | 4.2 | 31 |
| 70 | Epoxyeicosatrienoic Acid as Therapy for Diabetic and Ischemic Cardiomyopathy. <i>Trends in Pharmacological Sciences</i> , 2016 , 37, 945-962 | 13.2 | 30 |
| 69 | Modulation of cGMP by human HO-1 retrovirus gene transfer in pulmonary microvessel endothelial cells. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2002 , 283, L1117-24 | 5.8 | 30 |
| 68 | Epoxyeicosatrienoic intervention improves NAFLD in leptin receptor deficient mice by an increase in PGC1 β -HO-1-PGC1 β mitochondrial signaling. <i>Experimental Cell Research</i> , 2019 , 380, 180-187 | 4.2 | 28 |
| 67 | Downregulation of PGC-1 Prevents the Beneficial Effect of EET-Heme Oxygenase-1 on Mitochondrial Integrity and Associated Metabolic Function in Obese Mice. <i>Journal of Nutrition and Metabolism</i> , 2016 , 2016, 9039754 | 2.7 | 28 |
| 66 | Development of NASH in Obese Mice is Confounded by Adipose Tissue Increase in Inflammatory NOV and Oxidative Stress. <i>International Journal of Hepatology</i> , 2018 , 2018, 3484107 | 2.7 | 27 |
| 65 | Role of the heme oxygenases in abnormalities of the mesenteric circulation in cirrhotic rats. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2004 , 308, 636-43 | 4.7 | 27 |
| 64 | Oxidized HDL is a potent inducer of adipogenesis and causes activation of the Ang-II and 20-HETE systems in human obese females. <i>Prostaglandins and Other Lipid Mediators</i> , 2016 , 123, 68-77 | 3.7 | 27 |
| 63 | Heme oxygenase-2 deletion impairs macrophage function: implication in wound healing. <i>FASEB Journal</i> , 2015 , 29, 105-15 | 0.9 | 26 |
| 62 | Genetic Polymorphisms Complicate COVID-19 Therapy: Pivotal Role of HO-1 in Cytokine Storm. <i>Antioxidants</i> , 2020 , 9, | 7.1 | 26 |
| 61 | High-fat diet exacerbates renal dysfunction in SHR: reversal by induction of HO-1-adiponectin axis. <i>Obesity</i> , 2012 , 20, 945-53 | 8 | 25 |
| 60 | The renal cytochrome P-450 arachidonic acid system. <i>Pediatric Nephrology</i> , 1992 , 6, 490-8 | 3.2 | 25 |
| 59 | HO-1 Upregulation Attenuates Adipocyte Dysfunction, Obesity, and Isoprostane Levels in Mice Fed High Fructose Diets. <i>Journal of Nutrition and Metabolism</i> , 2014 , 2014, 980547 | 2.7 | 24 |
| 58 | The Na/K-ATPase Oxidant Amplification Loop Regulates Aging. <i>Scientific Reports</i> , 2018 , 8, 9721 | 4.9 | 24 |
| 57 | Elevated levels of heme oxygenase-1 activity and mRNA in peripheral blood adherent cells of acquired immunodeficiency syndrome patients. <i>American Journal of Hematology</i> , 1993 , 43, 19-23 | 7.1 | 23 |
| 56 | Oxidized HDL, Adipokines, and Endothelial Dysfunction: A Potential Biomarker Profile for Cardiovascular Risk in Women with Obesity. <i>Obesity</i> , 2019 , 27, 87-93 | 8 | 23 |
| 55 | Ablation of soluble epoxide hydrolase reprogram white fat to beige-like fat through an increase in mitochondrial integrity, HO-1-adiponectin in vitro and in vivo. <i>Prostaglandins and Other Lipid Mediators</i> , 2018 , 138, 1-8 | 3.7 | 21 |
| 54 | Cardioprotective Heme Oxygenase-1-PGC1 β Signaling in Epicardial Fat Attenuates Cardiovascular Risk in Humans as in Obese Mice. <i>Obesity</i> , 2019 , 27, 1634-1643 | 8 | 20 |

| | | | |
|----|---|-----|----|
| 53 | Positive Effects of Heme Oxygenase Upregulation on Adiposity and Vascular Dysfunction: Gene Targeting vs. Pharmacologic Therapy. <i>International Journal of Molecular Sciences</i> , 2019 , 20, | 6.3 | 19 |
| 52 | Existence of a Strong Correlation of Biomarkers and miRNA in Females with Metabolic Syndrome and Obesity in a Population of West Virginia. <i>International Journal of Medical Sciences</i> , 2017 , 14, 543-553 ³⁻⁷ | | 19 |
| 51 | Cold Press Pomegranate Seed Oil Attenuates Dietary-Obesity Induced Hepatic Steatosis and Fibrosis through Antioxidant and Mitochondrial Pathways in Obese Mice. <i>International Journal of Molecular Sciences</i> , 2020 , 21, | 6.3 | 19 |
| 50 | Ablation of adipose-HO-1 expression increases white fat over beige fat through inhibition of mitochondrial fusion and of PGC1 β in female mice. <i>Hormone Molecular Biology and Clinical Investigation</i> , 2017 , 31, | 1.3 | 18 |
| 49 | Caloric restriction ameliorates cardiomyopathy in animal model of diabetes. <i>Experimental Cell Research</i> , 2017 , 350, 147-153 | 4.2 | 17 |
| 48 | Functional expression of human heme oxygenase-1 (HO-1) driven by HO-1 promoter in vitro and in vivo. <i>Journal of Cellular Biochemistry</i> , 2002 , 85, 410-421 | 4.7 | 17 |
| 47 | Soluble epoxide hydrolase null mice exhibit female and male differences in regulation of vascular homeostasis. <i>Prostaglandins and Other Lipid Mediators</i> , 2015 , 120, 139-47 | 3.7 | 16 |
| 46 | Adipocyte Specific HO-1 Gene Therapy is Effective in Antioxidant Treatment of Insulin Resistance and Vascular Function in an Obese Mice Model. <i>Antioxidants</i> , 2020 , 9, | 7.1 | 16 |
| 45 | HO-1 induction improves the type-1 cardiorenal syndrome in mice with impaired angiotensin II-induced lymphocyte activation. <i>Hypertension</i> , 2013 , 62, 310-6 | 8.5 | 15 |
| 44 | Targeting Heme Oxygenase-1 in Cardiovascular and Kidney Disease. <i>Antioxidants</i> , 2019 , 8, | 7.1 | 14 |
| 43 | Cold-Pressed Oil Standardized to 3% Thymoquinone Potentiates Omega-3 Protection against Obesity-Induced Oxidative Stress, Inflammation, and Markers of Insulin Resistance Accompanied with Conversion of White to Beige Fat in Mice. <i>Antioxidants</i> , 2020 , 9, | 7.1 | 14 |
| 42 | Effect of acute and chronic treatment of tin on blood pressure in spontaneously hypertensive rats. <i>Tohoku Journal of Experimental Medicine</i> , 1992 , 166, 85-91 | 2.4 | 14 |
| 41 | Heme oxygenase-2/adiponectin protein-protein interaction in metabolic syndrome. <i>Biochemical and Biophysical Research Communications</i> , 2013 , 432, 606-11 | 3.4 | 13 |
| 40 | The Adipocyte Na/K-ATPase Oxidant Amplification Loop is the Central Regulator of Western Diet-Induced Obesity and Associated Comorbidities. <i>Scientific Reports</i> , 2019 , 9, 7927 | 4.9 | 12 |
| 39 | Uric Acid-Induced Adipocyte Dysfunction Is Attenuated by HO-1 Upregulation: Potential Role of Antioxidant Therapy to Target Obesity. <i>Stem Cells International</i> , 2016 , 2016, 8197325 | 5 | 12 |
| 38 | EET enhances renal function in obese mice resulting in restoration of HO-1-Mfn1/2 signaling, and decrease in hypertension through inhibition of sodium chloride co-transporter. <i>Prostaglandins and Other Lipid Mediators</i> , 2018 , 137, 30-39 | 3.7 | 12 |
| 37 | Kavain Reduces Induced Adipocyte Inflammation: Role of PGC-1 β Signaling. <i>Journal of Immunology</i> , 2018 , 201, 1491-1499 | 5.3 | 11 |
| 36 | Methods for measurements of heme oxygenase (HO) isoforms-mediated synthesis of carbon monoxide and HO-1 and HO-2 proteins. <i>Methods in Molecular Medicine</i> , 2003 , 86, 399-411 | | 11 |

| | | | |
|----|--|------|----|
| 35 | Functional expression of human heme oxygenase-1 gene in renal structure of spontaneously hypertensive rats. <i>Experimental Biology and Medicine</i> , 2003 , 228, 454-8 | 3.7 | 11 |
| 34 | Antioxidants condition pleiotropic vascular responses to exogenous H ₂ O ₂ : role of modulation of vascular TP receptors and the heme oxygenase system. <i>Antioxidants and Redox Signaling</i> , 2013 , 18, 471-80 | 8.4 | 10 |
| 33 | Central Role for Adipocyte Na,K-ATPase Oxidant Amplification Loop in the Pathogenesis of Experimental Uremic Cardiomyopathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2020 , 31, 1746-1760 | 12.7 | 9 |
| 32 | Beneficial Role of HO-1-SIRT1 Axis in Attenuating Angiotensin II-Induced Adipocyte Dysfunction. <i>International Journal of Molecular Sciences</i> , 2019 , 20, | 6.3 | 9 |
| 31 | The association of NOV/CCN3 with obstructive sleep apnea (OSA): preliminary evidence of a novel biomarker in OSA. <i>Hormone Molecular Biology and Clinical Investigation</i> , 2017 , 31, | 1.3 | 9 |
| 30 | Heme oxygenase-1 gene expression as a stress index to ocular irritation. <i>Current Eye Research</i> , 1999 , 19, 115-22 | 2.9 | 9 |
| 29 | Expression and Inducibility of Cytochrome P450 Iiia Family within Intrasplically Transplanted Fetal Hepatocytes. <i>Cell Transplantation</i> , 1996 , 5, 117-122 | 4 | 9 |
| 28 | Synergistic effect of heme and IL-1 on hematopoietic stromal regeneration after radiation. <i>American Journal of Hematology</i> , 1993 , 44, 172-8 | 7.1 | 8 |
| 27 | Heme Oxygenase-1 Upregulation: A Novel Approach in the Treatment of Cardiovascular Disease. <i>Antioxidants and Redox Signaling</i> , 2020 , 32, 1045-1060 | 8.4 | 8 |
| 26 | Milk thistle seed cold press oil attenuates markers of the metabolic syndrome in a mouse model of dietary-induced obesity. <i>Journal of Food Biochemistry</i> , 2020 , 44, e13522 | 3.3 | 7 |
| 25 | The Peroxisome Proliferator-Activated Receptor-Gamma Coactivator-1β-Heme Oxygenase 1 Axis, a Powerful Antioxidative Pathway with Potential to Attenuate Diabetic Cardiomyopathy. <i>Antioxidants and Redox Signaling</i> , 2020 , 32, 1273-1290 | 8.4 | 6 |
| 24 | Pathophysiology of chronic peripheral ischemia: new perspectives. <i>Therapeutic Advances in Chronic Disease</i> , 2020 , 11, 2040622319894466 | 4.9 | 6 |
| 23 | Oxidized HDL and Isoprostane Exert a Potent Adipogenic Effect on Stem Cells: Where in the Lineage? 2016 , 2, | | 6 |
| 22 | The pivotal role of heme Oxygenase-1 in reversing the pathophysiology and systemic complications of NAFLD. <i>Archives of Biochemistry and Biophysics</i> , 2021 , 697, 108679 | 4.1 | 6 |
| 21 | Comparative effect of heme analogues on hematopoiesis in lymphoproliferative disorders. <i>Leukemia and Lymphoma</i> , 1991 , 5, 179-85 | 1.9 | 5 |
| 20 | OX-HDL: A Starring Role in Cardiorenal Syndrome and the Effects of Heme Oxygenase-1 Intervention. <i>Diagnostics</i> , 2020 , 10, | 3.8 | 5 |
| 19 | Therapeutic approaches to diabetic cardiomyopathy: Targeting the antioxidant pathway. <i>Prostaglandins and Other Lipid Mediators</i> , 2020 , 150, 106454 | 3.7 | 4 |
| 18 | Coexpression of erythropoietin and heme oxygenase genes in Hep3B cells. <i>Hepatology</i> , 1993 , 17, 861-868 | 1.2 | 4 |

| | | |
|----|---|--------|
| 17 | Relationship between dietary sodium and sugar intake: A cross-sectional study of the National Health and Nutrition Examination Survey 2001-2016. <i>Journal of Clinical Hypertension</i> , 2020 , 22, 1694-1702 ^{2,3} | 4 |
| 16 | Heme-oxygenase and lipid mediators in obesity and associated cardiometabolic diseases: Therapeutic implications. <i>Pharmacology & Therapeutics</i> , 2021 , 107975 | 13.9 4 |
| 15 | Promotive Effects of a Silk Film on Epidermal Recovery from Full-Thickness Skin Wounds. <i>Proceedings of the Society for Experimental Biology and Medicine</i> , 2000 , 225, 58-64 | 3 |
| 14 | The Pivotal Role of Adipocyte-Na K peptide in Reversing Systemic Inflammation in Obesity and COVID-19 in the Development of Heart Failure. <i>Antioxidants</i> , 2020 , 9, | 7.1 3 |
| 13 | Enzymatic Activity and Expression of Cytochrome P450 LA within Intrasplenically Transplanted Fetal Hepatocytes in Spontaneously Hypertensive Rats. <i>Cell Transplantation</i> , 1997 , 6, 531-534 | 4 2 |
| 12 | Oxidant-Induced Alterations in the Adipocyte Transcriptome: Role of the Na,K-ATPase Oxidant Amplification Loop. <i>International Journal of Molecular Sciences</i> , 2020 , 21, | 6.3 2 |
| 11 | Cyclooxygenase activity is regulated by the heme oxygenase system in microvessel endothelial cells. <i>Advances in Experimental Medicine and Biology</i> , 2003 , 525, 59-66 | 3.6 2 |
| 10 | The Essential Role of the L4F-Adiponectin Regulatory Axis: Leading to Improvements in the Metabolic Profile of Diabetes Mellitus. <i>FASEB Journal</i> , 2008 , 22, 1226.43 | 0.9 1 |
| 9 | Can charcoal improve outcomes in COVID-19 infections?. <i>Medical Hypotheses</i> , 2020 , 144, 110176 | 3.8 1 |
| 8 | Treatment of Obese Diabetic Mice with an Heme Oxygenase Inducer Reduces Visceral and Abdominal Adiposity, Increases Adiponectin Levels and Improves Insulin Sensitivity and Glucose Tolerance. <i>FASEB Journal</i> , 2008 , 22, 642-642 | 0.9 |
| 7 | EET-agonist Prevents and Reverses Heart Failure in Obesity Induced Diabetic Cardiomyopathy. <i>FASEB Journal</i> , 2018 , 32, 561.7 | 0.9 |
| 6 | EET Enhances Renal Function in Obese Mice Resulting in Restoration of Mfn1/2 Signaling and a Decrease in Hypertension Through Inhibition of Sodium Chloride Co-Transporter. <i>FASEB Journal</i> , 2018 , 32, 561.13 | 0.9 |
| 5 | Characterization of Diabetic Cardiomyopathy: A Role for KDAC Activity. <i>FASEB Journal</i> , 2015 , 29, LB557 | 0.9 |
| 4 | Heme Oxygenase-1 Induction Modulates Hypoxic Pulmonary Vasoconstriction through Upregulation of ecSOD. <i>FASEB Journal</i> , 2009 , 23, 1002.9 | 0.9 |
| 3 | Apolipoprotein Mimetic Peptide (L-4F) Regulation of Adiposity via Increases in Estradiol/Testosterone Ratio in Obese Female Mice. <i>FASEB Journal</i> , 2010 , 24, 588.5 | 0.9 |
| 2 | L-4F Rescues the Metabolic Syndrome Phenotype of HO-2 Null Mice via Insulin, Adiponectin, & LKB1 Signaling Pathways. <i>FASEB Journal</i> , 2010 , 24, 570.2 | 0.9 |
| 1 | L-4F Improves Metabolic Syndrome Phenotype in HO-2 Null Mice by Decreasing NFkB Activity & Increasing Adiponectin Levels. <i>FASEB Journal</i> , 2010 , 24, 1035.5 | 0.9 |