Olivier Rohr

List of Publications by Citations

Source: https://exaly.com/author-pdf/1912754/olivier-rohr-publications-by-citations.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

36
papers

1,726
citations

h-index

38
ext. papers

20
h-index

7.5
avg, IF

L-index

#	Paper	IF	Citations
36	Recruitment of chromatin-modifying enzymes by CTIP2 promotes HIV-1 transcriptional silencing. <i>EMBO Journal</i> , 2007 , 26, 412-23	13	270
35	An In-Depth Comparison of Latency-Reversing Agent Combinations in Various In Vitro and Ex Vivo HIV-1 Latency Models Identified Bryostatin-1+JQ1 and Ingenol-B+JQ1 to Potently Reactivate Viral Gene Expression. <i>PLoS Pathogens</i> , 2015 , 11, e1005063	7.6	188
34	Molecular mechanisms of HIV-1 persistence in the monocyte-macrophage lineage. <i>Retrovirology</i> , 2010 , 7, 32	3.6	131
33	Regulation of HIV-1 gene transcription: from lymphocytes to microglial cells. <i>Journal of Leukocyte Biology</i> , 2003 , 74, 736-49	6.5	107
32	Microglial Cells: The Main HIV-1 Reservoir in the Brain. <i>Frontiers in Cellular and Infection Microbiology</i> , 2019 , 9, 362	5.9	97
31	p21(WAF1) gene promoter is epigenetically silenced by CTIP2 and SUV39H1. <i>Oncogene</i> , 2009 , 28, 3380-	-99.2	94
30	COUP-TF interacting protein 2 represses the initial phase of HIV-1 gene transcription in human microglial cells. <i>Nucleic Acids Research</i> , 2005 , 33, 2318-31	20.1	80
29	Targeting the Brain Reservoirs: Toward an HIV Cure. Frontiers in Immunology, 2016, 7, 397	8.4	68
28	CTIP2 is a negative regulator of P-TEFb. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 12655-60	11.5	66
27	Sequential treatment with 5-aza-2Rdeoxycytidine and deacetylase inhibitors reactivates HIV-1. <i>EMBO Molecular Medicine</i> , 2016 , 8, 117-38	12	58
26	Recruitment of Tat to heterochromatin protein HP1 via interaction with CTIP2 inhibits human immunodeficiency virus type 1 replication in microglial cells. <i>Journal of Virology</i> , 2003 , 77, 5415-27	6.6	56
25	LSD1 cooperates with CTIP2 to promote HIV-1 transcriptional silencing. <i>Nucleic Acids Research</i> , 2012 , 40, 1904-15	20.1	55
24	HIV-1 regulation of latency in the monocyte-macrophage lineage and in CD4+ T lymphocytes. <i>Journal of Leukocyte Biology</i> , 2010 , 87, 575-88	6.5	50
23	Functional interactions between C/EBP, Sp1, and COUP-TF regulate human immunodeficiency virus type 1 gene transcription in human brain cells. <i>Journal of Virology</i> , 2000 , 74, 65-73	6.6	48
22	Current Status of Latency Reversing Agents Facing the Heterogeneity of HIV-1 Cellular and Tissue Reservoirs. <i>Frontiers in Microbiology</i> , 2019 , 10, 3060	5.7	46
21	On the way to find a cure: Purging latent HIV-1 reservoirs. <i>Biochemical Pharmacology</i> , 2017 , 146, 10-22	6	40
20	Resveratrol Inhibits HCoV-229E and SARS-CoV-2 Coronavirus Replication In Vitro. <i>Viruses</i> , 2021 , 13,	6.2	40

(2021-2012)

19	Achieving a cure for HIV infection: do we have reasons to be optimistic?. <i>Journal of Antimicrobial Chemotherapy</i> , 2012 , 67, 1063-74	5.1	39
18	HMGA1 recruits CTIP2-repressed P-TEFb to the HIV-1 and cellular target promoters. <i>Nucleic Acids Research</i> , 2014 , 42, 4962-71	20.1	36
17	Genome-wide binding map of the HIV-1 Tat protein to the human genome. <i>PLoS ONE</i> , 2011 , 6, e26894	3.7	27
16	Protein Kinase C-Mediated Phosphorylation of BCL11B at Serine 2 Negatively Regulates Its Interaction with NuRD Complexes during CD4+ T-Cell Activation. <i>Molecular and Cellular Biology</i> , 2016 , 36, 1881-98	4.8	20
15	The many lives of CTIP2: from AIDS to cancer and cardiac hypertrophy. <i>Journal of Cellular Physiology</i> , 2014 , 229, 533-7	7	18
14	Human-Phosphate-Binding-Protein inhibits HIV-1 gene transcription and replication. <i>Virology Journal</i> , 2011 , 8, 352	6.1	16
13	HIC1 controls cellular- and HIV-1- gene transcription via interactions with CTIP2 and HMGA1. <i>Scientific Reports</i> , 2016 , 6, 34920	4.9	14
12	HIV-1 Vpr mediates the depletion of the cellular repressor CTIP2 to counteract viral gene silencing. <i>Scientific Reports</i> , 2019 , 9, 13154	4.9	12
11	Improving combination antiretroviral therapy by targeting HIV-1 gene transcription. <i>Expert Opinion on Therapeutic Targets</i> , 2016 , 20, 1311-1324	6.4	12
10	Targeting the DNA-PK complex: Its rationale use in cancer and HIV-1 infection. <i>Biochemical Pharmacology</i> , 2019 , 160, 80-91	6	10
10 9		4.9	10
	Pharmacology, 2019 , 160, 80-91	4.9	
9	Pharmacology, 2019 , 160, 80-91 Inhibition of HIV-1 gene transcription by KAP1 in myeloid lineage. <i>Scientific Reports</i> , 2021 , 11, 2692 In Trauma Patients, the Occurrence of Early-Onset Nosocomial Infections is Associated With	4.9	8
9	Pharmacology, 2019, 160, 80-91 Inhibition of HIV-1 gene transcription by KAP1 in myeloid lineage. Scientific Reports, 2021, 11, 2692 In Trauma Patients, the Occurrence of Early-Onset Nosocomial Infections is Associated With Increased Plasma Concentrations of Chromogranin A. Shock, 2018, 49, 522-528 Pseudomonas DING proteins as human transcriptional regulators and HIV-1 antagonists. Virology	4·9 3·4	8 5
9 8 7	Inhibition of HIV-1 gene transcription by KAP1 in myeloid lineage. <i>Scientific Reports</i> , 2021 , 11, 2692 In Trauma Patients, the Occurrence of Early-Onset Nosocomial Infections is Associated With Increased Plasma Concentrations of Chromogranin A. <i>Shock</i> , 2018 , 49, 522-528 Pseudomonas DING proteins as human transcriptional regulators and HIV-1 antagonists. <i>Virology Journal</i> , 2013 , 10, 234 Evolution of a concept: From accessory protein to key virulence factor, the case of HIV-1 Vpr.	4·9 3·4 6.1	855
9 8 7 6	Inhibition of HIV-1 gene transcription by KAP1 in myeloid lineage. <i>Scientific Reports</i> , 2021 , 11, 2692 In Trauma Patients, the Occurrence of Early-Onset Nosocomial Infections is Associated With Increased Plasma Concentrations of Chromogranin A. <i>Shock</i> , 2018 , 49, 522-528 Pseudomonas DING proteins as human transcriptional regulators and HIV-1 antagonists. <i>Virology Journal</i> , 2013 , 10, 234 Evolution of a concept: From accessory protein to key virulence factor, the case of HIV-1 Vpr. <i>Biochemical Pharmacology</i> , 2020 , 180, 114128 Analysis of RNA binding properties of human Ku protein reveals its interactions with 7SK snRNA	4.9 3.4 6.1	854
9 8 7 6	Inhibition of HIV-1 gene transcription by KAP1 in myeloid lineage. <i>Scientific Reports</i> , 2021 , 11, 2692 In Trauma Patients, the Occurrence of Early-Onset Nosocomial Infections is Associated With Increased Plasma Concentrations of Chromogranin A. <i>Shock</i> , 2018 , 49, 522-528 Pseudomonas DING proteins as human transcriptional regulators and HIV-1 antagonists. <i>Virology Journal</i> , 2013 , 10, 234 Evolution of a concept: From accessory protein to key virulence factor, the case of HIV-1 Vpr. <i>Biochemical Pharmacology</i> , 2020 , 180, 114128 Analysis of RNA binding properties of human Ku protein reveals its interactions with 7SK snRNA and protein components of 7SK snRNP complex. <i>Biochimie</i> , 2020 , 171-172, 110-123	4.9 3.4 6.1 6	8544

Novel role of UHRF1 in the epigenetic repression of the latent HIV-1.. EBioMedicine, 2022, 79, 103985 8.8 o