## Carmelo J Rizzo

## List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1919169/carmelo-j-rizzo-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

3,150 110 31 50 h-index g-index citations papers 4.69 6.3 112 3,370 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
110	Enzymatic bypass and the structural basis of miscoding opposite the DNA adduct 1,N-ethenodeoxyguanosine by human DNA translesion polymerase \( \partit{Journal of Biological Chemistry, \textbf{2021}, 296, 100642} \)	5.4	2
109	Kinetics of DNA Adducts and Abasic Site Formation in Tissues of Mice Treated with a Nitrogen Mustard. <i>Chemical Research in Toxicology</i> , <b>2020</b> , 33, 988-998	4	2
108	Quantitation of Apurinic/Apyrimidinic Sites in Isolated DNA and in Mammalian Tissue with a Reduced Level of Artifacts. <i>Analytical Chemistry</i> , <b>2019</b> , 91, 7403-7410	7.8	19
107	Configurational and Conformational Equilibria of N-(2-Deoxy-d-erythro-pentofuranosyl)-2,6-diamino-3,4-dihydro-4-oxo-5-N-methylformamidopyrimidine (MeFapy-dG) Lesion in DNA. <i>Chemical Research in Toxicology</i> , <b>2018</b> ,	4	2
106	31, 924-935 Error-prone replication bypass of the imidazole ring-opened formamidopyrimidine deoxyguanosine adduct. <i>Environmental and Molecular Mutagenesis</i> , <b>2017</b> , 58, 182-189	3.2	5
105	Mutagenic potential of nitrogen mustard-induced formamidopyrimidine DNA adduct: Contribution of the non-canonical nomer. <i>Journal of Biological Chemistry</i> , <b>2017</b> , 292, 18790-18799	5.4	7
104	Novel genomic island modifies DNA with 7-deazaguanine derivatives. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2016</b> , 113, E1452-9	11.5	66
103	Catalysts of DNA Strand Cleavage at Apurinic/Apyrimidinic Sites. Scientific Reports, 2016, 6, 28894	4.9	12
102	Translesion Synthesis of the N(2)-2'-Deoxyguanosine Adduct of the Dietary Mutagen IQ in Human Cells: Error-Free Replication by DNA Polymerase and Mutagenic Bypass by DNA Polymerases pand Rev1. Chemical Research in Toxicology, <b>2016</b> , 29, 1549-59	4	8
101	Base-Displaced Intercalated Conformation of the 2-Amino-3-methylimidazo[4,5-f]quinoline N(2)-dG DNA Adduct Positioned at the Nonreiterated G(1) in the Narl Restriction Site. <i>Chemical Research in Toxicology</i> , <b>2015</b> , 28, 1455-68	4	5
100	Structural Basis for Error-Free Bypass of the 5-N-Methylformamidopyrimidine-dG Lesion by Human DNA Polymerase and Sulfolobus solfataricus P2 Polymerase IV. <i>Journal of the American Chemical Society</i> , <b>2015</b> , 137, 7011-4	16.4	12
99	Differential repair of etheno-DNA adducts by bacterial and human AlkB proteins. <i>DNA Repair</i> , <b>2015</b> , 30, 1-10	4.3	26
98	Characterization of nitrogen mustard formamidopyrimidine adduct formation of bis(2-chloroethyl)ethylamine with calf thymus DNA and a human mammary cancer cell line. <i>Chemical Research in Toxicology</i> , <b>2015</b> , 28, 1850-60	4	16
97	Next-generation sequencing reveals the biological significance of the N(2),3-ethenoguanine lesion in vivo. <i>Nucleic Acids Research</i> , <b>2015</b> , 43, 5489-500	20.1	28
96	DNA polymerases and Leooperatively perform mutagenic translesion synthesis of the C8-2'-deoxyguanosine adduct of the dietary mutagen IQ in human cells. <i>Nucleic Acids Research</i> , <b>2015</b> , 43, 8340-51	20.1	11
95	Mechanism of repair of acrolein- and malondialdehyde-derived exocyclic guanine adducts by the Eketoglutarate/Fe(II) dioxygenase AlkB. <i>Chemical Research in Toxicology</i> , <b>2014</b> , 27, 1619-31	4	20
94	Synthesis of G(CH)G Trimethylene DNA interstrand cross-links. <i>Current Protocols in Nucleic Acid Chemistry</i> , <b>2014</b> , 5, 5.14.1-5.14.15	0.5	2

## (2011-2014)

93	Characterization of the deoxyguanosine-lysine cross-link of methylglyoxal. <i>Chemical Research in Toxicology</i> , <b>2014</b> , 27, 1019-29	4	10
92	Synthesis and characterization of oligonucleotides containing a nitrogen mustard formamidopyrimidine monoadduct of deoxyguanosine. <i>Chemical Research in Toxicology</i> , <b>2014</b> , 27, 1610	-8	12
91	Base-displaced intercalation of the 2-amino-3-methylimidazo[4,5-f]quinolone N2-dG adduct in the NarI DNA recognition sequence. <i>Nucleic Acids Research</i> , <b>2014</b> , 42, 3450-63	20.1	13
90	Xeroderma pigmentosum complementation group C protein (XPC) serves as a general sensor of damaged DNA. <i>DNA Repair</i> , <b>2013</b> , 12, 947-53	4.3	36
89	Ring-opening of the EDH-PdG adduct promotes error-free bypass by the Sulfolobus solfataricus DNA polymerase Dpo4. <i>Chemical Research in Toxicology</i> , <b>2013</b> , 26, 1348-60	4	3
88	Mutagenic spectra arising from replication bypass of the 2,6-diamino-4-hydroxy-N(5)-methyl formamidopyrimidine adduct in primate cells. <i>Chemical Research in Toxicology</i> , <b>2013</b> , 26, 1108-14	4	15
87	Simplified synthesis of individual stereoisomers of the 4-hydroxynonenal adducts of deoxyguanosine. <i>Tetrahedron Letters</i> , <b>2013</b> , 54, 4289-4291	2	6
86	Replication, repair, and translesion polymerase bypass of NEbxopropenyl-2'-deoxyadenosine. <i>Biochemistry</i> , <b>2013</b> , 52, 8766-76	3.2	6
85	Replication bypass of the trans-4-Hydroxynonenal-derived (6S,8R,11S)-1,N(2)-deoxyguanosine DNA adduct by the sulfolobus solfataricus DNA polymerase IV. <i>Chemical Research in Toxicology</i> , <b>2012</b> , 25, 42	2 <sup>4</sup> 35	8
84	Replication of the 2,6-diamino-4-hydroxy-N(5)-(methyl)-formamidopyrimidine (MeFapy-dGuo) adduct by eukaryotic DNA polymerases. <i>Chemical Research in Toxicology</i> , <b>2012</b> , 25, 1652-61	4	14
83	Replication of N2,3-Ethenoguanine by DNA Polymerases. <i>Angewandte Chemie</i> , <b>2012</b> , 124, 5562-5565	3.6	1
82	Replication of N2,3-ethenoguanine by DNA polymerases. <i>Angewandte Chemie - International Edition</i> , <b>2012</b> , 51, 5466-9	16.4	26
81	Basis of miscoding of the DNA adduct N2,3-ethenoguanine by human Y-family DNA polymerases. Journal of Biological Chemistry, <b>2012</b> , 287, 35516-35526	5.4	25
80	Synthesis of the four stereoisomers of 2,3-epoxy-4-hydroxynonanal and their reactivity with deoxyguanosine. <i>Organic and Biomolecular Chemistry</i> , <b>2011</b> , 9, 1960-71	3.9	7
79	EHydroxy-1,N2-propano-2'-deoxyguanosine DNA adduct conjugates the N-terminal amine of the KWKK peptide via a carbinolamine linkage. <i>Chemical Research in Toxicology</i> , <b>2011</b> , 24, 1123-33	4	5
78	1,N2-Etheno-2'-deoxyguanosine adopts the syn conformation about the glycosyl bond when mismatched with deoxyadenosine. <i>Chemical Research in Toxicology</i> , <b>2011</b> , 24, 1071-9	4	1
77	Deoxyguanosine forms a bis-adduct with E,E-muconaldehyde, an oxidative metabolite of benzene: implications for the carcinogenicity of benzene. <i>Chemical Research in Toxicology</i> , <b>2011</b> , 24, 1944-56	4	2
76	Formation of a N2-dG:N2-dG carbinolamine DNA cross-link by the trans-4-hydroxynonenal-derived (6S,8R,11S) 1,N2-dG adduct. <i>Journal of the American Chemical Society</i> , <b>2011</b> , 133, 16101-10	16.4	10

75	Selective Incision of the alpha-N-Methyl-Formamidopyrimidine Anomer by Escherichia coli Endonuclease IV. <i>Journal of Nucleic Acids</i> , <b>2010</b> , 2010,	2.3	23
74	The mouse ortholog of NEIL3 is a functional DNA glycosylase in vitro and in vivo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2010</b> , 107, 4925-30	11.5	144
73	In vitro bypass of the major malondialdehyde- and base propenal-derived DNA adduct by human Y-family DNA polymerases [[]]and Rev1. <i>Biochemistry</i> , <b>2010</b> , 49, 8415-24	3.2	21
7 <del>2</del>	Structure of the 1,N(2)-etheno-2'-deoxyguanosine lesion in the 3'-G(epsilon dG)T-5' sequence opposite a one-base deletion. <i>Biochemistry</i> , <b>2010</b> , 49, 2615-26	3.2	8
71	Novel enzymatic function of DNA polymerase nu in translesion DNA synthesis past major groove DNA-peptide and DNA-DNA cross-links. <i>Chemical Research in Toxicology</i> , <b>2010</b> , 23, 689-95	4	50
70	Formation of deoxyguanosine cross-links from calf thymus DNA treated with acrolein and 4-hydroxy-2-nonenal. <i>Chemical Research in Toxicology</i> , <b>2010</b> , 23, 1701-13	4	29
69	Minor groove orientation of the KWKK peptide tethered via the N-terminal amine to the acrolein-derived 1,N2-gamma-hydroxypropanodeoxyguanosine lesion with a trimethylene linkage. <i>Biochemistry</i> , <b>2010</b> , 49, 6155-64	3.2	11
68	Comparison of the in vitro replication of the 7-(2-oxoheptyl)-1,N2-etheno-2'-deoxyguanosine and 1,N2-etheno-2'-deoxyguanosine lesions by Sulfolobus solfataricus P2 DNA polymerase IV (Dpo4). <i>Chemical Research in Toxicology</i> , <b>2010</b> , 23, 1330-41	4	8
67	The C8-2'-deoxyguanosine adduct of 2-amino-3-methylimidazo[1,2-d]naphthalene, a carbocyclic analogue of the potent mutagen 2-amino-3-methylimidazo[4,5-f]quinoline, is a block to replication in vitro. <i>Chemical Research in Toxicology</i> , <b>2010</b> , 23, 1076-88	4	9
66	DNA cross-link induced by trans-4-hydroxynonenal. <i>Environmental and Molecular Mutagenesis</i> , <b>2010</b> , 51, 625-34	3.2	35
65	Structure-Function Relationships in Miscoding by Sulfolobus solfataricus DNA Polymerase Dpo4. Journal of Biological Chemistry, <b>2009</b> , 284, 17687-17699	5.4	31
64	Versatility of Y-family Sulfolobus solfataricus DNA polymerase Dpo4 in translesion synthesis past bulky N2-alkylguanine adducts. <i>Journal of Biological Chemistry</i> , <b>2009</b> , 284, 3563-76	5.4	59
63	Oxidation and glycolytic cleavage of etheno and propano DNA base adducts. <i>Biochemistry</i> , <b>2009</b> , 48, 800-9	3.2	17
62	Translesion DNA synthesis by human DNA polymerase eta on templates containing a pyrimidopurinone deoxyguanosine adduct, 3-(2'-deoxy-beta-d-erythro-pentofuranosyl)pyrimido-[1,2-a]purin-10(3H)-one. <i>Biochemistry</i> , <b>2009</b> ,	3.2	11
61	Conformational interconversion of the trans-4-hydroxynonenal-derived (6S,8R,11S) 1,N(2)-deoxyguanosine adduct when mismatched with deoxyadenosine in DNA. <i>Chemical Research in Toxicology</i> , <b>2009</b> , 22, 187-200	4	12
60	Structural and functional analysis of Sulfolobus solfataricus Y-family DNA polymerase Dpo4-catalyzed bypass of the malondialdehyde-deoxyguanosine adduct. <i>Biochemistry</i> , <b>2009</b> , 48, 7079-8	38 <sup>3.2</sup>	26
59	Stereospecific formation of the (R)-gamma-hydroxytrimethylene interstrand N2-dG:N2-dG cross-link arising from the gamma-OH-1,N2-propano-2'-deoxyguanosine adduct in the 5'-CpG-3' DNA sequence. <i>Journal of the American Chemical Society</i> , <b>2009</b> , 131, 8416-24	16.4	13
58	Chemistry and biology of DNA containing 1,N(2)-deoxyguanosine adducts of the alpha,beta-unsaturated aldehydes acrolein, crotonaldehyde, and 4-hydroxynonenal. <i>Chemical Research in Toxicology</i> , <b>2009</b> , 22, 759-78	4	141

57	Replication past the N5-methyl-formamidopyrimidine lesion of deoxyguanosine by DNA polymerases and an improved procedure for sequence analysis of in vitro bypass products by mass spectrometry. <i>Chemical Research in Toxicology</i> , <b>2009</b> , 22, 1086-95	4	25
56	Steric and electrostatic effects at the C2 atom substituent influence replication and miscoding of the DNA deamination product deoxyxanthosine and analogs by DNA polymerases. <i>Journal of Molecular Biology</i> , <b>2009</b> , 392, 251-69	6.5	26
55	Checkpoint signaling from a single DNA interstrand crosslink. <i>Molecular Cell</i> , <b>2009</b> , 35, 704-15	17.6	88
54	Structure-function relationships in miscoding by Sulfolobus solfataricus DNA polymerase Dpo4: guanine N2,N2-dimethyl substitution produces inactive and miscoding polymerase complexes. <i>Journal of Biological Chemistry</i> , <b>2009</b> , 284, 17687-99	5.4	32
53	Mutagenic potential of DNA-peptide crosslinks mediated by acrolein-derived DNA adducts. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , <b>2008</b> , 637, 161-72	3.3	45
52	Structure of the 1,N2-ethenodeoxyguanosine adduct opposite cytosine in duplex DNA: Hoogsteen base pairing at pH 5.2. <i>Chemical Research in Toxicology</i> , <b>2008</b> , 21, 1795-805	4	14
51	The stereochemistry of trans-4-hydroxynonenal-derived exocyclic 1,N2-2'-deoxyguanosine adducts modulates formation of interstrand cross-links in the 5'-CpG-3' sequence. <i>Biochemistry</i> , <b>2008</b> , 47, 11457	<del>-32</del>	17
50	The formamidopyrimidine derivative of 7-(2-oxoethyl)-2'-deoxyguanosine. <i>Chemical Research in Toxicology</i> , <b>2008</b> , 21, 1777-86	4	3
49	Rearrangement of the (6S,8R,11S) and (6R,8S,11R) exocyclic 1,N2-deoxyguanosine adducts of trans-4-hydroxynonenal to N2-deoxyguanosine cyclic hemiacetal adducts when placed complementary to cytosine in duplex DNA. <i>Journal of the American Chemical Society</i> , <b>2008</b> , 130, 10898-5	16.4 <b>906</b>	16
48	Replication bypass of the acrolein-mediated deoxyguanine DNA-peptide cross-links by DNA polymerases of the DinB family. <i>Chemical Research in Toxicology</i> , <b>2008</b> , 21, 1983-90	4	58
47	Interstrand DNA cross-links induced by alpha, beta-unsaturated aldehydes derived from lipid peroxidation and environmental sources. <i>Accounts of Chemical Research</i> , <b>2008</b> , 41, 793-804	24.3	142
46	Chemical properties of oxopropenyl adducts of purine and pyrimidine nucleosides and their reactivity toward amino acid cross-link formation. <i>Journal of the American Chemical Society</i> , <b>2008</b> , 130, 2195-201	16.4	16
45	Site-specific synthesis and characterization of oligonucleotides containing an N6-(2-deoxy-D-erythro-pentofuranosyl)-2,6-diamino-3,4-dihydro-4-oxo-5-N-methylformamidopyrimiding lesion, the consequence of the contact	e <sub>4</sub>	26
44	Toxicology, <b>2008</b> , 21, 2324-33 "One-pot" syntheses of malondialdehyde adducts of nucleosides. <i>Nucleosides, Nucleotides and Nucleic Acids</i> , <b>2008</b> , 27, 103-9	1.4	11
43	Role for DNA polymerase kappa in the processing of N2-N2-guanine interstrand cross-links. <i>Journal of Biological Chemistry</i> , <b>2008</b> , 283, 17075-82	5.4	111
42	The modulation of topoisomerase I-mediated DNA cleavage and the induction of DNA-topoisomerase I crosslinks by crotonaldehyde-derived DNA adducts. <i>Nucleic Acids Research</i> , <b>2008</b> , 36, 4128-36	20.1	22
41	Chemical and electrochemical oxidation of C8-arylamine adducts of 2'-deoxyguanosine. <i>Journal of the American Chemical Society</i> , <b>2007</b> , 129, 2074-81	16.4	22
40	Stereochemistry modulates the stability of reduced interstrand cross-links arising from R- and S-alpha-CH3-gamma-OH-1,N2-propano-2'-deoxyguanosine in the 5'-CpG-3' DNA sequence.  Biochemistry, 2007, 46, 2608-21	3.2	16

39	Conformational differences of the C8-deoxyguanosine adduct of 2-amino-3-methylimidazo[4,5-f]quinoline (IQ) within the Narl recognition sequence. <i>Chemical Research in Toxicology</i> , <b>2007</b> , 20, 445-54	4	28
38	DNA sequence modulates the conformation of the food mutagen 2-amino-3-methylimidazo[4,5-f]quinoline in the recognition sequence of the Narl restriction enzyme. <i>Biochemistry</i> , <b>2007</b> , 46, 8498-516	3.2	34
37	Structure of the 1,N2-etheno-2'-deoxyguanosine adduct in duplex DNA at pH 8.6. <i>Chemical Research in Toxicology</i> , <b>2007</b> , 20, 1601-11	4	15
36	Mechanism of 1,N2-etheno-2'-deoxyguanosine formation from epoxyaldehydes. <i>Chemical Research in Toxicology</i> , <b>2007</b> , 20, 1685-92	4	24
35	Metabolism and elimination of the endogenous DNA adduct, 3-(2-deoxy-beta-D-erythropentofuranosyl)-pyrimido[1,2-alpha]purine-10(3H)-one, in the rat. <i>Journal of Biological Chemistry</i> , <b>2007</b> , 282, 36257-64	5.4	13
34	Synthesis of oligonucleotides containing the N2-deoxyguanosine adduct of the dietary carcinogen 2-amino-3-methylimidazo[4,5-f]quinoline. <i>Chemical Research in Toxicology</i> , <b>2007</b> , 20, 1972-9	4	10
33	Biochemical basis of genotoxicity of heterocyclic arylamine food mutagens: Human DNA polymerase eta selectively produces a two-base deletion in copying the N2-guanyl adduct of 2-amino-3-methylimidazo[4,5-f]quinoline but not the C8 adduct at the Narl G3 site. <i>Journal of</i>	5.4	37
32	Base-displaced intercalated structure of the food mutagen 2-amino-3-methylimidazo[4,5-f]quinoline in the recognition sequence of the Narl restriction enzyme, a hotspot for -2 bp deletions. <i>Journal of the American Chemical Society</i> , <b>2006</b> , 128, 10085-95	16.4	57
31	Stereospecific formation of interstrand carbinolamine DNA cross-links by crotonaldehyde- and acetaldehyde-derived alpha-CH3-gamma-OH-1,N2-propano-2'-deoxyguanosine adducts in the 5'-CpG-3' sequence. <i>Chemical Research in Toxicology</i> , <b>2006</b> , 19, 195-208	4	49
30	Translesion synthesis across 1,N2-ethenoguanine by human DNA polymerases. <i>Chemical Research in Toxicology</i> , <b>2006</b> , 19, 879-86	4	52
29	Site-specific synthesis of oligonucleotides containing malondialdehyde adducts of deoxyguanosine and deoxyadenosine via a postsynthetic modification strategy. <i>Chemical Research in Toxicology</i> , <b>2006</b> , 19, 1467-74	4	27
28	Translesion synthesis past the C8- and N2-deoxyguanosine adducts of the dietary mutagen 2-Amino-3-methylimidazo[4,5-f]quinoline in the NarI recognition sequence by prokaryotic DNA polymerases. <i>Chemical Research in Toxicology</i> , <b>2006</b> , 19, 1506-17	4	40
27	Orientation of the crotonaldehyde-derived N2-[3-Oxo-1(S)-methyl-propyl]-dG DNA adduct hinders interstrand cross-link formation in the 5'-CpG-3' sequence. <i>Chemical Research in Toxicology</i> , <b>2006</b> , 19, 1019-29	4	19
26	Spectroscopic characterization of interstrand carbinolamine cross-links formed in the 5'-CpG-3' sequence by the acrolein-derived gamma-OH-1,N2-propano-2'-deoxyguanosine DNA adduct. <i>Journal of the American Chemical Society</i> , <b>2005</b> , 127, 17686-96	16.4	36
25	Site specific synthesis and polymerase bypass of oligonucleotides containing a 6-hydroxy-3,5,6,7-tetrahydro-9H-imidazo[1,2-a]purin-9-one base, an intermediate in the formation of 1,N2-etheno-2'-deoxyguanosine. <i>Chemical Research in Toxicology</i> , <b>2005</b> , 18, 1701-14	4	26
24	Structure of an oligodeoxynucleotide containing a butadiene oxide-derived N1 beta-hydroxyalkyl deoxyinosine adduct in the human N-ras codon 61 sequence. <i>Biochemistry</i> , <b>2005</b> , 44, 3327-37	3.2	9
23	DNA adduct bypass polymerization by Sulfolobus solfataricus DNA polymerase Dpo4: analysis and crystal structures of multiple base pair substitution and frameshift products with the adduct 1,N2-ethenoguanine. <i>Journal of Biological Chemistry</i> , <b>2005</b> , 280, 29750-64	5.4	132
22	Synthesis of the N2-deoxyguanosine adduct of the potent dietary mutagen IQ. <i>Organic Letters</i> , <b>2004</b> , 6, 4985-8	6.2	19

## (1995-2004)

21	Induction of apoptosis in colorectal carcinoma cells treated with 4-hydroxy-2-nonenal and structurally related aldehydic products of lipid peroxidation. <i>Chemical Research in Toxicology</i> , <b>2004</b> , 17, 453-62	4	65
20	Coupling products of nucleosides with the glyoxal adduct of deoxyguanosine. <i>Chemical Research in Toxicology</i> , <b>2004</b> , 17, 1047-56	4	20
19	A novel synthesis of malondialdehyde adducts of deoxyguanosine, deoxyadenosine, and deoxycytidine. <i>Chemical Research in Toxicology</i> , <b>2004</b> , 17, 144-9	4	24
18	Site-specific synthesis and properties of oligonucleotides containing C8-deoxyguanosine adducts of the dietary mutagen IQ. <i>Journal of the American Chemical Society</i> , <b>2004</b> , 126, 11189-201	16.4	63
17	Site-specific mutagenicity of stereochemically defined 1,N2-deoxyguanosine adducts of trans-4-hydroxynonenal in mammalian cells. <i>Environmental and Molecular Mutagenesis</i> , <b>2003</b> , 42, 68-74	3.2	46
16	Stereoselective synthesis of the 1,N2-deoxyguanosine adducts of cinnamaldehyde. A stereocontrolled route to deoxyguanosine adducts of flunsaturated aldehydes. <i>Tetrahedron Letters</i> , <b>2003</b> , 44, 7513-7516	2	8
15	Site-specific synthesis and reactivity of oligonucleotides containing stereochemically defined 1,N2-deoxyguanosine adducts of the lipid peroxidation product trans-4-hydroxynonenal. <i>Journal of the American Chemical Society</i> , <b>2003</b> , 125, 5687-700	16.4	66
14	DNA interchain cross-links formed by acrolein and crotonaldehyde. <i>Journal of the American Chemical Society</i> , <b>2003</b> , 125, 50-61	16.4	167
13	DNA adducts of acrolein: site-specific synthesis of an oligodeoxynucleotide containing 6-hydroxy-5,6,7,8-tetrahydropyrimido[1,2-a]purin-10(3H)-one, an acrolein adduct of guanine. <i>Chemical Research in Toxicology</i> , <b>2002</b> , 15, 607-13	4	31
12	Absolute configuration and conformational stability of (+)-2,5-dimethylthiolane and (-)-2,5-dimethylsulfolane. <i>Journal of Organic Chemistry</i> , <b>2001</b> , 66, 3507-12	4.2	14
11	Synthesis of the C8-deoxyguanosine adduct of the food mutagen IQ. Organic Letters, 2001, 3, 565-8	6.2	63
10	Stereocontrolled syntheses of all four stereoisomeric 1,N2-deoxyguanosine adducts of the lipid peroxidation product trans-4-hydroxynonenal. <i>Organic Letters</i> , <b>2001</b> , 3, 3603-5	6.2	26
9	Stereocontrolled syntheses of deoxyribonucleosides via photoinduced electron-transfer deoxygenation of benzoyl-protected ribo- and arabinonucleosides. <i>Journal of Organic Chemistry</i> , <b>2000</b> , 65, 5969-85	4.2	32
8	X-ray crystal structures of conformationally biased flavin models. <i>Organic Letters</i> , <b>2000</b> , 2, 903-6	6.2	7
7	Regioselective synthesis of beta-N1- and beta-N3-alloxazine nucleosides. <i>Organic Letters</i> , <b>2000</b> , 2, 227-3	<b>3%</b> .2	28
6	Conformational Effects on Flavin Redox Chemistry. <i>Journal of Organic Chemistry</i> , <b>1997</b> , 62, 5244-5245	4.2	42
5	A Dne-PotIPhase Transfer Alkylation/Hydrolysis of o-Nitrotrifluoroacetanilides. A Convenient Route to N-ALKYL o-Phenylenediamines. <i>Synthetic Communications</i> , <b>1996</b> , 26, 4065-4080	1.7	17
4	Facile Addition of Dichloroketene to Acetylenes Mediated by Zinc and Ultrasound. <i>Synthetic Communications</i> , <b>1995</b> , 25, 2781-2789	1.7	20

3 9-Ethyl-3,6-Dimethylcarbazole (DMECZ)153-153

2	Synthesis of 2?-Deoxyribonucleosides: EB?,5?-Di-O-Benzoylthymidine162-162	1
1	Interstrand DNA Cross-Linking 1,N2-Deoxyguanosine Adducts Derived from ∰Unsaturated Aldehydes: Structure∄unction Relationships201-216	2