

Terrance M Arthur

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1933305/terrance-m-arthur-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

95
papers

3,879
citations

37
h-index

60
g-index

97
ext. papers

4,459
ext. citations

3.9
avg, IF

4.91
L-index

#	Paper	IF	Citations
95	Resistomes and microbiome of meat trimmings and colon content from culled cows raised in conventional and organic production systems.. <i>Animal Microbiome</i> , 2022 , 4, 21	4.1	0
94	Rates of evolutionary change of resident Escherichia coli O157:H7 differ within the same ecological niche.. <i>BMC Genomics</i> , 2022 , 23, 275	4.5	0
93	Validation of Additional Approaches and Applications for Using the Continuous and Manual Sampling Devices for Raw Beef Trim. <i>Journal of Food Protection</i> , 2021 , 84, 536-544	2.5	0
92	Antimicrobial Resistance in U.S. Retail Ground Beef with and without Label Claims Regarding Antibiotic Use. <i>Journal of Food Protection</i> , 2021 , 84, 827-842	2.5	3
91	A Comparative Quantitative Assessment of Human Exposure to Various Antimicrobial-Resistant Bacteria among U.S. Ground Beef Consumers. <i>Journal of Food Protection</i> , 2021 , 84, 736-759	2.5	0
90	Strain and host-cell dependent role of type-1 fimbriae in the adherence phenotype of super-shed Escherichia coli O157:H7. <i>International Journal of Medical Microbiology</i> , 2021 , 311, 151511	3.7	2
89	Nonfimbrial Adhesin Mutants Reveal Divergent O157:H7 Adherence Mechanisms on Human and Cattle Epithelial Cells. <i>International Journal of Microbiology</i> , 2021 , 2021, 8868151	3.6	1
88	Supershed O157:H7 Has Potential for Increased Persistence on the Rectoanal Junction Squamous Epithelial Cells and Antibiotic Resistance. <i>International Journal of Microbiology</i> , 2020 , 2020, 2368154	3.6	2
87	In-Feed Tylosin Phosphate Administration to Feedlot Cattle Minimally Affects Antimicrobial Resistance. <i>Journal of Food Protection</i> , 2020 , 350-364	2.5	5
86	Antimicrobial Resistance at Two U.S. Cull Cow Processing Establishments. <i>Journal of Food Protection</i> , 2020 , 83, 2216-2228	2.5	3
85	Metagenomic Characterization of the Microbiome and Resistome of Retail Ground Beef Products. <i>Frontiers in Microbiology</i> , 2020 , 11, 541972	5.7	6
84	Evaluation of two commercially-available vaccines on in the peripheral lymph nodes of experimentally-infected cattle 2020 , 8, 2515135520957760	2.6	0
83	No Change in Risk for Antibiotic-Resistant Salmonellosis from Beef, United States, 2002-2010. <i>Emerging Infectious Diseases</i> , 2020 , 26, 2108-2117	10.2	2
82	Food Service Pork Chops from Three U.S. Regions Harbor Similar Levels of Antimicrobial Resistance Regardless of Antibiotic Use Claims. <i>Journal of Food Protection</i> , 2019 , 82, 1667-1676	2.5	5
81	Cropland Amendment with Beef Cattle Manure Minimally Affects Antimicrobial Resistance. <i>Journal of Environmental Quality</i> , 2019 , 48, 1683-1693	3.4	5
80	Surface pH of Fresh Beef as a Parameter To Validate Effectiveness of Lactic Acid Treatment against Escherichia coli O157:H7 and Salmonella. <i>Journal of Food Protection</i> , 2018 , 81, 1126-1133	2.5	5
79	Similar Levels of Antimicrobial Resistance in U.S. Food Service Ground Beef Products with and without a "Raised without Antibiotics" Claim. <i>Journal of Food Protection</i> , 2018 , 81, 2007-2018	2.5	21

78	Novel Continuous and Manual Sampling Methods for Beef Trim Microbiological Testing. <i>Journal of Food Protection</i> , 2018 , 81, 1605-1613	2.5	2
77	Effects of In-Feed Chlortetracycline Prophylaxis in Beef Cattle on Antimicrobial Resistance Genes. <i>Foodborne Pathogens and Disease</i> , 2018 , 15, 689-697	3.8	11
76	Biofilm Formation, Antimicrobial Resistance, and Sanitizer Tolerance of <i>Salmonella enterica</i> Strains Isolated from Beef Trim. <i>Foodborne Pathogens and Disease</i> , 2017 , 14, 687-695	3.8	14
75	Comparative genomics of two super-shedder isolates of <i>Escherichia coli</i> O157:H7. <i>PLoS ONE</i> , 2017 , 12, e0182940	3.7	10
74	Rapid Detection and Classification of <i>Salmonella enterica</i> Shedding in Feedlot Cattle Utilizing the Roka Bioscience Atlas <i>Salmonella</i> Detection Assay for the Analysis of Rectoanal Mucosal Swabs. <i>Journal of Food Protection</i> , 2017 , 80, 1760-1767	2.5	3
73	Evaluation of Rectoanal Mucosal Swab Sampling for Molecular Detection of Enterohemorrhagic <i>Escherichia coli</i> in Beef Cattle. <i>Journal of Food Protection</i> , 2017 , 80, 661-667	2.5	4
72	Impact of "Raised without Antibiotics" Beef Cattle Production Practices on Occurrences of Antimicrobial Resistance. <i>Applied and Environmental Microbiology</i> , 2017 , 83,	4.8	46
71	Evaluation of Bacteriophage Application to Cattle in Lairage at Beef Processing Plants to Reduce <i>Escherichia coli</i> O157:H7 Prevalence on Hides and Carcasses. <i>Foodborne Pathogens and Disease</i> , 2017 , 14, 17-22	3.8	22
70	in Peripheral Lymph Nodes of Healthy Cattle at Slaughter. <i>Frontiers in Microbiology</i> , 2017 , 8, 2214	5.7	30
69	Effects of In-Feed Chlortetracycline Prophylaxis in Beef Cattle on Animal Health and Antimicrobial-Resistant <i>Escherichia coli</i> . <i>Applied and Environmental Microbiology</i> , 2016 , 82, 7197-7204	4.8	30
68	Diagnostic Accuracy of Rectoanal Mucosal Swab of Feedlot Cattle for Detection and Enumeration of <i>Salmonella enterica</i> . <i>Journal of Food Protection</i> , 2016 , 79, 531-7	2.5	4
67	<i>Escherichia coli</i> O157:H7 Strains Isolated from High-Event Period Beef Contamination Have Strong Biofilm-Forming Ability and Low Sanitizer Susceptibility, Which Are Associated with High pO157 Plasmid Copy Number. <i>Journal of Food Protection</i> , 2016 , 79, 1875-1883	2.5	13
66	Antimicrobial-Resistant Fecal Bacteria from Ceftiofur-Treated and Nonantimicrobial-Treated Comingled Beef Cows at a Cow-Calf Operation. <i>Microbial Drug Resistance</i> , 2016 , 22, 598-608	2.9	19
65	Effect of Direct-Fed Microbial Dosage on the Fecal Concentrations of Enterohemorrhagic <i>Escherichia coli</i> in Feedlot Cattle. <i>Foodborne Pathogens and Disease</i> , 2016 , 13, 190-5	3.8	2
64	Occurrence of Antimicrobial-Resistant <i>Escherichia coli</i> and <i>Salmonella enterica</i> in the Beef Cattle Production and Processing Continuum. <i>Applied and Environmental Microbiology</i> , 2015 , 81, 713-25	4.8	59
63	Efficacy of antimicrobial compounds on surface decontamination of seven Shiga toxin-producing <i>Escherichia coli</i> and <i>Salmonella</i> inoculated onto fresh beef. <i>Journal of Food Protection</i> , 2015 , 78, 503-10	2.5	11
62	Complete Genome Sequence of SS52, a Strain of <i>Escherichia coli</i> O157:H7 Recovered from Supershedder Cattle. <i>Genome Announcements</i> , 2015 , 3,		10
61	Comparative analysis of super-shedder strains of <i>Escherichia coli</i> O157:H7 reveals distinctive genomic features and a strongly aggregative adherent phenotype on bovine rectoanal junction squamous epithelial cells. <i>PLoS ONE</i> , 2015 , 10, e0116743	3.7	24

60	Antimicrobial-Resistant Bacterial Populations and Antimicrobial Resistance Genes Obtained from Environments Impacted by Livestock and Municipal Waste. <i>PLoS ONE</i> , 2015 , 10, e0132586	3.7	68
59	Biofilm formation and sanitizer resistance of Escherichia coli O157:H7 strains isolated from "high event period" meat contamination. <i>Journal of Food Protection</i> , 2014 , 77, 1982-7	2.5	38
58	Characterization of Escherichia coli O157:H7 strains from contaminated raw beef trim during "high event periods". <i>Applied and Environmental Microbiology</i> , 2014 , 80, 506-14	4.8	19
57	The physiologic state of Escherichia coli O157:H7 does not affect its detection in two commercial real-time PCR-based tests. <i>Food Microbiology</i> , 2013 , 33, 205-12	6	2
56	Chromogenic agar medium for detection and isolation of Escherichia coli serogroups O26, O45, O103, O111, O121, and O145 from fresh beef and cattle feces. <i>Journal of Food Protection</i> , 2013 , 76, 192-95	2.5	46
55	Isolation and characterization of Clostridium difficile associated with beef cattle and commercially produced ground beef. <i>Journal of Food Protection</i> , 2013 , 76, 256-64	2.5	12
54	Disinfectant and antibiotic susceptibility profiles of Escherichia coli O157:H7 strains from cattle carcasses, feces, and hides and ground beef from the United States. <i>Journal of Food Protection</i> , 2013 , 76, 6-17	2.5	18
53	Characterization of Escherichia coli O157:H7 strains isolated from supershedding cattle. <i>Applied and Environmental Microbiology</i> , 2013 , 79, 4294-303	4.8	37
52	Cross-sectional study examining Salmonella enterica carriage in subiliac lymph nodes of cull and feedlot cattle at harvest. <i>Foodborne Pathogens and Disease</i> , 2013 , 10, 368-74	3.8	69
51	Microbiological analysis of bovine lymph nodes for the detection of Salmonella enterica. <i>Journal of Food Protection</i> , 2012 , 75, 854-8	2.5	43
50	Evaluation of commonly used antimicrobial interventions for fresh beef inoculated with Shiga toxin-producing Escherichia coli serotypes O26, O45, O103, O111, O121, O145, and O157:H7. <i>Journal of Food Protection</i> , 2012 , 75, 1207-12	2.5	59
49	Detection of Escherichia coli O157:H7 and Salmonella enterica in air and droplets at three U.S. commercial beef processing plants. <i>Journal of Food Protection</i> , 2012 , 75, 2213-8	2.5	22
48	The epitope for the polyol-responsive monoclonal antibody 8RB13 is in the flap-domain of the beta-subunit of bacterial RNA polymerase and can be used as an epitope tag for immunoaffinity chromatography. <i>Protein Expression and Purification</i> , 2011 , 77, 26-33	2	11
47	Diversity of multidrug-resistant salmonella enterica strains associated with cattle at harvest in the United States. <i>Applied and Environmental Microbiology</i> , 2011 , 77, 1783-96	4.8	53
46	Survival of Escherichia coli O157:H7 on cattle hides. <i>Applied and Environmental Microbiology</i> , 2011 , 77, 3002-8	4.8	31
45	Evaluation of a direct-fed microbial product effect on the prevalence and load of Escherichia coli O157:H7 in feedlot cattle. <i>Journal of Food Protection</i> , 2010 , 73, 366-71	2.5	25
44	Soil versus Pond Ash Surfacing of Feedlot Pens: Occurrence of Escherichia coli O157:H7 in Cattle and Persistence in Manure. <i>Journal of Food Protection</i> , 2010 , 73, 1269-77	2.5	16
43	Super shedding of Escherichia coli O157:H7 by cattle and the impact on beef carcass contamination. <i>Meat Science</i> , 2010 , 86, 32-7	6.4	105

42	Prevalence and enumeration of Escherichia coli O157:H7 and Salmonella in U.S. abattoirs that process fewer than 1000 head of cattle per day. <i>Journal of Food Protection</i> , 2009 , 72, 1272-8	2.5	35
41	Prevalence rates of Escherichia coli O157:H7 and Salmonella at different sampling sites on cattle hides at a feedlot and processing plant. <i>Journal of Food Protection</i> , 2009 , 72, 1267-71	2.5	24
40	Longitudinal study of Escherichia coli O157:H7 in a beef cattle feedlot and role of high-level shedders in hide contamination. <i>Applied and Environmental Microbiology</i> , 2009 , 75, 6515-23	4.8	102
39	Impacts of individual animal response to heat and handling stresses on Escherichia coli and E. coli O157:H7 fecal shedding by feedlot cattle. <i>Foodborne Pathogens and Disease</i> , 2009 , 6, 855-64	3.8	11
38	Effectiveness of 1,3-dibromo-5,5 dimethylhydantoin on reduction of Escherichia coli O157:H7- and Salmonella-inoculated fresh meat. <i>Journal of Food Protection</i> , 2009 , 72, 151-6	2.5	19
37	Salmonella and Escherichia coli O157:H7 contamination on hides and carcasses of cull cattle presented for slaughter in the United States: an evaluation of prevalence and bacterial loads by immunomagnetic separation and direct plating methods. <i>Applied and Environmental Microbiology</i> , 2009 , 75, 4000-07	4.8	115
36	Source tracking of Escherichia coli O157:H7 and Salmonella contamination in the lairage environment at commercial U.S. beef processing plants and identification of an effective intervention. <i>Journal of Food Protection</i> , 2008 , 71, 1752-60	2.5	74
35	Prevalence and characterization of Salmonella in bovine lymph nodes potentially destined for use in ground beef. <i>Journal of Food Protection</i> , 2008 , 71, 1685-8	2.5	65
34	Comparison of effects of antimicrobial interventions on multidrug-resistant Salmonella, susceptible Salmonella, and Escherichia coli O157:H7. <i>Journal of Food Protection</i> , 2008 , 71, 2177-81	2.5	27
33	Evaluation of various antimicrobial interventions for the reduction of Escherichia coli O157:H7 on bovine heads during processing. <i>Journal of Food Protection</i> , 2008 , 71, 621-4	2.5	39
32	Enumeration of Salmonella from poultry carcass rinses via direct plating methods. <i>Letters in Applied Microbiology</i> , 2008 , 46, 186-91	2.9	21
31	Transportation and lairage environment effects on prevalence, numbers, and diversity of Escherichia coli O157:H7 on hides and carcasses of beef cattle at processing. <i>Journal of Food Protection</i> , 2007 , 70, 280-6	2.5	112
30	Listeria prevalence and Listeria monocytogenes serovar diversity at cull cow and bull processing plants in the United States. <i>Journal of Food Protection</i> , 2007 , 70, 2578-82	2.5	40
29	Microbiological characterization of imported and domestic boneless beef trim used for ground beef. <i>Journal of Food Protection</i> , 2007 , 70, 440-9	2.5	47
28	Microbiological characterization of lamb carcasses at commercial processing plants in the United States. <i>Journal of Food Protection</i> , 2007 , 70, 1811-9	2.5	30
27	Effects of a minimal hide wash cabinet on the levels and prevalence of Escherichia coli O157:H7 and Salmonella on the hides of beef cattle at slaughter. <i>Journal of Food Protection</i> , 2007 , 70, 1076-9	2.5	50
26	Enumeration of Salmonella and Escherichia coli O157:H7 in ground beef, cattle carcass, hide and faecal samples using direct plating methods. <i>Journal of Applied Microbiology</i> , 2007 , 103, 1657-68	4.7	87
25	Interventions to reduce/eliminate Escherichia coli O157:H7 in ground beef. <i>Meat Science</i> , 2007 , 77, 90-6	6.4	41

24	Comparison of the molecular genotypes of Escherichia coli O157:H7 from the hides of beef cattle in different regions of North America. <i>Journal of Food Protection</i> , 2007 , 70, 1622-6	2.5	20
23	Treatments using hot water instead of lactic acid reduce levels of aerobic bacteria and Enterobacteriaceae and reduce the prevalence of Escherichia coli O157:H7 on previsceration beef carcasses. <i>Journal of Food Protection</i> , 2006 , 69, 1808-13	2.5	78
22	Improvement of immunomagnetic separation for Escherichia coli O157:H7 detection by the PickPen magnetic particle separation device. <i>Journal of Food Protection</i> , 2006 , 69, 2870-4	2.5	30
21	Evaluation of Escherichia coli O157:H7 growth media for use in test-and-hold procedures for ground beef processing. <i>Journal of Food Protection</i> , 2006 , 69, 1007-11	2.5	25
20	Post-harvest interventions to reduce/eliminate pathogens in beef. <i>Meat Science</i> , 2005 , 71, 79-91	6.4	171
19	Methods for recovering Escherichia coli O157:H7 from cattle fecal, hide, and carcass samples: sensitivity and improvements. <i>Journal of Food Protection</i> , 2005 , 68, 2264-8	2.5	36
18	Effects of low-dose, low-penetration electron beam irradiation of chilled beef carcass surface cuts on Escherichia coli O157:H7 and meat quality. <i>Journal of Food Protection</i> , 2005 , 68, 666-72	2.5	41
17	Evaluation of culture- and PCR-based detection methods for Escherichia coli O157:H7 in inoculated ground beef. <i>Journal of Food Protection</i> , 2005 , 68, 1566-74	2.5	38
16	Prevalence of Escherichia coli O157 and levels of aerobic bacteria and Enterobacteriaceae are reduced when hides are washed and treated with cetylpyridinium chloride at a commercial beef processing plant. <i>Journal of Food Protection</i> , 2004 , 67, 646-50	2.5	85
15	Protocol for evaluating the efficacy of cetylpyridinium chloride as a beef hide intervention. <i>Journal of Food Protection</i> , 2004 , 67, 303-9	2.5	29
14	Prevalence of Escherichia coli O157:H7, Listeria monocytogenes, and Salmonella in two geographically distant commercial beef processing plants in the United States. <i>Journal of Food Protection</i> , 2004 , 67, 295-302	2.5	105
13	Escherichia coli O157 prevalence and enumeration of aerobic bacteria, Enterobacteriaceae, and Escherichia coli O157 at various steps in commercial beef processing plants. <i>Journal of Food Protection</i> , 2004 , 67, 658-65	2.5	196
12	Characterization of O157:H7 and other Escherichia coli isolates recovered from cattle hides, feces, and carcasses. <i>Journal of Food Protection</i> , 2004 , 67, 993-8	2.5	22
11	Seasonal prevalence of Shiga toxin-producing Escherichia coli, including O157:H7 and non-O157 serotypes, and Salmonella in commercial beef processing plants. <i>Journal of Food Protection</i> , 2003 , 66, 1978-86	2.5	351
10	Development of an epitope tag for the gentle purification of proteins by immunoaffinity chromatography: application to epitope-tagged green fluorescent protein. <i>Analytical Biochemistry</i> , 2003 , 323, 171-9	3.1	26
9	Development of methods for the recovery of Escherichia coli O157:H7 and Salmonella from beef carcass sponge samples and bovine fecal and hide samples. <i>Journal of Food Protection</i> , 2002 , 65, 1527-34	2.5	83
8	Gas formation in ground beef chubs due to Hafnia alvei is reduced by multiple applications of antimicrobial interventions to artificially inoculated beef trim stock. <i>Journal of Food Protection</i> , 2002 , 65, 1651-5	2.5	7
7	Prevalence and characterization of non-O157 Shiga toxin-producing Escherichia coli on carcasses in commercial beef cattle processing plants. <i>Applied and Environmental Microbiology</i> , 2002 , 68, 4847-52	4.8	105

6	Genotypic analyses of Escherichia coli O157:H7 and O157 nonmotile isolates recovered from beef cattle and carcasses at processing plants in the Midwestern states of the United States. <i>Applied and Environmental Microbiology</i> , 2001 , 67, 3810-8	4.8	93
5	Binding of the initiation factor sigma(70) to core RNA polymerase is a multistep process. <i>Molecular Cell</i> , 2001 , 8, 21-31	17.6	58
4	A coiled-coil from the RNA polymerase betaTsubunit allosterically induces selective nontemplate strand binding by sigma(70). <i>Cell</i> , 2001 , 105, 935-44	56.2	84
3	Mutational analysis of beta T260-309, a sigma 70 binding site located on Escherichia coli core RNA polymerase. <i>Journal of Biological Chemistry</i> , 2000 , 275, 23113-9	5.4	64
2	Mapping protein-protein interaction domains using ordered fragment ladder far-western analysis of hexahistidine-tagged fusion proteins. <i>Methods in Enzymology</i> , 2000 , 328, 141-57	1.7	39
1	Localization of a sigma70 binding site on the N terminus of the Escherichia coli RNA polymerase betaTsubunit. <i>Journal of Biological Chemistry</i> , 1998 , 273, 31381-7	5.4	106