

Paulo Bonifácio

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1934141/publications.pdf>

Version: 2024-02-01

32
papers

411
citations

759233

12
h-index

839539

18
g-index

41
all docs

41
docs citations

41
times ranked

510
citing authors

#	ARTICLE	IF	CITATIONS
1	The Community Structure of Deep-Sea Macrofauna Associated with Polymetallic Nodules in the Eastern Part of the Clarion-Clipperton Fracture Zone. <i>Frontiers in Marine Science</i> , 2017, 4, .	2.5	64
2	Alpha and beta diversity patterns of polychaete assemblages across the nodule province of the eastern Clarion-Clipperton Fracture Zone (equatorial Pacific). <i>Biogeosciences</i> , 2020, 17, 865-886.	3.3	38
3	Patterns of Macrofaunal Biodiversity Across the Clarion-Clipperton Zone: An Area Targeted for Seabed Mining. <i>Frontiers in Marine Science</i> , 2021, 8, .	2.5	33
4	New genera and species from the Equatorial Pacific provide phylogenetic insights into deep-sea Polynoidae (Annelida). <i>Zoological Journal of the Linnean Society</i> , 2019, 185, 555-635.	2.3	32
5	Spatiotemporal changes in surface sediment characteristics and benthic macrofauna composition off the Rhône River in relation to its hydrological regime. <i>Estuarine, Coastal and Shelf Science</i> , 2014, 151, 196-209.	2.1	24
6	A new species of the <i>Marphysa sanguinea</i> complex from French waters (Bay of Biscay, NE Atlantic) (Annelida, Eunicidae). <i>ZooKeys</i> , 2017, 716, 1-17.	1.1	24
7	High diversity and pan-oceanic distribution of deep-sea polychaetes: <i>Prionospio</i> and <i>Aurospio</i> (Annelida: Spionidae) in the Atlantic and Pacific Ocean. <i>Organisms Diversity and Evolution</i> , 2020, 20, 171-187.	1.6	16
8	Toward a reliable assessment of potential ecological impacts of deep-sea polymetallic nodule mining on abyssal infauna. <i>Limnology and Oceanography: Methods</i> , 2021, 19, 626-650.	2.0	16
9	First record of <i>Naineris setosa</i> (Verrill, 1900) (Annelida: Polychaeta: Orbiniidae) in the Western Mediterranean Sea. <i>BioInvasions Records</i> , 2014, 3, 83-88.	1.1	15
10	Recent discovery of <i>Paranthura japonica</i> Richardson, 1909 (Crustacea: Isopoda: Paranthuridae) in		

#	ARTICLE	IF	CITATIONS
19	<i>Anobothrus amourouxi</i> sp. nov., a new species of Ampharetidae (Polychaeta) from the Capbreton Canyon (Bay of Biscay, NE Atlantic Ocean). Journal of the Marine Biological Association of the United Kingdom, 2015, 95, 961-969.	0.8	6
20	Long-term (1998–2010) large-scale comparison of the ecological quality status of gulf of lions (NW) Tj ETQq0 0 0 rgBT /Overlock 10 T	3.0	6
21	<i>Malmgrenia louiseae</i> sp. nov., a new scale worm species (Polychaeta: Polynoidae) from southern Europe with a key to European <i>Malmgrenia</i> species. Journal of the Marine Biological Association of the United Kingdom, 2015, 95, 947-952.	0.8	5
22	Diversity of Deep-Sea Scale-Worms (Annelida, Polynoidae) in the Clarion-Clipperton Fracture Zone. Frontiers in Marine Science, 2021, 8, .	2.5	5
23	Infestation of <i>Crassostrea</i> cf. <i>brasiliensis</i> by boring-polychaete polydorids in estuaries from Northeastern Brazil. Journal of Environmental Analysis and Progress, 2017, 2, 16-22.	0.2	5
24	A new species and phylogenetic insights in <i>Hesiospina</i> (Annelida, Hesionidae). Zootaxa, 2018, 4441, 59-75.	0.5	4
25	Climate-driven changes in macrobenthic communities in the Mediterranean Sea: A 10-year study in the Bay of Banyuls-sur-Mer. Ecology and Evolution, 2019, 9, 10483-10498.	1.9	4
26	New records of <i>Spio</i> symphyta and <i>Spio martinensis</i> (Polychaeta: Canalipalpata: Spionidae) from Arcachon Bay (France), NE Atlantic. Marine Biodiversity, 2015, 45, 77-86.	1.0	3
27	A new species of <i>Pista</i> Malmgren, 1866 (Polychaeta, Terebellidae) from the north-western Mediterranean Sea. ZooKeys, 2019, 838, 71-84.	1.1	3
28	A new species of <i>Thorecta</i> (Porifera: Demospongiae) from the western Atlantic, with remarks on the taxonomy of the genus. Journal of the Marine Biological Association of the United Kingdom, 2010, 90, 775-782.	0.8	2
29	Shallow-water polychaete assemblages in the northwestern Mediterranean Sea and its possible use in the evaluation of good environmental state. Memoirs of Museum Victoria, 2014, 71, 289-301.	0.6	2
30	<p class="ZootaxaTitle">New insights in the biogeographical distributions of two Spionidae (Annelida) from the NE Atlantic and Mediterranean French coasts</p>. Zoosymposia, 2020, 19, 173-184.	0.3	2
31	Molecular analysis and new records of the invasive polychaete <i>Boccardia proboscidea</i> (Annelida:) Tj ETQq1 1 0.784314 rgBT /Overlock 1	1.6	1
32	<i>Anobothrus amourouxi</i> sp. nov., a new species of Ampharetidae (Polychaeta) from the Capbreton Canyon (Bay of Biscay, NE Atlantic Ocean) – ERRATUM. Journal of the Marine Biological Association of the United Kingdom, 2015, 95, 971-971.	0.8	0