## Pedro Anda

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1955689/publications.pdf

Version: 2024-02-01

206112 172457 2,438 57 29 48 citations h-index g-index papers 64 64 64 2654 all docs docs citations times ranked citing authors

| #  | Article  | IF   | CITATIONS |
|----|--|------|-----------|
| 1  | Molecular Survey of <i>Rickettsia</i> spp., <i>Anaplasma</i> spp., <i>Ehrlichia</i> spp., <i>Bartonella</i> spp., and <i>Borrelia</i> spp. in Fleas and Lice in Ethiopia. Vector-Borne and Zoonotic Diseases, 2020, 20, 10-14.                   | 1.5  | 8         |
| 2  | Density-Dependent Prevalence of <i>Francisella tularensis</i> in Fluctuating Vole Populations, Northwestern Spain. Emerging Infectious Diseases, 2017, 23, 1377-1379.  | 4.3  | 30        |
| 3  | Irruptive mammal host populations shape tularemia epidemiology. PLoS Pathogens, 2017, 13, e1006622.  | 4.7  | 40        |
| 4  | Genotypes of <i>Coxiella burnetii</i> in wildlife: disentangling the molecular epidemiology of a multiâ€host pathogen. Environmental Microbiology Reports, 2016, 8, 708-714.   | 2.4  | 11        |
| 5  | Unique human immune signature of Ebola virus disease in Guinea. Nature, 2016, 533, 100-104.  | 27.8 | 170       |
| 6  | Towards Development of Improved Serodiagnostics for Tularemia by Use of Francisella tularensis Proteome Microarrays. Journal of Clinical Microbiology, 2016, 54, 1755-1765.  | 3.9  | 13        |
| 7  | Long-range dispersal moved Francisella tularensis into Western Europe from the East. Microbial<br>Genomics, 2016, 2, e000100.  | 2.0  | 32        |
| 8  | <i>Coxiella burnetii</i> total immunoglobulin G, phase I and phase II immunoglobulin G antibodies, and bacterial shedding in young dams in persistently infected dairy herds. Journal of Veterinary Diagnostic Investigation, 2015, 27, 167-176. | 1.1  | 7         |
| 9  | Tularemia Outbreaks and Common Vole ( <i>Microtus arvalis</i> ) Irruptive Population Dynamics in Northwestern Spain, 1997–2014. Vector-Borne and Zoonotic Diseases, 2015, 15, 568-570.   | 1.5  | 30        |
| 10 | The importance of lizards and small mammals as reservoirs for <scp><i>B</i></scp> <i>orrelia lusitaniae</i> in <scp>P</scp> ortugal. Environmental Microbiology Reports, 2015, 7, 188-193.   | 2.4  | 15        |
| 11 | Identification of essential outstanding questions for an adequate European laboratory response to Ebolavirus Zaire West Africa 2014. Journal of Clinical Virology, 2015, 62, 124-134.  | 3.1  | 27        |
| 12 | Molecular characterization of Rickettsia massiliae and Anaplasma platys infecting Rhipicephalus sanguineus ticks and domestic dogs, Buenos Aires (Argentina). Ticks and Tick-borne Diseases, 2014, 5, 484-488.                                   | 2.7  | 42        |
| 13 | Melioidosis in Traveler from Africa to Spain. Emerging Infectious Diseases, 2013, 19, 1656-1659.   | 4.3  | 26        |
| 14 | Human Infection withRickettsia sibirica mongolitimonae, Spain, 2007–2011. Emerging Infectious Diseases, 2013, 19, 267-269.   | 4.3  | 40        |
| 15 | Distribution of Bartonella henselae Variants in Patients, Reservoir Hosts and Vectors in Spain. PLoS<br>ONE, 2013, 8, e68248.  | 2.5  | 27        |
| 16 | Fatal Bacillary Angiomatosis Mimicking An Infiltrative Vascular Tumour in the Immune Restoration Phase of An HIV-Infected Patient. Antiviral Therapy, 2012, 17, 405-407.   | 1.0  | 5         |
| 17 | Presence of Bartonella Species in Wild Carnivores of Northern Spain. Applied and Environmental<br>Microbiology, 2012, 78, 885-888.   | 3.1  | 39        |
| 18 | Molecular method for the characterization of Coxiella burnetii from clinical and environmental samples: variability of genotypes in Spain. BMC Microbiology, 2012, 12, 91.   | 3.3  | 28        |

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Recombinase Polymerase Amplification Assay for Rapid Detection of Francisella tularensis. Journal of Clinical Microbiology, 2012, 50, 2234-2238.   | 3.9 | 144       |
| 20 | Differences in Questing Tick Species Distribution Between Atlantic and Continental Climate Regions in Spain. Journal of Medical Entomology, 2011, 48, 13-19.   | 1.8 | 46        |
| 21 | Infeccion por Rickettsia sibirica subsp. monogolitimonae. Piel, 2011, 26, 224-226.   | 0.0 | 4         |
| 22 | Melioidosis Imported from West Africa to Europe. American Journal of Tropical Medicine and Hygiene, 2011, 85, 282-284.   | 1.4 | 26        |
| 23 | A possible novel Francisellagenomic species isolated from blood and urine of a patient with severe illness. Clinical Microbiology and Infection, 2010, 16, 1026-1030.  | 6.0 | 12        |
| 24 | Description of Francisella hispaniensis sp. nov., isolated from human blood, reclassification of Francisella novicida (Larson et al. 1955) Olsufiev et al. 1959 as Francisella tularensis subsp. novicida comb. nov. and emended description of the genus Francisella. International Journal of Systematic and Evolutionary Microbiology, 2010, 60, 1887-1896. | 1.7 | 101       |
| 25 | Phylogenetic Analysis of a Virulent <i>Borrelia</i> Species Isolated from Patients with Relapsing Fever. Journal of Clinical Microbiology, 2010, 48, 2484-2489.  | 3.9 | 41        |
| 26 | Variability of <i>Bartonella</i> Genotypes among Small Mammals in Spain. Applied and Environmental Microbiology, 2010, 76, 8062-8070.  | 3.1 | 47        |
| 27 | Objections to the transfer of Francisella novicida to the subspecies rank of Francisella tularensis $\hat{a} \in \mathbb{C}^*$ response to Johansson et al International Journal of Systematic and Evolutionary Microbiology, 2010, 60, 1718-1720.   | 1.7 | 36        |
| 28 | Objections to the transfer of Francisella novicida to the subspecies rank of Francisella tularensis. International Journal of Systematic and Evolutionary Microbiology, 2010, 60, 1717-1718.   | 1.7 | 62        |
| 29 | Detection ofCoxiella burnetiiin Ticks Collected from Central Spain. Vector-Borne and Zoonotic Diseases, 2009, 9, 465-468.  | 1.5 | 42        |
| 30 | Complement factor H binding by different Lyme disease and relapsing fever Borrelia in animals and human. BMC Research Notes, 2009, 2, 134.   | 1.4 | 44        |
| 31 | Anaplasma phagocytophilum is not an aetiological agent of fever of intermediate duration in Gran<br>Canaria (Spain). Clinical Microbiology and Infection, 2009, 15, 6-7.   | 6.0 | O         |
| 32 | Tick-Borne Zoonotic Bacteria in Ticks Collected from Central Spain. American Journal of Tropical Medicine and Hygiene, 2009, 81, 67-74.  | 1.4 | 80        |
| 33 | Tick-borne zoonotic bacteria in ticks collected from central Spain. American Journal of Tropical Medicine and Hygiene, 2009, 81, 67-74.  | 1.4 | 28        |
| 34 | Prevalence of Tick-Borne Zoonotic Bacteria in Questing Adult Ticks from Northern Spain. Vector-Borne and Zoonotic Diseases, 2008, 8, 829-836.  | 1.5 | 67        |
| 35 | Molecular Method for Bartonella Species Identification in Clinical and Environmental Samples.<br>Journal of Clinical Microbiology, 2008, 46, 776-779.  | 3.9 | 37        |
| 36 | Molecular Method for Discrimination between <i>Francisella tularensis</i> and <i>Francisella</i> -Like Endosymbionts. Journal of Clinical Microbiology, 2008, 46, 3139-3143.   | 3.9 | 33        |

| #  | Article  | IF          | CITATIONS |
|----|--|-------------|-----------|
| 37 | Tick-Borne Zoonotic Bacteria in Wild and Domestic Small Mammals in Northern Spain. Applied and Environmental Microbiology, 2007, 73, 6166-6171.  | 3.1         | 73        |
| 38 | <i>Rickettsia monacensis</i> i>and Human Disease, Spain. Emerging Infectious Diseases, 2007, 13, 1405-1407.  | 4.3         | 188       |
| 39 | <i>Francisella tularensis, </i> Portugal. Emerging Infectious Diseases, 2007, 13, 666-667.   | 4.3         | 20        |
| 40 | Risk factors associated with ixodid tick species distributions in the Basque region in Spain. Medical and Veterinary Entomology, 2006, 20, 177-188.  | 1.5         | 62        |
| 41 | Molecular Method for Identification of Rickettsia Species in Clinical and Environmental Samples.<br>Journal of Clinical Microbiology, 2006, 44, 4572-4576.                                       | 3.9         | 75        |
| 42 | Identification of a New Borrelia Species among Small Mammals in Areas of Northern Spain Where Lyme Disease Is Endemic. Applied and Environmental Microbiology, 2005, 71, 1336-1345.              | 3.1         | 38        |
| 43 | Dermacentor-borne necrosis erythema and lymphadenopathy: clinical and epidemiological features of a new tick-borne disease. Clinical Microbiology and Infection, 2004, 10, 327-331.              | 6.0         | 90        |
| 44 | Hallazgo infrecuente en sangre perifà ©rica. Enfermedades Infecciosas Y MicrobiologÃa ClÃnica, 2003, 21, 59-60.  | 0.5         | 1         |
| 45 | Distribution of <i>Borrelia burgdorferi</i> sensu lato in <i>lxodes ricinus</i> (Acari: lxodidae) Ticks from the Basque Country, Spain. Journal of Medical Entomology, 2002, 39, 177-184.        | 1.8         | 35        |
| 46 | In Vitro Culture of Borrelia garinii Results in Loss of Flagella and Decreased Invasiveness. Infection and Immunity, 2002, 70, 4851-4858.  | 2.2         | 28        |
| 47 | Waterborne Outbreak of Tularemia Associated with Crayfish Fishing. Emerging Infectious Diseases, 2001, 7, 575-582.   | 4.3         | 61        |
| 48 | Molecular and Pathogenic Characterization of <i>Borrelia burgdorferi</i> Sensu Lato Isolates from Spain. Journal of Clinical Microbiology, 2000, 38, 4026-4033.                                  | 3.9         | 60        |
| 49 | A European multi-centre comparison of immunoblot in the serodiagnosis of Lyme borreliosis.<br>Zentralblatt Fur Bakteriologie: International Journal of Medical Microbiology, 1999, 289, 678-680. | 0.5         | 1         |
| 50 | Q fever in pregnancy: case report after a 2-year follow-up. Journal of Infection, 1998, 37, 79-81.   | 3.3         | 11        |
| 51 | Use of the C3H/He Lyme disease mouse model for the recovery of a Spanish isolate of Borrelia garinii from erythema migrans lesions. Research in Microbiology, 1998, 149, 39-46.                  | 2.1         | 28        |
| 52 | Disparity Between Serological Reactivity to <i>Borrelia burgdorferi</i> a Highâ€Risk Group. Clinical Infectious Diseases, 1998, 27, 1210-1213.   | <b>5.</b> 8 | 22        |
| 53 | A Mouse Model of <i>Borrelia</i> Meningitis after Intradermal Injection. Journal of Infectious Diseases, 1997, 175, 1243-1245.   | 4.0         | 36        |
| 54 | Glycolytic enzyme operon of Borrelia burgdorferi: characterization and evolutionary implications. Gene, 1997, 188, 221-228.  | 2.2         | 23        |

## PEDRO ANDA

| #  | Article   | IF   | CITATIONS |
|----|---|------|-----------|
| 55 | A new Borrelia species isolated from patients with relapsing fever in Spain. Lancet, The, 1996, 348, 162-165.                                       | 13.7 | 80        |
| 56 | Study of C. burnetii human and animal seroprevalence in a rural population in Madrid community. European Journal of Epidemiology, 1989, 5, 444-446. | 5.7  | 18        |
| 57 | LEGIONNAIRES' DISEASE IN SPAIN. Lancet, The, 1983, 321, 759.  | 13.7 | 1         |