

# Signe Modvig

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1970642/publications.pdf>

Version: 2024-02-01

13  
papers

769  
citations

1039880

9  
h-index

1281743

11  
g-index

14  
all docs

14  
docs citations

14  
times ranked

1890  
citing authors

#	ARTICLE	IF	CITATIONS
1	High CD34 surface expression in BCP-ALL predicts poor induction therapy response and is associated with altered expression of genes related to cell migration and adhesion. <i>Molecular Oncology</i> , 2022, 16, 2015-2030.	2.1	5
2	Flow cytometric analysis of cerebrospinal fluid improves detection of leukaemic blasts in infants with acute lymphoblastic leukaemia. <i>British Journal of Haematology</i> , 2021, 195, 119-122.	1.2	4
3	Value of flow cytometry for MRD-based relapse prediction in B-cell precursor ALL in a multicenter setting. <i>Leukemia</i> , 2020, 35, 1894-1906.	3.3	23
4	Integrin-Mediated Adhesion and Chemoresistance of Acute Lymphoblastic Leukemia Cells Residing in the Bone Marrow or the Central Nervous System. <i>Frontiers in Oncology</i> , 2020, 10, 775.	1.3	16
5	A comprehensive clinical study of integrins in acute lymphoblastic leukemia indicates a role of $\alpha 6/\text{CD}49\text{f}$ in persistent minimal residual disease and $\alpha 5$ in the colonization of cerebrospinal fluid. <i>Leukemia and Lymphoma</i> , 2020, 61, 1714-1718.	0.6	13
6	Diagnostic Value of Cerebrospinal Fluid Neurofilament Light Protein in Neurology. <i>JAMA Neurology</i> , 2019, 76, 1035.	4.5	455
7	Minimal residual disease quantification by flow cytometry provides reliable risk stratification in T-cell acute lymphoblastic leukemia. <i>Leukemia</i> , 2019, 33, 1324-1336.	3.3	42
8	Value of Flow Cytometry for MRD-Based Relapse Prediction in B-Cell Precursor Acute Lymphoblastic Leukemia in a Multi-Center Setting. <i>Blood</i> , 2019, 134, 2755-2755.	0.6	0
9	Increased prevalence of lymphoid tissue inducer cells in the cerebrospinal fluid of patients with early multiple sclerosis. <i>Multiple Sclerosis Journal</i> , 2016, 22, 1013-1020.	1.4	20
10	Cerebrospinal fluid neurofilament light chain levels predict visual outcome after optic neuritis. <i>Multiple Sclerosis Journal</i> , 2016, 22, 590-598.	1.4	39
11	An Immature Immunophenotype (CD34pos, CD38dim, nTdTdim) on Malignant B-Cell Precursor Blasts at Diagnosis Predicts High Minimal Residual Disease in BCP-ALL. <i>Blood</i> , 2016, 128, 1728-1728.	0.6	0
12	Cerebrospinal fluid levels of chitinase 3-like 1 and neurofilament light chain predict multiple sclerosis development and disability after optic neuritis. <i>Multiple Sclerosis Journal</i> , 2015, 21, 1761-1770.	1.4	97
13	Relationship between Cerebrospinal Fluid Biomarkers for Inflammation, Demyelination and Neurodegeneration in Acute Optic Neuritis. <i>PLoS ONE</i> , 2013, 8, e77163.	1.1	55