

Geesje M Dallinga-Thie

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/197161/geesje-m-dallinga-thie-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

148
papers

8,027
citations

42
h-index

87
g-index

152
ext. papers

9,257
ext. citations

6.5
avg, IF

5.32
L-index

#	Paper	IF	Citations
148	The Role of Heparan Sulfate and Neuropilin 2 in VEGFA Signaling in Human Endothelial Tip Cells and Non-Tip Cells during Angiogenesis In Vitro. <i>Cells</i> , 2021 , 10,	7.9	5
147	ANGPTL3 Inhibition With Evinacumab Results in Faster Clearance of IDL and LDL apoB in Patients With Homozygous Familial Hypercholesterolemia-Brief Report. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2021 , 41, 1753-1759	9.4	18
146	Parenteral nutrition impairs plasma bile acid and gut hormone responses to mixed meal testing in lean healthy men. <i>Clinical Nutrition</i> , 2021 , 40, 1013-1021	5.9	2
145	Targeting apoC-III and ANGPTL3 in the treatment of hypertriglyceridemia. <i>Expert Review of Cardiovascular Therapy</i> , 2020 , 18, 355-361	2.5	9
144	Taking One Step Back in Familial Hypercholesterolemia: Does Not Alter Plasma LDL (Low-Density Lipoprotein) Cholesterol in Mice and Humans. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2020 , 40, 973-985	9.4	22
143	Oral butyrate does not affect innate immunity and islet autoimmunity in individuals with longstanding type 1 diabetes: a randomised controlled trial. <i>Diabetologia</i> , 2020 , 63, 597-610	10.3	26
142	Reduced CETP glycosylation and activity in patients with homozygous B4GALT1 mutations. <i>Journal of Inherited Metabolic Disease</i> , 2020 , 43, 611-617	5.4	7
141	Genetic variants in SUSP2 are associated with the risk of ischemic heart disease. <i>Journal of Clinical Lipidology</i> , 2020 , 14, 470-481	4.9	1
140	Diabetic Nephropathy Alters the Distribution of Circulating Angiogenic MicroRNAs Among Extracellular Vesicles, HDL, and Ago-2. <i>Diabetes</i> , 2019 , 68, 2287-2300	0.9	25
139	N-Glycosylation Defects in Humans Lower Low-Density Lipoprotein Cholesterol Through Increased Low-Density Lipoprotein Receptor Expression. <i>Circulation</i> , 2019 , 140, 280-292	16.7	18
138	A 3-SNP gene risk score and a metabolic risk score both predict hypertriglyceridemia and cardiovascular disease risk. <i>Journal of Clinical Lipidology</i> , 2019 , 13, 492-501	4.9	0
137	Oral vancomycin treatment does not alter markers of postprandial inflammation in lean and obese subjects. <i>Physiological Reports</i> , 2019 , 7, e14199	2.6	4
136	The Role of (Modified) Lipoproteins in Vascular Function: A Duet Between Monocytes and the Endothelium. <i>Current Medicinal Chemistry</i> , 2019 , 26, 1594-1609	4.3	11
135	Potential epigenetic therapeutics for atherosclerosis treatment. <i>Atherosclerosis</i> , 2019 , 281, 189-197	3.1	31
134	Effect of Vegan Fecal Microbiota Transplantation on Carnitine- and Choline-Derived Trimethylamine-N-Oxide Production and Vascular Inflammation in Patients With Metabolic Syndrome. <i>Journal of the American Heart Association</i> , 2018 , 7,	6	100
133	Decreased GPIHBP1 protein levels in visceral adipose tissue partly underlie the hypertriglyceridemic phenotype in insulin resistance. <i>PLoS ONE</i> , 2018 , 13, e0205858	3.7	2
132	A Deep Intronic Variant in LDLR in Familial Hypercholesterolemia. <i>Circulation Genomic and Precision Medicine</i> , 2018 , 11, e002385	5.2	12

131	Use of plasma metabolomics to analyze phenotype-genotype relationships in young hypercholesterolemic females. <i>Journal of Lipid Research</i> , 2018 , 59, 2174-2180	6.3	1
130	Proteoglycan 4 regulates macrophage function without altering atherosclerotic lesion formation in a murine bone marrow-specific deletion model. <i>Atherosclerosis</i> , 2018 , 274, 120-127	3.1	9
129	Naturally Occurring Variants in LRP1 (Low-Density Lipoprotein Receptor-Related Protein 1) Affect HDL (High-Density Lipoprotein) Metabolism Through ABCA1 (ATP-Binding Cassette A1) and SR-B1 (Scavenger Receptor Class B Type 1) in Humans. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018 , 38, 1440-1453	9.4	10
128	GPIHBP1 autoantibodies in a patient with unexplained chylomicronemia. <i>Journal of Clinical Lipidology</i> , 2017 , 11, 964-971	4.9	19
127	Improvement of Insulin Sensitivity after Lean Donor Feces in Metabolic Syndrome Is Driven by Baseline Intestinal Microbiota Composition. <i>Cell Metabolism</i> , 2017 , 26, 611-619.e6	24.6	440
126	Nile Red Quantifier: a novel and quantitative tool to study lipid accumulation in patient-derived circulating monocytes using confocal microscopy. <i>Journal of Lipid Research</i> , 2017 , 58, 2210-2219	6.3	11
125	ABCA8 Regulates Cholesterol Efflux and High-Density Lipoprotein Cholesterol Levels. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2017 , 37, 2147-2155	9.4	34
124	HDL functionality in type 1 diabetes. <i>Atherosclerosis</i> , 2017 , 267, 99-109	3.1	35
123	Intestinal <i>Ralstonia pickettii</i> augments glucose intolerance in obesity. <i>PLoS ONE</i> , 2017 , 12, e0181693	3.7	28
122	A rare variant in MCF2L identified using exclusion linkage in a pedigree with premature atherosclerosis. <i>European Journal of Human Genetics</i> , 2016 , 24, 86-91	5.3	7
121	Paraoxonase-1 activity is positively related to phospholipid transfer protein activity in type 2 diabetes mellitus: Role of large HDL particles. <i>Clinical Biochemistry</i> , 2016 , 49, 508-510	3.5	7
120	Triglyceride-Rich Lipoproteins and Remnants: Targets for Therapy?. <i>Current Cardiology Reports</i> , 2016 , 18, 67	4.2	55
119	Role of sulfatase 2 in lipoprotein metabolism and angiogenesis. <i>Current Opinion in Lipidology</i> , 2016 , 27, 181-6	4.4	5
118	Oral treatment with improves insulin sensitivity in mice. <i>Npj Biofilms and Microbiomes</i> , 2016 , 2, 16009	8.2	101
117	Liposomal prednisolone promotes macrophage lipotoxicity in experimental atherosclerosis. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2016 , 12, 1463-70	6	30
116	A loss-of-function variant in OSBPL1A predisposes to low plasma HDL cholesterol levels and impaired cholesterol efflux capacity. <i>Atherosclerosis</i> , 2016 , 249, 140-7	3.1	21
115	Causality of small and large intestinal microbiota in weight regulation and insulin resistance. <i>Molecular Metabolism</i> , 2016 , 5, 759-70	8.8	102
114	Prednisolone-containing liposomes accumulate in human atherosclerotic macrophages upon intravenous administration. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2015 , 11, 1039-46	6	97

113	Adipocyte ATP-binding cassette G1 promotes triglyceride storage, fat mass growth, and human obesity. <i>Diabetes</i> , 2015 , 64, 840-55	0.9	43
112	Annexin A5 haplotypes in familial hypercholesterolemia: lack of association with carotid intima-media thickness and cardiovascular disease risk. <i>Atherosclerosis</i> , 2015 , 238, 195-200	3.1	7
111	Phospholipid transfer protein activity and incident type 2 diabetes mellitus. <i>Clinica Chimica Acta</i> , 2015 , 439, 38-41	6.2	12
110	Increased large VLDL particles confer elevated cholesteryl ester transfer in diabetes. <i>European Journal of Clinical Investigation</i> , 2015 , 45, 36-44	4.6	11
109	Impact of oral vancomycin on gut microbiota, bile acid metabolism, and insulin sensitivity. <i>Journal of Hepatology</i> , 2014 , 60, 824-31	13.4	353
108	SULF2 strongly predisposes to fasting and postprandial triglycerides in patients with obesity and type 2 diabetes mellitus. <i>Obesity</i> , 2014 , 22, 1309-16	8	23
107	Targeting macrophage Histone deacetylase 3 stabilizes atherosclerotic lesions. <i>EMBO Molecular Medicine</i> , 2014 , 6, 1124-32	12	110
106	Towards network analysis to understand the genetic architecture of blood lipids: do the inclusion of age-dependency helps to identify seven novel loci?. <i>Atherosclerosis</i> , 2014 , 235, 642-3	3.1	
105	Abnormal hemostatic parameters in patients with myocardial infarction but angiographically normal coronary arteries. <i>International Journal of Cardiology</i> , 2014 , 174, 734-5	3.2	1
104	The endoplasmic reticulum coat protein II transport machinery coordinates cellular lipid secretion and cholesterol biosynthesis. <i>Journal of Biological Chemistry</i> , 2014 , 289, 4244-61	5.4	30
103	Akt2/LDLr double knockout mice display impaired glucose tolerance and develop more complex atherosclerotic plaques than LDLr knockout mice. <i>Cardiovascular Research</i> , 2014 , 101, 277-87	9.9	25
102	Mutations in STAP1 are associated with autosomal dominant hypercholesterolemia. <i>Circulation Research</i> , 2014 , 115, 552-5	15.7	114
101	Two novel mutations in apolipoprotein C3 underlie atheroprotective lipid profiles in families. <i>Clinical Genetics</i> , 2014 , 85, 433-40	4	15
100	HDL does not influence the polarization of human monocytes toward an alternative phenotype. <i>International Journal of Cardiology</i> , 2014 , 172, 179-84	3.2	18
99	Hepatic transforming growth factor- β stimulated clone-22 D1 controls systemic cholesterol metabolism. <i>Molecular Metabolism</i> , 2014 , 3, 155-66	8.8	3
98	Mutation in KERA identified by linkage analysis and targeted resequencing in a pedigree with premature atherosclerosis. <i>PLoS ONE</i> , 2014 , 9, e98289	3.7	5
97	Carriers of loss-of-function mutations in EXT display impaired pancreatic beta-cell reserve due to smaller pancreas volume. <i>PLoS ONE</i> , 2014 , 9, e115662	3.7	11
96	Lactate increases hepatic secretion of VLDL-triglycerides in humans. <i>Atherosclerosis</i> , 2013 , 228, 443-50	3.1	5

95	The positive relationship of serum paraoxonase-1 activity with apolipoprotein E is abrogated in metabolic syndrome. <i>Atherosclerosis</i> , 2013 , 230, 6-11	3.1	15
94	Hypertriglyceridemia: the future of genetics to guide individualized therapeutic strategies. <i>Clinical Lipidology</i> , 2013 , 8, 321-328		
93	Low normal thyroid function enhances plasma cholesteryl ester transfer in Type 2 diabetes mellitus. <i>Atherosclerosis</i> , 2013 , 228, 466-71	3.1	18
92	Pro-atherogenic lipid changes and decreased hepatic LDL receptor expression by tocilizumab in rheumatoid arthritis. <i>Atherosclerosis</i> , 2013 , 229, 174-81	3.1	72
91	ABCA1 mutation carriers with low high-density lipoprotein cholesterol are characterized by a larger atherosclerotic burden. <i>European Heart Journal</i> , 2013 , 34, 286-91	9.5	54
90	Identification of a loss-of-function inducible degrader of the low-density lipoprotein receptor variant in individuals with low circulating low-density lipoprotein. <i>European Heart Journal</i> , 2013 , 34, 1292-7	9.5	43
89	High density lipoprotein as a source of cholesterol for adrenal steroidogenesis: a study in individuals with low plasma HDL-C. <i>Journal of Lipid Research</i> , 2013 , 54, 1698-1704	6.3	37
88	Monocyte gene expression and coronary artery disease. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2013 , 16, 411-7	3.8	13
87	TSC22D4 is a molecular output of hepatic wasting metabolism. <i>EMBO Molecular Medicine</i> , 2013 , 5, 294-308		42
86	Pathophysiology of hypertriglyceridemia. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2012 , 1821, 826-32	5	50
85	Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. <i>Gastroenterology</i> , 2012 , 143, 913-6.e7	13.3	1766
84	Inhibition of hepatic sulfatase-2 in vivo: a novel strategy to correct diabetic dyslipidemia. <i>Hepatology</i> , 2012 , 55, 1746-53	11.2	36
83	Human ATP-binding cassette G1 controls macrophage lipoprotein lipase bioavailability and promotes foam cell formation. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2012 , 32, 2223-31	9.4	37
82	In Vivo Inflammation Does Not Impair ABCA1-Mediated Cholesterol Efflux Capacity of HDL. <i>Cholesterol</i> , 2012 , 2012, 610741		7
81	Impact of LDL apheresis on atheroprotective reverse cholesterol transport pathway in familial hypercholesterolemia. <i>Journal of Lipid Research</i> , 2012 , 53, 767-75	6.3	17
80	PS3 - 15. Genetic Variation at the SULF2 Locus Affects Hepatic Postprandial Remnant Clearance in Patients with Type 2 Diabetes Mellitus. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012 , 10, 109-109	0	
79	PS14 - 68. Differential effects of antibiotics on bile acid metabolism, intestinal microbiota composition and insulin resistance in obese humans; a randomised controlled trial. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012 , 10, 147-147	0	
78	PS14 - 70. Intestinal microbiota translocation is associated with inflamed visceral adipose tissue. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2012 , 10, 148-148	0	

77	Infusion of a lipid emulsion in healthy men decreases the serotonergic response. <i>Neuroendocrinology</i> , 2012 , 95, 325-31	5.6	4
76	Hepatic deficiency in transcriptional cofactor TBL1 promotes liver steatosis and hypertriglyceridemia. <i>Cell Metabolism</i> , 2011 , 13, 389-400	24.6	37
75	Heterozygosity for a loss-of-function mutation in GALNT2 improves plasma triglyceride clearance in man. <i>Cell Metabolism</i> , 2011 , 14, 811-8	24.6	75
74	High plasma cholesteryl ester transfer but not CETP mass predicts incident cardiovascular disease: a nested case-control study. <i>Atherosclerosis</i> , 2011 , 217, 249-52	3.1	27
73	Plasma phospholipid transfer protein activity is independently determined by obesity and insulin resistance in non-diabetic subjects. <i>Atherosclerosis</i> , 2011 , 217, 253-9	3.1	16
72	Genome-wide association studies in atherosclerosis. <i>Current Atherosclerosis Reports</i> , 2011 , 13, 225-32	6	21
71	Genetic variant of the scavenger receptor BI in humans. <i>New England Journal of Medicine</i> , 2011 , 364, 136-45	59.2	253
70	An increased burden of common and rare lipid-associated risk alleles contributes to the phenotypic spectrum of hypertriglyceridemia. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011 , 31, 1916-26	9.4	73
69	Excess of rare variants in genes identified by genome-wide association study of hypertriglyceridemia. <i>Nature Genetics</i> , 2010 , 42, 684-7	36.3	365
68	Genetic variation at the phospholipid transfer protein locus affects its activity and high-density lipoprotein size and is a novel marker of cardiovascular disease susceptibility. <i>Circulation</i> , 2010 , 122, 470-7	16.7	75
67	ABCG5/G8 polymorphisms and markers of cholesterol metabolism: systematic review and meta-analysis. <i>Journal of Lipid Research</i> , 2010 , 51, 3016-23	6.3	46
66	Chylomicronemia with low postheparin lipoprotein lipase levels in the setting of GPIIb/IIIa defects. <i>Circulation: Cardiovascular Genetics</i> , 2010 , 3, 169-78		90
65	Effect of increasing doses of Rosuvastatin and Atorvastatin on apolipoproteins, enzymes and lipid transfer proteins involved in lipoprotein metabolism and inflammatory parameters. <i>Current Medical Research and Opinion</i> , 2010 , 26, 2301-13	2.5	20
64	HDL cholesterol response to GH replacement is associated with common cholesteryl ester transfer protein gene variation (-629C>A) and modified by glucocorticoid treatment. <i>European Journal of Endocrinology</i> , 2010 , 162, 227-34	6.5	3
63	Atorvastatin treatment lowers fasting remnant-like particle cholesterol and LDL subfraction cholesterol without affecting LDL size in type 2 diabetes mellitus: Relevance for non-HDL cholesterol and apolipoprotein B guideline targets. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2010 , 1801, 89-94	5	18
62	Do genome-wide association scans provide additional information on the variation of plasma adiponectin concentrations?. <i>Atherosclerosis</i> , 2010 , 208, 328-9	3.1	14
61	The metabolism of triglyceride-rich lipoproteins revisited: new players, new insight. <i>Atherosclerosis</i> , 2010 , 211, 1-8	3.1	127
60	Carotid intima media thickness is associated with plasma adiponectin but not with the leptin:adiponectin ratio independently of metabolic syndrome. <i>Atherosclerosis</i> , 2010 , 211, 393-6	3.1	24

59	Lack of association between common genetic variation in endothelial lipase (LIPG) and the risk for CAD and DVT. <i>Atherosclerosis</i> , 2010 , 211, 558-64	3.1	35
58	Identification and characterization of novel loss of function mutations in ATP-binding cassette transporter A1 in patients with low plasma high-density lipoprotein cholesterol. <i>Atherosclerosis</i> , 2010 , 213, 492-8	3.1	43
57	Beneficial effects of reconstituted HDL on ex vivo and in vitro platelet reactivity. <i>Clinical Lipidology</i> , 2010 , 5, 167-171		1
56	Fasting cholesteryl ester transfer protein concentration is independently associated with the postprandial decrease in high-density lipoprotein cholesterol concentration after fat-rich meals: the Hoorn prandial study. <i>Metabolism: Clinical and Experimental</i> , 2010 , 59, 854-60	12.7	9
55	Chylomicronemia with a mutant GPIIIBP1 (Q115P) that cannot bind lipoprotein lipase. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2009 , 29, 956-62	9.4	134
54	Complement factor H Y402H decreases cardiovascular disease risk in patients with familial hypercholesterolaemia. <i>European Heart Journal</i> , 2009 , 30, 618-23	9.5	12
53	A plant stanol yogurt drink alone or combined with a low-dose statin lowers serum triacylglycerol and non-HDL cholesterol in metabolic syndrome patients. <i>Journal of Nutrition</i> , 2009 , 139, 1143-9	4.1	70
52	Prolactin levels and the risk of future coronary artery disease in apparently healthy men and women. <i>Circulation: Cardiovascular Genetics</i> , 2009 , 2, 389-95		35
51	Diagnostic value of post-heparin lipase testing in detecting common genetic variants in the LPL and LIPC genes. <i>European Journal of Human Genetics</i> , 2009 , 17, 1386-93	5.3	8
50	Plasma pre beta-HDL formation is decreased by atorvastatin treatment in type 2 diabetes mellitus: Role of phospholipid transfer protein. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2009 , 1791, 714-8	5	16
49	Novel roles of hepatic lipase and phospholipid transfer protein in VLDL as well as HDL metabolism. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2009 , 1791, 1031-6	5	14
48	Role of plasma adiponectin on the HDL-cholesterol raising effect of atorvastatin in patients with type 2 diabetes. <i>Current Medical Research and Opinion</i> , 2009 , 25, 93-101	2.5	15
47	Atorvastatin affects low density lipoprotein and non-high density lipoprotein cholesterol relations with apolipoprotein B in type 2 diabetes mellitus: modification by triglycerides and cholesteryl ester transfer protein. <i>Expert Opinion on Therapeutic Targets</i> , 2009 , 13, 743-51	6.4	7
46	The effect of statin alone or in combination with ezetimibe on postprandial lipoprotein composition in obese metabolic syndrome patients. <i>Atherosclerosis</i> , 2009 , 202, 216-24	3.1	35
45	ABCG8 gene polymorphisms, plasma cholesterol concentrations, and risk of cardiovascular disease in familial hypercholesterolemia. <i>Atherosclerosis</i> , 2009 , 204, 453-8	3.1	36
44	Phospholipid transfer protein activity is determined by type 2 diabetes mellitus and metabolic syndrome, and is positively associated with serum transaminases. <i>Clinical Endocrinology</i> , 2008 , 68, 375-81	3.4	19
43	Lipid-lowering therapy does not affect the postprandial drop in high density lipoprotein-cholesterol (HDL-c) plasma levels in obese men with metabolic syndrome: a randomized double blind crossover trial. <i>Clinical Endocrinology</i> , 2008 , 69, 870-7	3.4	16
42	Effects of short-term and long-term danazol treatment on lipoproteins, coagulation, and progression of atherosclerosis: two clinical trials in healthy volunteers and patients with hereditary angioedema. <i>Clinical Therapeutics</i> , 2008 , 30, 2314-23	3.5	35

41	Atorvastatin increases HDL cholesterol by reducing CETP expression in cholesterol-fed APOE*3-Leiden.CETP mice. <i>Atherosclerosis</i> , 2008 , 197, 57-63	3.1	69
40	High HDL cholesterol does not protect against coronary artery disease when associated with combined cholesteryl ester transfer protein and hepatic lipase gene variants. <i>Atherosclerosis</i> , 2008 , 200, 161-7	3.1	49
39	The efficacy of statin monotherapy uptitration versus switching to ezetimibe/simvastatin: results of the EASEGO study. <i>Current Medical Research and Opinion</i> , 2008 , 24, 685-94	2.5	37
38	Fibroblast cholesterol efflux to plasma from metabolic syndrome subjects is not defective despite low high-density lipoprotein cholesterol. <i>European Journal of Endocrinology</i> , 2008 , 158, 53-60	6.5	31
37	Fasting and postprandial remnant-like particle cholesterol concentrations in obese participants are associated with plasma triglycerides, insulin resistance, and body fat distribution. <i>Journal of Nutrition</i> , 2008 , 138, 2399-405	4.1	9
36	An apolipoprotein A-V gene SNP is associated with marked hypertriglyceridemia among Asian-American patients. <i>Journal of Lipid Research</i> , 2008 , 49, 1846-54	6.3	48
35	Gene-load score of the renin-angiotensin-aldosterone system is associated with coronary heart disease in familial hypercholesterolaemia. <i>European Heart Journal</i> , 2008 , 29, 1370-6	9.5	14
34	A more atherogenic serum lipoprotein profile is present in women with polycystic ovary syndrome: a case-control study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 470-6	5.6	129
33	Role of lipoprotein lipase in triglyceride metabolism: potential therapeutic target. <i>Future Lipidology</i> , 2008 , 3, 385-397		16
32	Abnormal patterns of lipoprotein lipase release into the plasma in GPIHBP1-deficient mice. <i>Journal of Biological Chemistry</i> , 2008 , 283, 34511-8	5.4	59
31	Renin-angiotensin-aldosterone responsiveness to low sodium and blood pressure reactivity to angiotensin-II are unrelated to cholesteryl ester transfer protein mass in healthy subjects. <i>Expert Opinion on Therapeutic Targets</i> , 2008 , 12, 1321-8	6.4	4
30	The effects of low-dose simvastatin and ezetimibe compared to high-dose simvastatin alone on post-fat load endothelial function in patients with metabolic syndrome: a randomized double-blind crossover trial. <i>Journal of Cardiovascular Pharmacology</i> , 2008 , 52, 145-50	3.1	37
29	Evaluation of phospholipid transfer protein as a therapeutic target. <i>Future Lipidology</i> , 2008 , 3, 327-335		9
28	Haplotype of the angiotensinogen gene is associated with coronary heart disease in familial hypercholesterolemia. <i>Journal of Hypertension</i> , 2008 , 26, 462-7	1.9	7
27	Plasma apolipoprotein CI correlates with increased survival in patients with severe sepsis. <i>Intensive Care Medicine</i> , 2008 , 34, 907-11	14.5	18
26	Derangements of intravascular remodeling of lipoproteins in type 2 diabetes mellitus: consequences for atherosclerosis development. <i>Current Diabetes Reports</i> , 2008 , 8, 65-70	5.6	6
25	Plasma phospholipid transfer protein activity, a determinant of HDL kinetics in vivo. <i>Clinical Endocrinology</i> , 2007 , 67, 316-7; author reply 317	3.4	1
24	Apolipoprotein A-II is inversely associated with risk of future coronary artery disease. <i>Circulation</i> , 2007 , 116, 2029-35	16.7	88

23	Cholesteryl ester transfer protein and hyperalphalipoproteinemia in Caucasians. <i>Journal of Lipid Research</i> , 2007 , 48, 674-82	6.3	26
22	Angiotensin II type 1 receptor blockade improves hyperglycemia-induced endothelial dysfunction and reduces proinflammatory cytokine release from leukocytes. <i>Journal of Cardiovascular Pharmacology</i> , 2007 , 49, 6-12	3.1	29
21	High plasma cholesteryl ester transfer protein levels may favour reduced incidence of cardiovascular events in men with low triglycerides. <i>European Heart Journal</i> , 2007 , 28, 1012-8	9.5	63
20	High plasma level of remnant-like particles cholesterol in familial combined hyperlipidemia. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 1269-75	5.6	19
19	Fenofibrate increases HDL-cholesterol by reducing cholesteryl ester transfer protein expression. <i>Journal of Lipid Research</i> , 2007 , 48, 1763-71	6.3	74
18	Concerted actions of cholesteryl ester transfer protein and phospholipid transfer protein in type 2 diabetes: effects of apolipoproteins. <i>Current Opinion in Lipidology</i> , 2007 , 18, 251-7	4.4	43
17	Haplotype analyses of the APOA5 gene in patients with familial combined hyperlipidemia. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2007 , 1772, 81-8	6.9	20
16	A functional polymorphism in the glucocorticoid receptor gene and its relation to cardiovascular disease risk in familial hypercholesterolemia. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2006 , 91, 4131-6	5.6	13
15	Plasma phospholipid transfer protein activity is decreased in type 2 diabetes during treatment with atorvastatin: a role for apolipoprotein E?. <i>Diabetes</i> , 2006 , 55, 1491-6	0.9	19
14	Cholesteryl ester transfer protein decreases high-density lipoprotein and severely aggravates atherosclerosis in APOE*3-Leiden mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006 , 26, 2552-9	9.4	169
13	Role of transcription factor KLF11 and its diabetes-associated gene variants in pancreatic beta cell function. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 4807-12	11.5	197
12	Plasma cholesteryl ester transfer is a determinant of intima-media thickness in type 2 diabetic and nondiabetic subjects: role of CETP and triglycerides. <i>Diabetes</i> , 2005 , 54, 3554-9	0.9	68
11	Alteration in lipoprotein lipase activity bound to triglyceride-rich lipoproteins in the postprandial state in type 2 diabetes. <i>Journal of Lipid Research</i> , 2004 , 45, 859-65	6.3	28
10	Atorvastatin decreases apolipoprotein C-III in apolipoprotein B-containing lipoprotein and HDL in type 2 diabetes: a potential mechanism to lower plasma triglycerides. <i>Diabetes Care</i> , 2004 , 27, 1358-64	14.6	42
9	Screening for functional sequence variations and mutations in ABCA1. <i>Atherosclerosis</i> , 2004 , 175, 269-79	3.1	39
8	Common cholesteryl ester transfer protein gene polymorphisms and the effect of atorvastatin therapy in type 2 diabetes. <i>Diabetes Care</i> , 2003 , 26, 1216-23	14.6	98
7	Elevated remnant-like particles in heterozygous familial hypercholesterolemia and response to statin therapy. <i>Circulation</i> , 2002 , 106, 788-92	16.7	29
6	Reduced structural and functional skin capillaries in familial combined hyperlipidemia affected men, associated with increased remnant-like lipoprotein cholesterol levels. <i>Atherosclerosis</i> , 2002 , 163, 355-62	3.1	10

5	The role of high density lipoprotein in sepsis. <i>Netherlands Journal of Medicine</i> , 2001 , 59, 102-10	0.5	40
4	Linkage of a candidate gene locus to familial combined hyperlipidemia: lecithin:cholesterol acyltransferase on 16q. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1999 , 19, 2730-6	9.4	46
3	A genome scan for familial combined hyperlipidemia reveals evidence of linkage with a locus on chromosome 11. <i>American Journal of Human Genetics</i> , 1999 , 65, 397-412	11	105
2	Cholesterol crystallization in human gallbladder bile: relation to gallstone number, bile composition, and apolipoprotein E4 isoform. <i>Hepatology</i> , 1998 , 27, 1508-16	11.2	31
1	Families with familial combined hyperlipidemia and families enriched for coronary artery disease share genetic determinants for the atherogenic lipoprotein phenotype. <i>American Journal of Human Genetics</i> , 1998 , 63, 577-85	11	53