

Stefano Bonapace

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1974970/publications.pdf>

Version: 2024-02-01

55
papers

2,069
citations

279701

23
h-index

233338

45
g-index

55
all docs

55
docs citations

55
times ranked

2967
citing authors

#	ARTICLE	IF	CITATIONS
1	Risk of cardiovascular, cardiac and arrhythmic complications in patients with non-alcoholic fatty liver disease. <i>World Journal of Gastroenterology</i> , 2014, 20, 1724.	1.4	207
2	Nonalcoholic Fatty Liver Disease Is Associated With Left Ventricular Diastolic Dysfunction in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 389-395.	4.3	159
3	Non-Alcoholic Fatty Liver Disease Is Associated with an Increased Incidence of Atrial Fibrillation in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2013, 8, e57183.	1.1	153
4	Prognostic Relevance of Pulmonary Arterial Compliance in Patients With Chronic Heart Failure. <i>Chest</i> , 2014, 145, 1064-1070.	0.4	127
5	Nonalcoholic Fatty Liver Disease Is Associated With Ventricular Arrhythmias in Patients With Type 2 Diabetes Referred for Clinically Indicated 24-Hour Holter Monitoring. <i>Diabetes Care</i> , 2016, 39, 1416-1423.	4.3	95
6	Independent and additional prognostic value of aminoterminal propeptide of type III procollagen circulating levels in patients with chronic heart failure. <i>Journal of Cardiac Failure</i> , 2004, 10, 403-411.	0.7	91
7	Heart valve calcification in patients with type 2 diabetes and nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2015, 64, 879-887.	1.5	82
8	Prognostic Impact of Diabetes on Long-term Survival Outcomes in Patients With Heart Failure: A Meta-analysis. <i>Diabetes Care</i> , 2017, 40, 1597-1605.	4.3	82
9	Nonalcoholic Fatty Liver Disease Is Independently Associated with Early Left Ventricular Diastolic Dysfunction in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2015, 10, e0135329.	1.1	81
10	Association between non-alcoholic fatty liver disease and risk of atrial fibrillation in adult individuals: An updated meta-analysis. <i>Liver International</i> , 2019, 39, 758-769.	1.9	75
11	Aortic Distensibility Independently Affects Exercise Tolerance in Patients With Dilated Cardiomyopathy. <i>Circulation</i> , 2003, 107, 1603-1608.	1.6	74
12	Aortic and Mitral Annular Calcifications Are Predictive of All-Cause and Cardiovascular Mortality in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 1781-1786.	4.3	62
13	Relation of Elevated Serum Uric Acid Levels to Incidence of Atrial Fibrillation in Patients With Type 2 Diabetes Mellitus. <i>American Journal of Cardiology</i> , 2013, 112, 499-504.	0.7	58
14	Left atrial volume provides independent and incremental information compared with exercise tolerance parameters in patients with heart failure and left ventricular systolic dysfunction. <i>Heart</i> , 2007, 93, 1420-1425.	1.2	52
15	Nonalcoholic Fatty Liver Disease Is Associated with Aortic Valve Sclerosis in Patients with Type 2 Diabetes Mellitus. <i>PLoS ONE</i> , 2014, 9, e88371.	1.1	49
16	Risk of Heart Failure in Patients With Nonalcoholic Fatty Liver Disease. <i>Journal of the American College of Cardiology</i> , 2022, 79, 180-191.	1.2	46
17	Effects of ACE gene insertion/deletion polymorphism on response to spironolactone in patients with chronic heart failure. <i>American Journal of Medicine</i> , 2004, 116, 657-661.	0.6	43
18	Nonalcoholic fatty liver disease is associated with an increased risk of heart block in hospitalized patients with type 2 diabetes mellitus. <i>PLoS ONE</i> , 2017, 12, e0185459.	1.1	42

#	ARTICLE	IF	CITATIONS
19	Non-alcoholic fatty liver disease and increased risk of all-cause mortality in elderly patients admitted for acute heart failure. <i>International Journal of Cardiology</i> , 2018, 265, 162-168.	0.8	41
20	Nonalcoholic fatty liver disease and increased risk of 1-year all-cause and cardiac hospital readmissions in elderly patients admitted for acute heart failure. <i>PLoS ONE</i> , 2017, 12, e0173398.	1.1	38
21	Carotid atherosclerotic plaque instability in patients with acute myocardial infarction. <i>International Journal of Cardiology</i> , 2006, 111, 263-266.	0.8	32
22	Increased Aortic Pulse Wave Velocity as Measured by Echocardiography Is Strongly Associated with Poor Prognosis in Patients with Heart Failure. <i>Journal of the American Society of Echocardiography</i> , 2013, 26, 714-720.	1.2	31
23	Association of Plasma Ceramides With Myocardial Perfusion in Patients With Coronary Artery Disease Undergoing Stress Myocardial Perfusion Scintigraphy. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 2854-2861.	1.1	29
24	Early impairment in left ventricular longitudinal systolic function is associated with an increased risk of incident atrial fibrillation in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 413-418.	1.2	24
25	Aortic stiffness correlates with an increased extracellular matrix turnover in patients with dilated cardiomyopathy. <i>American Heart Journal</i> , 2006, 152, 93.e1-93.e6.	1.2	23
26	Antiaggregating and vasodilatory effects of a new nitroderivative of acetylsalicylic acid. <i>Thrombosis Research</i> , 1995, 80, 367-376.	0.8	22
27	Aortic Valve Sclerosis: A Marker of Significant Obstructive Coronary Artery Disease in Patients with Chest Pain?. <i>Journal of the American Society of Echocardiography</i> , 2007, 20, 703-708.	1.2	22
28	Mitral and aortic valve sclerosis/calcification and carotid atherosclerosis: results from 1065 patients. <i>Heart and Vessels</i> , 2014, 29, 776-783.	0.5	22
29	Effect of single and repeated doses of a new nitroderivative of acetylsalicylic acid on platelet TXA2 production in rats. <i>Life Sciences</i> , 1996, 58, PL207-PL210.	2.0	21
30	Increased Pulse Pressure Independently Predicts Incident Atrial Fibrillation in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 2337-2339.	4.3	20
31	Mitral Regurgitation and Increased Risk of All-Cause and Cardiovascular Mortality in Patients with Type 2 Diabetes. <i>American Journal of Medicine</i> , 2017, 130, 70-76.e1.	0.6	18
32	Nonalcoholic Fatty Liver Disease Is Associated With Higher 1-year All-Cause Rehospitalization Rates in Patients Admitted for Acute Heart Failure. <i>Medicine (United States)</i> , 2016, 95, e2760.	0.4	17
33	Association between plasma ceramides and inducible myocardial ischemia in patients with established or suspected coronary artery disease undergoing myocardial perfusion scintigraphy. <i>Metabolism: Clinical and Experimental</i> , 2018, 85, 305-312.	1.5	15
34	Aortic stiffness: an old concept for new insights into the pathophysiology of functional mitral regurgitation. <i>Heart and Vessels</i> , 2013, 28, 606-612.	0.5	14
35	Relationship Between Early Diastolic Dysfunction and Abnormal Microvolt T-Wave Alternans in Patients With Type 2 Diabetes. <i>Circulation: Cardiovascular Imaging</i> , 2011, 4, 408-414.	1.3	12
36	Relationship between increased left atrial volume and microvascular complications in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 822-828.	1.2	12

#	ARTICLE	IF	CITATIONS
37	Relation between plasma ceramides and cardiovascular death in chronic heart failure: A subset analysis of the GISSI-HF trial. <i>ESC Heart Failure</i> , 2020, 7, 3288-3297.	1.4	12
38	Association between subclinical left ventricular systolic dysfunction and glycemic control in asymptomatic type 2 diabetic patients with preserved left ventricular function. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1035-1040.	1.2	11
39	Echocardiographically Derived Pulse Wave Velocity and Diastolic Dysfunction Are Associated with an Increased Incidence of Atrial Fibrillation in Patients with Systolic Heart Failure. <i>Echocardiography</i> , 2016, 33, 1024-1031.	0.3	10
40	Relation of elevated serum uric acid levels to first-degree heart block and other cardiac conduction defects in hospitalized patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1691-1697.	1.2	10
41	Renal arterial pulsatility predicts progression of chronic kidney disease in chronic heart failure patients. <i>International Journal of Cardiology</i> , 2013, 167, 3050-3051.	0.8	6
42	Aortic valve sclerosis is a marker of atherosclerosis independently of traditional clinical risk factors. Analysis in 712 patients without ischemic heart disease. <i>International Journal of Cardiology</i> , 2012, 158, 163-164.	0.8	5
43	A higher body mass index is associated with reduced prevalence of unstable atherosclerotic plaque: A possible explanation of the obesity paradox. <i>International Journal of Cardiology</i> , 2013, 168, 2912-2913.	0.8	5
44	Left ventricular chamber dilation and filling pressure may help to categorise patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2018, 6, e000529.	1.2	4
45	Left atrial appendage closure in a patient with cor triatriatum and ASD: the added value of 3D echocardiography. <i>European Heart Journal Cardiovascular Imaging</i> , 2016, 17, 753-753.	0.5	3
46	Unusual echocardiographic image of a left circumflex coronary artery with anomalous origin from the right coronary sinus. <i>European Heart Journal</i> , 2016, 37, 792-792.	1.0	2
47	Increased aortic stiffness in adults with chronic indeterminate Chagas disease. <i>PLoS ONE</i> , 2019, 14, e0220689.	1.1	2
48	Brachial pulse pressure in acute heart failure. Results of the Heart Failure Registry. <i>ESC Heart Failure</i> , 2019, 6, 1167-1177.	1.4	2
49	Does high LDL-cholesterol cause cardiovascular disease?. <i>Expert Review of Clinical Pharmacology</i> , 2019, 12, 91-91.	1.3	2
50	Role of Multimodality Imaging in a Case of Aortic Prosthetic Endocarditis. <i>Circulation: Cardiovascular Imaging</i> , 2019, 12, e009460.	1.3	1
51	Regurgitant Volume/Left Ventricular End-Diastolic Volume Ratio. <i>JACC: Cardiovascular Imaging</i> , 2021, 14, 880.	2.3	1
52	Association between KLF6 rs3750861 polymorphism and plasma ceramide concentrations in post-menopausal women with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2022, 32, 1283-1287.	1.1	1
53	Association between Higher Circulating Leucine-Rich Î±-2 Glycoprotein 1 Concentrations and Specific Plasma Ceramides in Postmenopausal Women with Type 2 Diabetes. <i>Biomolecules</i> , 2022, 12, 943.	1.8	1
54	Inter-atrial shunt inversion by the sitting position in a patient with a patent foramen ovale and acute pulmonary embolism. <i>European Heart Journal</i> , 2014, 35, 1032-1032.	1.0	0

#	ARTICLE	IF	CITATIONS
55	Prognostic Role of Pericardial Fat on the Incidence of Heart Failure. Journal of the American College of Cardiology, 2021, 78, e111.	1.2	0