
## Stefano Bonapace

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1974970/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Risk of HeartÂFailure in Patients With Nonalcoholic Fatty Liver Disease. Journal of the American College of Cardiology, 2022, 79, 180-191.	1.2	46
2	Association between KLF6 rs3750861 polymorphism and plasma ceramide concentrations in post-menopausal women with type 2 diabetes. Nutrition, Metabolism and Cardiovascular Diseases, 2022, 32, 1283-1287.	1.1	1
3	Association between Higher Circulating Leucine-Rich α-2 Glycoprotein 1 Concentrations and Specific Plasma Ceramides in Postmenopausal Women with Type 2 Diabetes. Biomolecules, 2022, 12, 943.	1.8	1
4	Regurgitant Volume/Left Ventricular End-Diastolic Volume Ratio. JACC: Cardiovascular Imaging, 2021, 14, 880.	2.3	1
5	Prognostic Role of Pericardial Fat on the Incidence of HeartÂFailure. Journal of the American College of Cardiology, 2021, 78, e111.	1.2	0
6	Relation between plasma ceramides and cardiovascular death in chronic heart failure: A subset analysis of the GISSIâ€HF trial. ESC Heart Failure, 2020, 7, 3288-3297.	1.4	12
7	Increased aortic stiffness in adults with chronic indeterminate Chagas disease. PLoS ONE, 2019, 14, e0220689.	1.1	2
8	Association between nonâ€alcoholic fatty liver disease and risk of atrial fibrillation in adult individuals: An updated metaâ€analysis. Liver International, 2019, 39, 758-769.	1.9	75
9	Brachial pulse pressure in acute heart failure. Results of the Heart Failure Registry. ESC Heart Failure, 2019, 6, 1167-1177.	1.4	2
10	Role of Multimodality Imaging in a Case of Aortic Prosthetic Endocarditis. Circulation: Cardiovascular Imaging, 2019, 12, e009460.	1.3	1
11	Does high LDL-cholesterol cause cardiovascular disease?. Expert Review of Clinical Pharmacology, 2019, 12, 91-91.	1.3	2
12	Non-alcoholic fatty liver disease and increased risk of all-cause mortality in elderly patients admitted for acute heart failure. International Journal of Cardiology, 2018, 265, 162-168.	0.8	41
13	Association of Plasma Ceramides With Myocardial Perfusion in Patients With Coronary Artery Disease Undergoing Stress Myocardial Perfusion Scintigraphy. Arteriosclerosis, Thrombosis, and Vascular Biology, 2018, 38, 2854-2861.	1.1	29
14	Association between plasma ceramides and inducible myocardial ischemia in patients with established or suspected coronary artery disease undergoing myocardial perfusion scintigraphy. Metabolism: Clinical and Experimental, 2018, 85, 305-312.	1.5	15
15	Left ventricular chamber dilation and filling pressure may help to categorise patients with type 2 diabetes. BMJ Open Diabetes Research and Care, 2018, 6, e000529.	1.2	4
16	Early impairment in left ventricular longitudinal systolic function is associated with an increased risk of incident atrial fibrillation in patients with type 2 diabetes. Journal of Diabetes and Its Complications, 2017, 31, 413-418.	1.2	24
17	Association between subclinical left ventricular systolic dysfunction and glycemic control in asymptomatic type 2 diabetic patients with preserved left ventricular function. Journal of Diabetes and Its Complications, 2017, 31, 1035-1040.	1.2	11
18	Relation of elevated serum uric acid levels to first-degree heart block and other cardiac conduction defects in hospitalized patients with type 2 diabetes. Journal of Diabetes and Its Complications, 2017, 31, 1691-1697.	1.2	10

STEFANO BONAPACE

#	Article	IF	CITATIONS
19	Prognostic Impact of Diabetes on Long-term Survival Outcomes in Patients With Heart Failure: A Meta-analysis. Diabetes Care, 2017, 40, 1597-1605.	4.3	82
20	Mitral Regurgitation and Increased Risk of All-Cause and Cardiovascular Mortality in Patients with Type 2 Diabetes. American Journal of Medicine, 2017, 130, 70-76.e1.	0.6	18
21	Nonalcoholic fatty liver disease and increased risk of 1-year all-cause and cardiac hospital readmissions in elderly patients admitted for acute heart failure. PLoS ONE, 2017, 12, e0173398.	1.1	38
22	Nonalcoholic fatty liver disease is associated with an increased risk of heart block in hospitalized patients with type 2 diabetes mellitus. PLoS ONE, 2017, 12, e0185459.	1.1	42
23	Echocardiographically Derived Pulse Wave Velocity and Diastolic Dysfunction Are Associated with an Increased Incidence of Atrial Fibrillation in Patients with Systolic Heart Failure. Echocardiography, 2016, 33, 1024-1031.	0.3	10
24	Nonalcoholic Fatty Liver Disease Is Associated With Ventricular Arrhythmias in Patients With Type 2 Diabetes Referred for Clinically Indicated 24-Hour Holter Monitoring. Diabetes Care, 2016, 39, 1416-1423.	4.3	95
25	Left atrial appendage closure in a patient with cor triatriatum and ASD: the added value of 3D echocardiography. European Heart Journal Cardiovascular Imaging, 2016, 17, 753-753.	0.5	3
26	Unusual echocardiographic image of a left circumflex coronary artery with anomalous origin from the right coronary sinus. European Heart Journal, 2016, 37, 792-792.	1.0	2
27	Nonalcoholic Fatty Liver Disease Is Associated With Higher 1-year All-Cause Rehospitalization Rates in Patients Admitted for Acute Heart Failure. Medicine (United States), 2016, 95, e2760.	0.4	17
28	Nonalcoholic Fatty Liver Disease Is Independently Associated with Early Left Ventricular Diastolic Dysfunction in Patients with Type 2 Diabetes. PLoS ONE, 2015, 10, e0135329.	1.1	81
29	Heart valve calcification in patients with type 2 diabetes and nonalcoholic fatty liver disease. Metabolism: Clinical and Experimental, 2015, 64, 879-887.	1.5	82
30	Relationship between increased left atrial volume and microvascular complications in patients with type 2 diabetes. Journal of Diabetes and Its Complications, 2015, 29, 822-828.	1.2	12
31	Inter-atrial shunt inversion by the sitting position in a patient with a patent foramen ovale and acute pulmonary embolism. European Heart Journal, 2014, 35, 1032-1032.	1.0	0
32	Mitral and aortic valve sclerosis/calcification and carotid atherosclerosis: results from 1065 patients. Heart and Vessels, 2014, 29, 776-783.	0.5	22
33	Risk of cardiovascular, cardiac and arrhythmic complications in patients with non-alcoholic fatty liver disease. World Journal of Gastroenterology, 2014, 20, 1724.	1.4	207
34	Prognostic Relevance of Pulmonary Arterial Compliance in Patients With Chronic Heart Failure. Chest, 2014, 145, 1064-1070.	0.4	127
35	Nonalcoholic Fatty Liver Disease Is Associated with Aortic Valve Sclerosis in Patients with Type 2 Diabetes Mellitus. PLoS ONE, 2014, 9, e88371.	1.1	49
36	Increased Aortic Pulse Wave Velocity as Measured by Echocardiography Is Strongly Associated with Poor Prognosis in Patients with Heart Failure. Journal of the American Society of Echocardiography, 2013, 26, 714-720.	1.2	31

STEFANO BONAPACE

#	Article	IF	CITATIONS
37	Aortic stiffness: an old concept for new insights into the pathophysiology of functional mitral regurgitation. Heart and Vessels, 2013, 28, 606-612.	0.5	14
38	Relation of Elevated Serum Uric Acid Levels to Incidence of Atrial Fibrillation in Patients With Type 2 Diabetes Mellitus. American Journal of Cardiology, 2013, 112, 499-504.	0.7	58
39	Renal arterial pulsatility predicts progression of chronic kidney disease in chronic heart failure patients. International Journal of Cardiology, 2013, 167, 3050-3051.	0.8	6
40	A higher body mass index is associated with reduced prevalence of unstable atherosclerotic plaque: A possible explanation of the obesity paradox. International Journal of Cardiology, 2013, 168, 2912-2913.	0.8	5
41	Non-Alcoholic Fatty Liver Disease Is Associated with an Increased Incidence of Atrial Fibrillation in Patients with Type 2 Diabetes. PLoS ONE, 2013, 8, e57183.	1.1	153
42	Increased Pulse Pressure Independently Predicts Incident Atrial Fibrillation in Patients With Type 2 Diabetes. Diabetes Care, 2012, 35, 2337-2339.	4.3	20
43	Nonalcoholic Fatty Liver Disease Is Associated With Left Ventricular Diastolic Dysfunction in Patients With Type 2 Diabetes. Diabetes Care, 2012, 35, 389-395.	4.3	159
44	Aortic valve sclerosis is a marker of atherosclerosis independently of traditional clinical risk factors. Analysis in 712 patients without ischemic heart disease. International Journal of Cardiology, 2012, 158, 163-164.	0.8	5
45	Aortic and Mitral Annular Calcifications Are Predictive of All-Cause and Cardiovascular Mortality in Patients With Type 2 Diabetes. Diabetes Care, 2012, 35, 1781-1786.	4.3	62
46	Relationship Between Early Diastolic Dysfunction and Abnormal Microvolt T-Wave Alternans in Patients With Type 2 Diabetes. Circulation: Cardiovascular Imaging, 2011, 4, 408-414.	1.3	12
47	Left atrial volume provides independent and incremental information compared with exercise tolerance parameters in patients with heart failure and left ventricular systolic dysfunction. Heart, 2007, 93, 1420-1425.	1.2	52
48	Aortic Valve Sclerosis: A Marker of Significant Obstructive Coronary Artery Disease in Patients with Chest Pain?. Journal of the American Society of Echocardiography, 2007, 20, 703-708.	1.2	22
49	Aortic stiffness correlates with an increased extracellular matrix turnover in patients with dilated cardiomyopathy. American Heart Journal, 2006, 152, 93.e1-93.e6.	1.2	23
50	Carotid atherosclerotic plaque instability in patients with acute myocardial infarction. International Journal of Cardiology, 2006, 111, 263-266.	0.8	32
51	Independent and additional prognostic value of aminoterminal propeptide of type III procollagen circulating levels in patients with chronic heart failure. Journal of Cardiac Failure, 2004, 10, 403-411.	0.7	91
52	Effects of ACE gene insertion/deletion polymorphism on response to spironolactone in patients with chronic heart failure. American Journal of Medicine, 2004, 116, 657-661.	0.6	43
53	Aortic Distensibility Independently Affects Exercise Tolerance in Patients With Dilated Cardiomyopathy. Circulation, 2003, 107, 1603-1608.	1.6	74
54	Effect of single and repeated doses of a new nitroderivative of acetylsalicylic acid on platelet TXA2 production in rats. Life Sciences, 1996, 58, PL207-PL210.	2.0	21

#	Article	IF	CITATIONS
55	Antiaggregating and vasodilatory effects of a new nitroderivative of acetylsalicylic acid. Thrombosis Research, 1995, 80, 367-376.	0.8	22