

# Chin-Chan Lee

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1977791/publications.pdf>

Version: 2024-02-01

37  
papers

658  
citations

687363

13  
h-index

610901

24  
g-index

38  
all docs

38  
docs citations

38  
times ranked

1139  
citing authors

#	ARTICLE	IF	CITATIONS
1	Neutralization Assessments Reveal High Cardiothoracic Ratio and Old Age as Independent Predictors of Low Neutralizing Antibody Titers in Hemodialysis Patients Receiving a Single Dose of COVID-19 Vaccine. <i>Journal of Personalized Medicine</i> , 2022, 12, 68.	2.5	7
2	Hospitalization in patients with dialysis in Taiwan: A nationwide population-based observational study. <i>Journal of the Formosan Medical Association</i> , 2022, 121, S39-S46.	1.7	5
3	Ketoanalogue supplements reduce mortality in patients with pre-dialysis advanced diabetic kidney disease: A nationwide population-based study. <i>Clinical Nutrition</i> , 2021, 40, 4149-4160.	5.0	8
4	Renal function trajectories in hepatitis C infection: differences between renal healthy and chronic kidney disease individuals. <i>Scientific Reports</i> , 2021, 11, 17197.	3.3	5
5	Effective Preventive Strategies to Prevent Secondary Transmission of COVID-19 in Hemodialysis Unit: The First Month of Community Outbreak in Taiwan. <i>Healthcare (Switzerland)</i> , 2021, 9, 1173.	2.0	5
6	Xenoestrogen exposure and kidney function in the general population: Results of a community-based study by laboratory tests and questionnaire-based interviewing. <i>Environment International</i> , 2021, 155, 106585.	10.0	7
7	Incidence and Clinical Impacts of COVID-19 Infection in Patients with Hemodialysis: Systematic Review and Meta-Analysis of 396,062 Hemodialysis Patients. <i>Healthcare (Switzerland)</i> , 2021, 9, 47.	2.0	31
8	The association between low protein diet and body composition, muscle function, inflammation, and amino acid-based metabolic profile in chronic kidney disease stage 3-5 patients. <i>Clinical Nutrition ESPEN</i> , 2021, 46, 405-415.	1.2	12
9	The prognostic value of peripheral total and differential leukocyte count in renal progression: A community-based study. <i>PLoS ONE</i> , 2021, 16, e0258210.	2.5	7
10	Investigation of Possible Correlation Between Retinal Neurovascular Biomarkers and Early Cognitive Impairment in Patients With Chronic Kidney Disease. <i>Translational Vision Science and Technology</i> , 2021, 10, 9.	2.2	6
11	Dynamic Echocardiographic Assessments Reveal Septal E/e <sup>™</sup> Ratio as Independent Predictor of Intradialytic Hypotension in Maintenance for Hemodialysis Patients with Preserved Ejection Fraction. <i>Diagnostics</i> , 2021, 11, 2266.	2.6	0
12	Impact of blood pressure control on retinal microvasculature in patients with chronic kidney disease. <i>Scientific Reports</i> , 2020, 10, 14275.	3.3	7
13	Compositional and Functional Adaptations of Intestinal Microbiota and Related Metabolites in CKD Patients Receiving Dietary Protein Restriction. <i>Nutrients</i> , 2020, 12, 2799.	4.1	20
14	Vitamin D deficiency, cardiothoracic ratio, and long-term mortality in hemodialysis patients. <i>Scientific Reports</i> , 2020, 10, 7533.	3.3	4
15	Retinal neurovascular changes in chronic kidney disease. <i>Acta Ophthalmologica</i> , 2020, 98, e848-e855.	1.1	14
16	Gut Microbiota as Diagnostic Tools for Mirroring Disease Progression and Circulating Nephrotoxin Levels in Chronic Kidney Disease: Discovery and Validation Study. <i>International Journal of Biological Sciences</i> , 2020, 16, 420-434.	6.4	64
17	Early retinal microvascular abnormalities in patients with chronic kidney disease. <i>Microcirculation</i> , 2019, 26, e12555.	1.8	38
18	The association between chronic musculoskeletal pain and clinical outcome in chronic kidney disease patients: a prospective cohort study. <i>Renal Failure</i> , 2019, 41, 257-266.	2.1	12

#	ARTICLE	IF	CITATIONS
19	Vitamin D deficiency in northern Taiwan: a community-based cohort study. <i>BMC Public Health</i> , 2019, 19, 337.	2.9	47
20	Correlation between spot and 24h proteinuria: Derivation and validation of equation to estimate daily proteinuria. <i>PLoS ONE</i> , 2019, 14, e0214614.	2.5	15
21	Clinical association between the metabolite of healthy gut microbiota, 3-indolepropionic acid and chronic kidney disease. <i>Clinical Nutrition</i> , 2019, 38, 2945-2948.	5.0	42
22	Encapsulating Peritoneal Sclerosis in Long-Termed Peritoneal Dialysis Patients. <i>BioMed Research International</i> , 2018, 2018, 1-7.	1.9	6
23	Multidisciplinary predialysis education reduces incidence of peritonitis and subsequent death in peritoneal dialysis patients: 5-year cohort study. <i>PLoS ONE</i> , 2018, 13, e0202781.	2.5	15
24	Association between macular degeneration and mild to moderate chronic kidney disease. <i>Medicine (United States)</i> , 2017, 96, e6405.	1.0	11
25	Association between Parathyroid Hormone, 25 (OH) Vitamin D, and Chronic Kidney Disease: A Population-Based Study. <i>BioMed Research International</i> , 2017, 2017, 1-9.	1.9	12
26	Protein-bound uremic toxins impaired mitochondrial dynamics and functions. <i>Oncotarget</i> , 2017, 8, 77722-77733.	1.8	34
27	Association between human leucocyte antigen subtypes and risk of end stage renal disease in Taiwanese: a retrospective study. <i>BMC Nephrology</i> , 2015, 16, 177.	1.8	16
28	Correlation and presentation of thyroid functional status with thyroid autoantibodies in long-term follow-up of autoimmune thyroiditis: A study of 116 cases. <i>Journal of the Formosan Medical Association</i> , 2015, 114, 1039-1046.	1.7	5
29	Protein-Bound Uremic Toxins Induce Tissue Remodeling by Targeting the EGF Receptor. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 281-290.	6.1	34
30	The Association of Uremic Toxins and Inflammation in Hemodialysis Patients. <i>PLoS ONE</i> , 2014, 9, e102691.	2.5	23
31	Chronic Kidney Disease Stage Is a Modulator on the Association between High-Sensitivity C-Reactive Protein and Coronary Vasospastic Angina. <i>Scientific World Journal</i> , The, 2014, 2014, 1-9.	2.1	5
32	Depression, 5HTTLPR and BDNF Val66Met polymorphisms, and plasma BDNF levels in hemodialysis patients with chronic renal failure. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 1235.	2.2	10
33	Oral adsorbent AST-120 potentiates the effect of erythropoietin-stimulating agents on Stage 5 chronic kidney disease patients: a randomized crossover study. <i>Nephrology Dialysis Transplantation</i> , 2014, 29, 1719-1727.	0.7	43
34	Multidisciplinary Predialysis Education Reduced the Inpatient and Total Medical Costs of the First 6 Months of Dialysis in Incident Hemodialysis Patients. <i>PLoS ONE</i> , 2014, 9, e112820.	2.5	41
35	Re-evaluating the predictive roles of metabolic complications and clinical outcome according to eGFR levels "a four-years prospective cohort study in Taiwan. <i>BMC Nephrology</i> , 2013, 14, 92.	1.8	9
36	Positive Dialysate Gram Stain Predicts Outcome of Empirical Antibiotic Therapy for Peritoneal Dialysis-associated Peritonitis. <i>Therapeutic Apheresis and Dialysis</i> , 2010, 14, 201-208.	0.9	13

#	ARTICLE	IF	CITATIONS
37	Long-term modality-related mortality analysis in incident dialysis patients. <i>Peritoneal Dialysis International</i> , 2009, 29, 182-90.	2.3	25