

# Nicolas Fabresse

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1979186/publications.pdf>

Version: 2024-02-01

16  
papers

324  
citations

933447

10  
h-index

996975

15  
g-index

18  
all docs

18  
docs citations

18  
times ranked

321  
citing authors

#	ARTICLE	IF	CITATIONS
1	High-resolution mass spectrometry: Theoretical and technological aspects. <i>Toxicologie Analytique Et Clinique</i> , 2022, 34, 3-18.	0.1	2
2	MÃ©usage de prÃ©gabaline: Ã  propos de sept cas de dÃ©cÃ©s en rÃ©gion marseillaise. <i>Toxicologie Analytique Et Clinique</i> , 2022, 34, 151-158.	0.1	1
3	Les gabapentinoÃ©des: une revue de la littÃ©rature. <i>Toxicologie Analytique Et Clinique</i> , 2021, 33, 44-63.	0.1	0
4	Hair testing for 3-fluorofentanyl, furanylfentanyl, methoxyacetylfentanyl, carfentanil, acetylfentanyl and fentanyl by LC-MS/MS after unintentional overdose. <i>Forensic Toxicology</i> , 2020, 38, 277-286.	2.4	16
5	P1392ANALYSIS OF CALCIFYING POTENTIAL OF UREMIC SERUM FROM HAEMODIALYSIS PATIENTS TREATED WITH A MEDIUM CUT-OFF (THERANOVA) DIALYSER: A PROSPECTIVE, CROSS-OVER STUDY. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
6	Prevalence of New Psychoactive Substances (NPS) and conventional drugs of abuse (DOA) in high risk populations from Paris (France) and its suburbs. <i>Drug and Alcohol Dependence</i> , 2019, 204, 107508.	3.2	53
7	Novel synthetic opioids: A review of the literature. <i>Toxicologie Analytique Et Clinique</i> , 2019, 31, 298-316.	0.1	17
8	Development and validation of a liquid chromatography-tandem mass spectrometry method for simultaneous detection of 10 illicit drugs in oral fluid collected with FLOQSwabs, and application to real samples. <i>Drug Testing and Analysis</i> , 2019, 11, 824-832.	2.6	20
9	Development of a sensitive untargeted liquid chromatography-high resolution mass spectrometry screening devoted to hair analysis through a shared MS2 spectra database: A step toward early detection of new psychoactive substances. <i>Drug Testing and Analysis</i> , 2019, 11, 697-708.	2.6	57
10	Validation of an UPLC-MS/MS method for the determination of sixteen synthetic cannabinoids in human hair. Application to document chronic use of JWH-122 following a non-fatal overdose. <i>Toxicologie Analytique Et Clinique</i> , 2019, 31, 283-292.	0.1	3
11	Drug-facilitated sexual assault (DFSA) involving 4-methylethcathinone (4-MEC), 3,4-methylenedioxypyrovalerone (MDPV), and doxylamine highlighted by hair analysis. <i>Drug Testing and Analysis</i> , 2018, 10, 1280-1284.	2.6	64
12	Hair analysis does not allow to discriminate between acute and chronic administrations of a drug in young children. <i>International Journal of Legal Medicine</i> , 2018, 132, 165-172.	2.2	34
13	LC-MS/MS method for quantification of baclofen in hair: A useful tool to assess compliance in alcohol dependent patients?. <i>Drug Testing and Analysis</i> , 2018, 10, 694-700.	2.6	13
14	Prevalence and Surveillance of Synthetic Cathinones Use by Hair Analysis: An Update Review. <i>Current Pharmaceutical Design</i> , 2018, 23, 5487-5495.	1.9	11
15	Identification and quantification of diphenidine in hair by LC-MS/MS after single administration. <i>Toxicologie Analytique Et Clinique</i> , 2017, 29, 64-70.	0.1	7
16	Identification and quantification of 4-methylethcathinone (4-MEC) and 3,4-methylenedioxypyrovalerone (MDPV) in hair by LC-MS/MS after chronic administration. <i>Forensic Science International</i> , 2017, 270, 39-45.	2.2	25