

Stephen C Jameson

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1982013/stephen-c-jameson-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

168
papers

21,198
citations

70
h-index

145
g-index

199
ext. papers

23,914
ext. citations

16
avg, IF

6.97
L-index

#	Paper	IF	Citations
168	Sphingosine 1-phosphate receptor 5 (S1PR5) regulates the peripheral retention of tissue-resident lymphocytes. <i>Journal of Experimental Medicine</i> , 2022 , 219,	16.6	11
167	CoAching CD8 T cells for tumor immunotherapy-the pantothenate way. <i>Cell Metabolism</i> , 2021 , 33, 2305-2316	14.6	0
166	Engagement of the costimulatory molecule ICOS in tissues promotes establishment of CD8 tissue-resident memory T cells. <i>Immunity</i> , 2021 ,	32.3	4
165	CD8 T cell self-tolerance permits responsiveness but limits tissue damage. <i>ELife</i> , 2021 , 10,	8.9	3
164	Classical MHC expression by DP thymocytes impairs the selection of non-classical MHC restricted innate-like T cells. <i>Nature Communications</i> , 2021 , 12, 2308	17.4	6
163	Senolytics reduce coronavirus-related mortality in old mice. <i>Science</i> , 2021 , 373,	33.3	60
162	Inflating the role of stromal cells in CD8 T cell memory. <i>Nature Immunology</i> , 2021 , 22, 942-944	19.1	0
161	The Naming of Memory T-Cell Subsets. <i>Cold Spring Harbor Perspectives in Biology</i> , 2021 , 13,	10.2	1
160	T Cell Memory: Understanding COVID-19. <i>Immunity</i> , 2021 , 54, 14-18	32.3	55
159	New Insights into the Immune System Using Dirty Mice. <i>Journal of Immunology</i> , 2020 , 205, 3-11	5.3	26
158	Sensing of ATP via the Purinergic Receptor P2RX7 Promotes CD8 Trm Cell Generation by Enhancing Their Sensitivity to the Cytokine TGF- β <i>Immunity</i> , 2020 , 53, 158-171.e6	32.3	25
157	The relationship between CD4+ follicular helper T cells and CD8+ resident memory T cells: sisters or distant cousins?. <i>International Immunology</i> , 2020 , 32, 583-587	4.9	4
156	ZipSeq: barcoding for real-time mapping of single cell transcriptomes. <i>Nature Methods</i> , 2020 , 17, 833-843	11.6	33
155	VISTA is a checkpoint regulator for naive T cell quiescence and peripheral tolerance. <i>Science</i> , 2020 , 367,	33.3	63
154	Danger-associated extracellular ATP counters MDSC therapeutic efficacy in acute GVHD. <i>Blood</i> , 2019 , 134, 1670-1682	2.2	33
153	NK Cell IL-10 Production Requires IL-15 and IL-10 Driven STAT3 Activation. <i>Frontiers in Immunology</i> , 2019 , 10, 2087	8.4	15
152	Microbial Exposure Enhances Immunity to Pathogens Recognized by TLR2 but Increases Susceptibility to Cytokine Storm through TLR4 Sensitization. <i>Cell Reports</i> , 2019 , 28, 1729-1743.e5	10.6	43

151	The Functional Requirement for CD69 in Establishment of Resident Memory CD8 T Cells Varies with Tissue Location. <i>Journal of Immunology</i> , 2019 , 203, 946-955	5.3	59
150	Self-Regulation of Memory CD8 T Cell Metabolism through Extracellular ATP Signaling. <i>Immunometabolism</i> , 2019 , 1,	4.1	7
149	ARTC2.2/P2RX7 Signaling during Cell Isolation Distorts Function and Quantification of Tissue-Resident CD8 T Cell and Invariant NKT Subsets. <i>Journal of Immunology</i> , 2019 , 202, 2153-2163	5.3	25
148	Myeloid cells activate iNKT cells to produce IL-4 in the thymic medulla. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 22262-22268	11.5	17
147	Embracing microbial exposure in mouse research. <i>Journal of Leukocyte Biology</i> , 2019 , 105, 73-79	6.5	13
146	Understanding Subset Diversity in T Cell Memory. <i>Immunity</i> , 2018 , 48, 214-226	32.3	215
145	Interleukin-15 Complex Treatment Protects Mice from Cerebral Malaria by Inducing Interleukin-10-Producing Natural Killer Cells. <i>Immunity</i> , 2018 , 48, 760-772.e4	32.3	30
144	Is a Human CD8 T-Cell Vaccine Possible, and if So, What Would It Take? CD8 T-Cell Vaccines: To B or Not to B?. <i>Cold Spring Harbor Perspectives in Biology</i> , 2018 , 10,	10.2	4
143	What Is the Predictive Value of Animal Models for Vaccine Efficacy in Humans? Reevaluating the Potential of Mouse Models for the Human Immune System. <i>Cold Spring Harbor Perspectives in Biology</i> , 2018 , 10,	10.2	12
142	The virtuous self-tolerance of virtual memory T cells. <i>EMBO Journal</i> , 2018 , 37,	13	4
141	The purinergic receptor P2RX7 directs metabolic fitness of long-lived memory CD8 T cells. <i>Nature</i> , 2018 , 559, 264-268	50.4	128
140	Retrieving short-term memories of flu. <i>Science Immunology</i> , 2017 , 2,	28	1
139	CD8 α intraepithelial lymphocytes arise from two main thymic precursors. <i>Nature Immunology</i> , 2017 , 18, 771-779	19.1	58
138	Hemodynamic Forces Sculpt Developing Heart Valves through a KLF2-WNT9B Paracrine Signaling Axis. <i>Developmental Cell</i> , 2017 , 43, 274-289.e5	10.2	70
137	Of Mice, Dirty Mice, and Men: Using Mice To Understand Human Immunology. <i>Journal of Immunology</i> , 2017 , 199, 383-388	5.3	131
136	IL-4 sensitivity shapes the peripheral CD8 $^+$ T cell pool and response to infection. <i>Journal of Experimental Medicine</i> , 2016 , 213, 1319-29	16.6	34
135	Lineage-Specific Effector Signatures of Invariant NKT Cells Are Shared amongst α , Innate Lymphoid, and Th Cells. <i>Journal of Immunology</i> , 2016 , 197, 1460-70	5.3	75
134	Late stages of T cell maturation in the thymus involve NF- κ B and tonic type I interferon signaling. <i>Nature Immunology</i> , 2016 , 17, 565-73	19.1	95

133	Spontaneous partial loss of the OT-I transgene. <i>Nature Immunology</i> , 2016 , 17, 471	19.1	3
132	Normalizing the environment recapitulates adult human immune traits in laboratory mice. <i>Nature</i> , 2016 , 532, 512-6	50.4	590
131	Sequential Infection with Common Pathogens Promotes Human-like Immune Gene Expression and Altered Vaccine Response. <i>Cell Host and Microbe</i> , 2016 , 19, 713-9	23.4	144
130	Innate memory T cells. <i>Advances in Immunology</i> , 2015 , 126, 173-213	5.6	75
129	TCR affinity for thymoproteasome-dependent positively selecting peptides conditions antigen responsiveness in CD8(+) T cells. <i>Nature Immunology</i> , 2015 , 16, 1069-76	19.1	51
128	Tissue-Specific Distribution of iNKT Cells Impacts Their Cytokine Response. <i>Immunity</i> , 2015 , 43, 566-78	32.3	167
127	Effective effector generation of CD8+ T cells and NK cells: A need for T-bet and ZEB-too. <i>Journal of Experimental Medicine</i> , 2015 , 212, 1990	16.6	4
126	The TCRs sensitivity to self peptide-MHC dictates the ability of naive CD8(+) T cells to respond to foreign antigens. <i>Nature Immunology</i> , 2015 , 16, 107-17	19.1	136
125	The transcription factor KLF2 restrains CD4+ T follicular helper cell differentiation. <i>Immunity</i> , 2015 , 42, 252-264	32.3	105
124	The self-obsession of T cells: how TCR signaling thresholds affect fate decisions and effector function. <i>Nature Immunology</i> , 2014 , 15, 815-23	19.1	171
123	An uncommon tail about the common chain. <i>Immunity</i> , 2014 , 40, 859-60	32.3	3
122	Antigen-specific culture of memory-like CD8 T cells for adoptive immunotherapy. <i>Cancer Immunology Research</i> , 2014 , 2, 839-45	12.5	4
121	Transcriptional downregulation of S1pr1 is required for the establishment of resident memory CD8+ T cells. <i>Nature Immunology</i> , 2013 , 14, 1285-93	19.1	459
120	T Cell Memory: without Prompting. <i>Journal of Immunology</i> , 2013 , 190, 4443-4	5.3	
119	Effector-like CD8+ T cells in the memory population mediate potent protective immunity. <i>Immunity</i> , 2013 , 38, 1250-60	32.3	176
118	Preexisting high frequencies of memory CD8+ T cells favor rapid memory differentiation and preservation of proliferative potential upon boosting. <i>Immunity</i> , 2013 , 39, 171-83	32.3	59
117	Murine thymic selection quantified using a unique method to capture deleted T cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 4679-84	11.5	120
116	Steady-state production of IL-4 modulates immunity in mouse strains and is determined by lineage diversity of iNKT cells. <i>Nature Immunology</i> , 2013 , 14, 1146-54	19.1	387

115	Cutting edge: The signals for the generation of T cell memory are qualitatively different depending on TCR ligand strength. <i>Journal of Immunology</i> , 2013 , 191, 5797-801	5.3	16
114	Thymoproteasome subunit- β T generates peptide-MHC complexes specialized for positive selection. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 6979-84	11.5	70
113	Virtual memory CD8 T cells display unique functional properties. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 13498-503	11.5	87
112	Kruppel-like factor 2 protects against ischemic stroke by regulating endothelial blood brain barrier function. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2013 , 304, H796-805	5.2	57
111	CD8 T cell quiescence revisited. <i>Trends in Immunology</i> , 2012 , 33, 224-30	14.4	47
110	TGF- β sensitivity restrains CD8+ T cell homeostatic proliferation by enforcing sensitivity to IL-7 and IL-15. <i>PLoS ONE</i> , 2012 , 7, e42268	3.7	22
109	Selection of self-reactive T cells in the thymus. <i>Annual Review of Immunology</i> , 2012 , 30, 95-114	34.7	232
108	Cutting edge: Kr \ddot{u} ppel-like factor 2 is required for phenotypic maintenance but not development of B1 B cells. <i>Journal of Immunology</i> , 2012 , 189, 3293-7	5.3	5
107	Immunology. Remembering to be tolerant. <i>Science</i> , 2012 , 335, 667-8	33.3	
106	Derivation and maintenance of virtual memory CD8 T cells. <i>Journal of Immunology</i> , 2012 , 188, 2516-23	5.3	95
105	Kr \ddot{u} ppel-like factors in lymphocyte biology. <i>Journal of Immunology</i> , 2012 , 188, 521-6	5.3	38
104	Cholera toxin activates nonconventional adjuvant pathways that induce protective CD8 T-cell responses after epicutaneous vaccination. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 2072-7	11.5	26
103	CD8 T cell memory: it takes all kinds. <i>Frontiers in Immunology</i> , 2012 , 3, 353	8.4	11
102	Keeping STATs on memory CD8+ T cells. <i>Immunity</i> , 2011 , 35, 663-5	32.3	10
101	Alternative memory in the CD8 T cell lineage. <i>Trends in Immunology</i> , 2011 , 32, 50-6	14.4	105
100	Kruppel-like factor 2 is required for trafficking but not quiescence in postactivated T cells. <i>Journal of Immunology</i> , 2011 , 186, 775-83	5.3	39
99	Kr \ddot{u} ppel-like factor 2 (KLF2) regulates B-cell reactivity, subset differentiation, and trafficking molecule expression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 716-21	11.5	67
98	Postselection thymocyte maturation and emigration are independent of IL-7 and ERK5. <i>Journal of Immunology</i> , 2011 , 186, 1343-7	5.3	19

97	T cells expressing the transcription factor PLZF regulate the development of memory-like CD8+ T cells. <i>Nature Immunology</i> , 2010 , 11, 709-16	19.1	192
96	IL-15 regulates both quantitative and qualitative features of the memory CD8 T cell pool. <i>Journal of Immunology</i> , 2010 , 184, 35-44	5.3	63
95	Self-specific CD8+ T cells maintain a semi-naive state following lymphopenia-induced proliferation. <i>Journal of Immunology</i> , 2010 , 184, 5604-11	5.3	17
94	Kr μ pel-like factor 2 regulates trafficking and homeostasis of gammadelta T cells. <i>Journal of Immunology</i> , 2010 , 184, 6060-6	5.3	37
93	CD4+CD25+Foxp3+ regulatory T cells optimize diversity of the conventional T cell repertoire during reconstitution from lymphopenia. <i>Journal of Immunology</i> , 2010 , 184, 4749-60	5.3	30
92	IL-2 complex treatment can protect naive mice from bacterial and viral infection. <i>Journal of Immunology</i> , 2010 , 185, 6584-90	5.3	28
91	Positive selection optimizes the number and function of MHCII-restricted CD4+ T cell clones in the naive polyclonal repertoire. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 11241-5	11.5	34
90	The antigen-specific CD8+ T cell repertoire in unimmunized mice includes memory phenotype cells bearing markers of homeostatic expansion. <i>Journal of Experimental Medicine</i> , 2009 , 206, 435-48	16.6	237
89	Epidermal Langerhans cells are not required for UV-induced immunosuppression. <i>Journal of Immunology</i> , 2009 , 183, 5548-53	5.3	35
88	Programming for CD8 T cell memory development requires IL-12 or type I IFN. <i>Journal of Immunology</i> , 2009 , 182, 2786-94	5.3	166
87	Self-class I MHC molecules support survival of naive CD8 T cells, but depress their functional sensitivity through regulation of CD8 expression levels. <i>Journal of Experimental Medicine</i> , 2009 , 206, 2253-69	16.6	59
86	Naive T cell homeostasis: from awareness of space to a sense of place. <i>Nature Reviews Immunology</i> , 2009 , 9, 823-32	36.5	277
85	KLF2 transcription-factor deficiency in T cells results in unrestrained cytokine production and upregulation of bystander chemokine receptors. <i>Immunity</i> , 2009 , 31, 122-30	32.3	157
84	Diversity in T cell memory: an embarrassment of riches. <i>Immunity</i> , 2009 , 31, 859-71	32.3	282
83	Immunology. A chronic need for IL-21. <i>Science</i> , 2009 , 324, 1525-6	33.3	32
82	Different T cell receptor signals determine CD8+ memory versus effector development. <i>Science</i> , 2009 , 323, 502-5	33.3	151
81	Roles of Kr μ pel-like Factors in Lymphocytes 2009 , 95-106		
80	The nature of the lymphopenic environment dictates protective function of homeostatic-memory CD8+ T cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 18484-9	11.5	31

79	Langerin expressing cells promote skin immune responses under defined conditions. <i>Journal of Immunology</i> , 2008 , 180, 4722-7	5.3	98
78	Selective regulation of CD8 effector T cell migration by the p110 gamma isoform of phosphatidylinositol 3-kinase. <i>Journal of Immunology</i> , 2008 , 180, 2081-8	5.3	60
77	Regulation of KLF2 in the Thymus. <i>FASEB Journal</i> , 2008 , 22, 346-346	0.9	
76	IL-15 is required for sustained lymphopenia-driven proliferation and accumulation of CD8 T cells. <i>Journal of Immunology</i> , 2007 , 179, 120-5	5.3	56
75	The CD8 T cell response to vaccinia virus exhibits site-dependent heterogeneity of functional responses. <i>International Immunology</i> , 2007 , 19, 733-43	4.9	16
74	Detuning CD8 T cells: down-regulation of CD8 expression, tetramer binding, and response during CTL activation. <i>Journal of Experimental Medicine</i> , 2007 , 204, 2667-77	16.6	86
73	Naive CD4(+) T cell frequency varies for different epitopes and predicts repertoire diversity and response magnitude. <i>Immunity</i> , 2007 , 27, 203-13	32.3	728
72	CD8(+) T cell differentiation: choosing a path through T-bet. <i>Immunity</i> , 2007 , 27, 180-2	32.3	15
71	The sialyltransferase ST3Gal-I is not required for regulation of CD8-class I MHC binding during T cell development. <i>Journal of Immunology</i> , 2006 , 176, 7421-30	5.3	13
70	T cells climb on board Blimp-1. <i>Trends in Immunology</i> , 2006 , 27, 349-51	14.4	2
69	The generation of protective memory-like CD8+ T cells during homeostatic proliferation requires CD4+ T cells. <i>Nature Immunology</i> , 2006 , 7, 475-81	19.1	176
68	Kruppel-like factor 2 regulates thymocyte and T-cell migration. <i>Nature</i> , 2006 , 442, 299-302	50.4	400
67	The timing of TCR alpha expression critically influences T cell development and selection. <i>Journal of Experimental Medicine</i> , 2005 , 202, 111-21	16.6	130
66	T cell homeostasis: keeping useful T cells alive and live T cells useful. <i>Seminars in Immunology</i> , 2005 , 17, 231-7	10.7	99
65	Central tolerance: learning self-control in the thymus. <i>Nature Reviews Immunology</i> , 2005 , 5, 772-82	36.5	487
64	Loss of CD8 and TCR binding to Class I MHC ligands following T cell activation. <i>International Immunology</i> , 2005 , 17, 1607-17	4.9	26
63	Characterizing the impact of CD8 antibodies on class I MHC multimer binding. <i>Journal of Immunology</i> , 2005 , 174, 3986-91	5.3	14
62	Characteristics of NK cell migration early after vaccinia infection. <i>Journal of Immunology</i> , 2005 , 175, 2152-3	5.3	24

61	Cutting edge: transpresentation of IL-15 by bone marrow-derived cells necessitates expression of IL-15 and IL-15R alpha by the same cells. <i>Journal of Immunology</i> , 2004 , 173, 6537-41	5.3	156
60	A role for CD28 in lymphopenia-induced proliferation of CD4 T cells. <i>Journal of Immunology</i> , 2004 , 173, 3909-15	5.3	47
59	Cutting edge: LFA-1 integrin-dependent T cell adhesion is regulated by both ag specificity and sensitivity. <i>Journal of Immunology</i> , 2004 , 173, 2222-6	5.3	27
58	The fourth way? Harnessing aggressive tendencies in the thymus. <i>Journal of Immunology</i> , 2004 , 173, 6515-20	5.3	75
57	Location of the epitope for an anti-CD8alpha antibody 53.6.7 which enhances CD8alpha-MHC class I interaction indicates antibody stabilization of a higher affinity CD8 conformation. <i>Immunology Letters</i> , 2004 , 93, 123-30	4.1	15
56	Thymocyte sensitivity and supramolecular activation cluster formation are developmentally regulated: a partial role for sialylation. <i>Journal of Immunology</i> , 2003 , 171, 4512-20	5.3	45
55	Distinct effects of STAT5 activation on CD4+ and CD8+ T cell homeostasis: development of CD4+CD25+ regulatory T cells versus CD8+ memory T cells. <i>Journal of Immunology</i> , 2003 , 171, 5853-64	5.3	171
54	Competition for self ligands restrains homeostatic proliferation of naive CD4 T cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 1185-90	11.5	103
53	Differential role for IL-7 in inducing lung Kruppel-like factor (Kruppel-like factor 2) expression by naive versus activated T cells. <i>International Immunology</i> , 2003 , 15, 1341-8	4.9	18
52	Receptor sensitivity: when T cells lose their sense of self. <i>Current Biology</i> , 2003 , 13, R239-41	6.3	20
51	Positive and negative selection of T cells. <i>Annual Review of Immunology</i> , 2003 , 21, 139-76	34.7	1178
50	In vivo survival and homeostatic proliferation of natural killer cells. <i>Journal of Experimental Medicine</i> , 2003 , 197, 967-76	16.6	188
49	Homeostatic expansion versus antigen-driven proliferation: common ends by different means?. <i>Microbes and Infection</i> , 2002 , 4, 531-7	9.3	31
48	Sweet & Sour: the impact of differential glycosylation on T cell responses. <i>Nature Immunology</i> , 2002 , 3, 903-10	19.1	220
47	Maintaining the norm: T-cell homeostasis. <i>Nature Reviews Immunology</i> , 2002 , 2, 547-56	36.5	488
46	A spontaneous CD8 T cell-dependent autoimmune disease to an antigen expressed under the human keratin 14 promoter. <i>Journal of Immunology</i> , 2002 , 169, 2141-7	5.3	49
45	Multiple choices: regulation of memory CD8 T cell generation and homeostasis by interleukin (IL)-7 and IL-15. <i>Journal of Experimental Medicine</i> , 2002 , 195, F49-52	16.6	127
44	Rare, structurally homologous self-peptides promote thymocyte positive selection. <i>Immunity</i> , 2002 , 17, 131-42	32.3	81

43	Il-12 enhances CD8 T cell homeostatic expansion. <i>Journal of Immunology</i> , 2001 , 166, 5515-21	5.3	90
42	A low affinity TCR ligand restores positive selection of CD8+ T cells in vivo. <i>Journal of Immunology</i> , 2001 , 166, 6602-7	5.3	33
41	Homeostatic expansion occurs independently of costimulatory signals. <i>Journal of Immunology</i> , 2001 , 167, 5664-8	5.3	103
40	The impact of duration versus extent of TCR occupancy on T cell activation: a revision of the kinetic proofreading model. <i>Immunity</i> , 2001 , 15, 59-70	32.3	197
39	CD8 binding to MHC class I molecules is influenced by T cell maturation and glycosylation. <i>Immunity</i> , 2001 , 15, 1051-61	32.3	152
38	Cutting edge: In situ tetramer staining of antigen-specific T cells in tissues. <i>Journal of Immunology</i> , 2000 , 165, 613-7	5.3	121
37	Interleukin-7 mediates the homeostasis of naïve and memory CD8 T cells in vivo. <i>Nature Immunology</i> , 2000 , 1, 426-32	19.1	1310
36	Critical role for CD8 in T cell receptor binding and activation by peptide/major histocompatibility complex multimers. <i>Journal of Experimental Medicine</i> , 2000 , 191, 335-46	16.6	224
35	Positive selection is limited by available peptide-dependent MHC conformations. <i>Journal of Immunology</i> , 2000 , 164, 3519-26	5.3	12
34	Role of 2CT cell receptor residues in the binding of self- and allo-major histocompatibility complexes. <i>Journal of Experimental Medicine</i> , 2000 , 191, 1355-64	16.6	48
33	Homeostatic expansion and phenotypic conversion of naïve T cells in response to self peptide/MHC ligands. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999 , 96, 13306-11	11.5	292
32	A divalent major histocompatibility complex/IgG1 fusion protein induces antigen-specific T cell activation in vitro and in vivo. <i>Cellular Immunology</i> , 1999 , 192, 54-62	4.4	10
31	Qualitative and quantitative differences in T cell receptor binding of agonist and antagonist ligands. <i>Immunity</i> , 1999 , 10, 227-37	32.3	189
30	Enhanced sensitivity for sequence determination of major histocompatibility complex class I peptides by membrane preconcentration-capillary electrophoresis-microspray-tandem mass spectrometry. <i>Electrophoresis</i> , 1998 , 19, 2207-12	3.6	36
29	T-cell selection. <i>Current Opinion in Immunology</i> , 1998 , 10, 214-9	7.8	135
28	Preselection thymocytes are more sensitive to T cell receptor stimulation than mature T cells. <i>Journal of Experimental Medicine</i> , 1998 , 188, 1867-74	16.6	174
27	T cell receptor antagonism in vivo, at last. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998 , 95, 14001-2	11.5	8
26	Identification of a naturally occurring ligand for thymic positive selection. <i>Immunity</i> , 1997 , 6, 389-99	32.3	156

25	Utility of membrane preconcentration-capillary electrophoresis-mass spectrometry in overcoming limited sample loading for analysis of biologically derived drug metabolites, peptides, and proteins. <i>Journal of the American Society for Mass Spectrometry</i> , 1997 , 8, 15-24	3.5	75
24	Strategy for isolating and sequencing biologically derived MHC class I peptides. <i>Journal of Chromatography A</i> , 1996 , 744, 273-8	4.5	40
23	Rapid loading of large sample volumes, analyte cleanup, and modified moving boundary transient isotachopheresis conditions for membrane preconcentration-capillary electrophoresis in small diameter capillaries. <i>Electrophoresis</i> , 1996 , 17, 1801-7	3.6	50
22	T-cell-receptor affinity and thymocyte positive selection. <i>Nature</i> , 1996 , 381, 616-20	50.4	542
21	T cell receptor (TCR) recognition of MHC class I variants: intermolecular second-site reversion provides evidence for peptide/MHC conformational variation. <i>Journal of Experimental Medicine</i> , 1996 , 184, 253-8	16.6	25
20	Options for TCR Interactions: TCR Agonists, Antagonists and Partial Agonists 1996 , 181-190		
19	Positive selection of thymocytes. <i>Annual Review of Immunology</i> , 1995 , 13, 93-126	34.7	512
18	T cell receptor antagonists and partial agonists. <i>Immunity</i> , 1995 , 2, 1-11	32.3	267
17	Strong agonist ligands for the T cell receptor do not mediate positive selection of functional CD8+ T cells. <i>Immunity</i> , 1995 , 3, 79-86	32.3	145
16	Selecting the T cell receptor repertoire. <i>Science</i> , 1994 , 264, 796-7	33.3	66
15	A thymic epithelial cell line induces both positive and negative selection in the thymus. <i>International Immunology</i> , 1994 , 6, 239-46	4.9	26
14	The ligand for positive selection of T lymphocytes in the thymus. <i>Current Opinion in Immunology</i> , 1994 , 6, 273-8	7.8	51
13	Specificity and flexibility in thymic selection. <i>Nature</i> , 1994 , 369, 750-2	50.4	197
12	T cell receptor antagonist peptides induce positive selection. <i>Cell</i> , 1994 , 76, 17-27	56.2	2206
11	Clone-specific T cell receptor antagonists of major histocompatibility complex class I-restricted cytotoxic T cells. <i>Journal of Experimental Medicine</i> , 1993 , 177, 1541-50	16.6	258
10	Variable binding affinities of listeriolysin O peptides for the H-2Kd class I molecule. <i>European Journal of Immunology</i> , 1993 , 23, 2005-10	6.1	31
9	Peptide-induced conformational changes in class I heavy chains alter major histocompatibility complex recognition. <i>Journal of Experimental Medicine</i> , 1992 , 176, 1757-61	16.6	109
8	Cloning and expression of class I major histocompatibility complex genes of the rat. <i>Journal of Experimental Medicine</i> , 1992 , 175, 1749-57	16.6	42

7	Chromosome 14 in B10.A(18R) mice is recombinant and includes Tcra-Va alleles. <i>Immunogenetics</i> , 1992 , 35, 190-8	3.2	3
6	Ham-2 corrects the class I antigen-processing defect in RMA-S cells. <i>Nature</i> , 1992 , 355, 647-9	50.4	272
5	Dissection of major histocompatibility complex (MHC) and T cell receptor contact residues in a Kb-restricted ovalbumin peptide and an assessment of the predictive power of MHC-binding motifs. <i>European Journal of Immunology</i> , 1992 , 22, 2663-7	6.1	113
4	Profound alteration in an alpha beta T-cell antigen receptor repertoire due to polymorphism in the first complementarity-determining region of the beta chain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991 , 88, 10267-71	11.5	24
3	Selective development of CD4+ T cells in transgenic mice expressing a class II MHC-restricted antigen receptor. <i>Nature</i> , 1989 , 341, 746-9	50.4	570
2	ZipSeq : Barcoding for Real-time Mapping of Single Cell Transcriptomes		4
1	The extracellular ATP receptor P2RX7 imprints a pro-memory transcriptional signature in effector CD8+ T cells		1