

# Ingemar If Petersson

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/1982987/ingemar-if-petersson-publications-by-year.pdf>  
**Version:** 2024-04-11

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.  
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

123 papers	4,771 citations	38 h-index	64 g-index
129 ext. papers	5,439 ext. citations	3.9 avg, IF	5.33 L-index

#	Paper	IF	Citations
123	Utilization of healthcare and prescription medicines after non-pharmacological interventions for depression - A 3-year register follow-up of an RCT in primary care.. <i>Preventive Medicine Reports</i> , <b>2022</b> , 25, 101658	2.6	1
122	Mortality over 14 years in MTX-refractory patients randomized to a strategy of addition of infliximab or sulfasalazine and hydroxychloroquine. <i>Rheumatology</i> , <b>2021</b> , 60, 2217-2222	3.9	
121	Evaluation of a multimodal pain rehabilitation programme in primary care based on clinical register data: a feasibility study. <i>Primary Health Care Research and Development</i> , <b>2020</b> , 21, e2	1.6	5
120	Long-term effects on function, health-related quality of life and work ability after structured physiotherapy including a workplace intervention. A secondary analysis of a randomised controlled trial (WorkUp) in primary care for patients with neck and/or back pain. <i>Scandinavian Journal of Primary Health Care</i> , <b>2020</b> , 38, 92-100	2.7	2
119	Educational inequalities in mortality associated with rheumatoid arthritis and other musculoskeletal disorders in Sweden. <i>BMC Musculoskeletal Disorders</i> , <b>2019</b> , 20, 83	2.8	4
118	Temporal trend and regional disparity in osteoarthritis hospitalisations in Sweden 1998-2015. <i>Scandinavian Journal of Public Health</i> , <b>2019</b> , 47, 53-60	3	6
117	The revised Bristol Rheumatoid Arthritis Fatigue measures and the Rheumatoid Arthritis Impact of Disease scale: validation in six countries. <i>Rheumatology</i> , <b>2018</b> , 57, 300-308	3.9	21
116	Income disparities in healthcare use remain after controlling for healthcare need: evidence from Swedish register data on psoriasis and psoriatic arthritis. <i>European Journal of Health Economics</i> , <b>2018</b> , 19, 447-462	3.6	6
115	Secular trends in sickness absence among Swedish patients with ankylosing spondylitis and psoriatic arthritis. <i>Rheumatology International</i> , <b>2018</b> , 38, 275-282	3.6	2
114	High and rising burden of hip and knee osteoarthritis in the Nordic region, 1990-2015. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2018</b> , 89, 177-183	4.3	36
113	Protocol for evaluating and implementing a pragmatic value-based healthcare management model for patients with inflammatory arthritis: a Danish population-based regional cohort and qualitative implementation study. <i>BMJ Open</i> , <b>2018</b> , 8, e023915	3	6
112	Can the STarT Back Tool predict health-related quality of life and work ability after an acute/subacute episode with back or neck pain? A psychometric validation study in primary care. <i>BMJ Open</i> , <b>2018</b> , 8, e021748	3	6
111	Secular trends in work disability and its relationship to musculoskeletal pain and mental health: a time-trend analysis using five cross-sectional surveys (2002-2010) in the general population. <i>Occupational and Environmental Medicine</i> , <b>2018</b> , 75, 877-883	2.1	8
110	Understanding Occupation, Sick Leave, and Disability Pension Due to Knee and Hip Osteoarthritis From a Sex Perspective. <i>Arthritis Care and Research</i> , <b>2017</b> , 69, 226-233	4.7	18
109	Predictors of work disability after start of anti-TNF therapy in a national cohort of Swedish patients with rheumatoid arthritis: does early anti-TNF therapy bring patients back to work?. <i>Annals of the Rheumatic Diseases</i> , <b>2017</b> , 76, 1245-1252	2.4	22
108	Severe Extraarticular Manifestations in a Community-based Cohort of Patients with Rheumatoid Arthritis: Risk Factors and Incidence in Relation to Treatment with Tumor Necrosis Factor Inhibitors. <i>Journal of Rheumatology</i> , <b>2017</b> , 44, 981-987	4.1	17
107	Sick leave in Sweden before and after total joint replacement in hip and knee osteoarthritis patients. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2017</b> , 88, 152-157	4.3	20

106	A Multi-Biomarker Disease Activity Score and the Choice of Second-Line Therapy in Early Rheumatoid Arthritis After Methotrexate Failure. <i>Arthritis and Rheumatology</i> , <b>2017</b> , 69, 953-963	9.5	22
105	Comparison of the Swedish STarT Back Screening Tool and the Short Form of the Fibro Musculoskeletal Pain Screening Questionnaire in patients with acute or subacute back and neck pain. <i>BMC Musculoskeletal Disorders</i> , <b>2017</b> , 18, 89	2.8	13
104	Referring to multimodal rehabilitation for patients with musculoskeletal disorders - a register study in primary health care. <i>BMC Health Services Research</i> , <b>2017</b> , 17, 15	2.9	5
103	Direct and indirect costs for systemic lupus erythematosus in Sweden. A nationwide health economic study based on five defined cohorts. <i>Seminars in Arthritis and Rheumatism</i> , <b>2016</b> , 45, 684-90	5.3	12
102	Does disease activity at start of biologic therapy influence work-loss in RA patients?. <i>Rheumatology</i> , <b>2016</b> , 55, 729-34	3.9	5
101	Changes in the anticitrullinated peptide antibody response in relation to therapeutic outcome in early rheumatoid arthritis: results from the SWEFOT trial. <i>Annals of the Rheumatic Diseases</i> , <b>2016</b> , 75, 356-61	2.4	23
100	Content validity of global measures for at-work productivity in patients with rheumatic diseases: an international qualitative study. <i>Rheumatology</i> , <b>2016</b> , 55, 1364-73	3.9	15
99	Back pain and health status in patients with clinically diagnosed ankylosing spondylitis, psoriatic arthritis and other spondyloarthritis: a cross-sectional population-based study. <i>BMC Musculoskeletal Disorders</i> , <b>2016</b> , 17, 106	2.8	14
98	Incremental Costs for Psoriasis and Psoriatic Arthritis in a Population-based Cohort in Southern Sweden: Is It All Psoriasis-attributable Morbidity?. <i>Journal of Rheumatology</i> , <b>2016</b> , 43, 640-7	4.1	10
97	Test-retest Reliability and Correlations of 5 Global Measures Addressing At-work Productivity Loss in Patients with Rheumatic Diseases. <i>Journal of Rheumatology</i> , <b>2016</b> , 43, 433-9	4.1	15
96	Economic Burden of Psoriasis and Potential Cost Offsets with Biologic Treatment: A Swedish Register Analysis. <i>Acta Dermato-Venereologica</i> , <b>2016</b> , 96, 651-7	2.2	11
95	Infliximab Versus Conventional Combination Treatment and Seven-Year Work Loss in Early Rheumatoid Arthritis: Results of a Randomized Swedish Trial. <i>Arthritis Care and Research</i> , <b>2016</b> , 68, 1758-1766	4.7	6
94	Predictors of presenteeism and activity impairment outside work in patients with spondyloarthritis. <i>Journal of Occupational Rehabilitation</i> , <b>2015</b> , 25, 288-95	3.6	20
93	Pretreatment multi-biomarker disease activity score and radiographic progression in early RA: results from the SWEFOT trial. <i>Annals of the Rheumatic Diseases</i> , <b>2015</b> , 74, 1102-9	2.4	76
92	Smoking is associated with a worse self-reported health status in patients with psoriatic arthritis: data from a Swedish population-based cohort. <i>Clinical Rheumatology</i> , <b>2015</b> , 34, 579-83	3.9	24
91	Factors affecting recall rate and false positive fraction in breast cancer screening with breast tomosynthesis - A statistical approach. <i>Breast</i> , <b>2015</b> , 24, 680-6	3.6	6
90	Work Disability in Early Systemic Sclerosis: A Longitudinal Population-based Cohort Study. <i>Journal of Rheumatology</i> , <b>2015</b> , 42, 1794-800	4.1	7
89	The risk of pneumococcal infections after immunization with pneumococcal conjugate vaccine compared to non-vaccinated inflammatory arthritis patients. <i>Scandinavian Journal of Rheumatology</i> , <b>2015</b> , 44, 271-9	1.9	11

88	Cost-effectiveness of infliximab versus conventional combination treatment in methotrexate-refractory early rheumatoid arthritis: 2-year results of the register-enriched randomised controlled SWEFOT trial. <i>Annals of the Rheumatic Diseases</i> , <b>2015</b> , 74, 1094-101	2.4	35
87	Development of patient-centred standards of care for osteoarthritis in Europe: the eumusc.net-project. <i>Annals of the Rheumatic Diseases</i> , <b>2015</b> , 74, 1145-9	2.4	35
86	The 21st-century landscape of adult fractures: cohort study of a complete adult regional population. <i>Journal of Bone and Mineral Research</i> , <b>2015</b> , 30, 535-42	6.3	36
85	Brief report: enhancement of patient recruitment in rheumatoid arthritis clinical trials using a multi-biomarker disease activity score as an inclusion criterion. <i>Arthritis and Rheumatology</i> , <b>2015</b> , 67, 2855-60	9.5	7
84	Comparison of non-radiographic axial spondyloarthritis and ankylosing spondylitis patients--baseline characteristics, treatment adherence, and development of clinical variables during three years of anti-TNF therapy in clinical practice. <i>Arthritis Research and Therapy</i> , <b>2015</b> , 17, 378	5.7	43
83	The association between antibody levels before and after 7-valent pneumococcal conjugate vaccine immunization and subsequent pneumococcal infection in chronic arthritis patients. <i>Arthritis Research and Therapy</i> , <b>2015</b> , 17, 124	5.7	12
82	Increased cause-specific mortality in patients with mild and severe psoriasis: a population-based Swedish register study. <i>Acta Dermato-Venereologica</i> , <b>2015</b> , 95, 809-15	2.2	21
81	Sickness absence from work among persons with new physician-diagnosed carpal tunnel syndrome: a population-based matched-cohort study. <i>PLoS ONE</i> , <b>2015</b> , 10, e0119795	3.7	15
80	International comparisons of the consultation prevalence of musculoskeletal conditions using population-based healthcare data from England and Sweden. <i>Annals of the Rheumatic Diseases</i> , <b>2014</b> , 73, 212-8	2.4	100
79	Current and future impact of osteoarthritis on health care: a population-based study with projections to year 2032. <i>Osteoarthritis and Cartilage</i> , <b>2014</b> , 22, 1826-32	6.2	202
78	Development of patient-centred standards of care for rheumatoid arthritis in Europe: the eumusc.net project. <i>Annals of the Rheumatic Diseases</i> , <b>2014</b> , 73, 902-5	2.4	69
77	Development of healthcare quality indicators for rheumatoid arthritis in Europe: the eumusc.net project. <i>Annals of the Rheumatic Diseases</i> , <b>2014</b> , 73, 906-8	2.4	35
76	Predictors of work disability during the first 3 years after diagnosis in a national rheumatoid arthritis inception cohort. <i>Annals of the Rheumatic Diseases</i> , <b>2014</b> , 73, 845-53	2.4	38
75	Socioeconomic status and the risk for being diagnosed with spondyloarthritis and chronic pain: a nested case-control study. <i>Rheumatology International</i> , <b>2014</b> , 34, 1291-8	3.6	12
74	Sociodemographic and disease-related factors are associated with patient-reported anxiety and depression in spondyloarthritis patients in the Swedish SpAScania cohort. <i>Clinical Rheumatology</i> , <b>2014</b> , 33, 1649-56	3.9	15
73	Sick leave patterns in common musculoskeletal disorders--a study of doctor prescribed sick leave. <i>BMC Musculoskeletal Disorders</i> , <b>2014</b> , 15, 176	2.8	35
72	Muscle strength in adolescent men and risk of cardiovascular disease events and mortality in middle age: a prospective cohort study. <i>BMC Medicine</i> , <b>2014</b> , 12, 62	11.4	60
71	Facilitators to implement standards of care for rheumatoid arthritis and osteoarthritis: the EUMUSC.NET project. <i>Annals of the Rheumatic Diseases</i> , <b>2014</b> , 73, 1545-8	2.4	8

70	Validity of diagnostic codes and prevalence of physician-diagnosed psoriasis and psoriatic arthritis in southern Sweden--a population-based register study. <i>PLoS ONE</i> , <b>2014</b> , 9, e98024	3.7	75
69	Nature-assisted rehabilitation for reactions to severe stress and/or depression in a rehabilitation garden: long-term follow-up including comparisons with a matched population-based reference cohort. <i>Journal of Rehabilitation Medicine</i> , <b>2014</b> , 46, 271-6	3.4	38
68	Which patients improve the most after arthritis rehabilitation? A study of predictors in patients with inflammatory arthritis in Northern Europe, the STAR-ETIC collaboration. <i>Journal of Rehabilitation Medicine</i> , <b>2014</b> , 46, 250-7	3.4	6
67	The risk for depression in patients with ankylosing spondylitis: a population-based cohort study. <i>Arthritis Research and Therapy</i> , <b>2014</b> , 16, 418	5.7	55
66	Worker productivity outcome measures: OMERACT filter evidence and agenda for future research. <i>Journal of Rheumatology</i> , <b>2014</b> , 41, 165-76	4.1	33
65	Evaluation of hand bone loss by digital X-ray radiogrammetry as a complement to clinical and radiographic assessment in early rheumatoid arthritis: results from the SWEFOT trial. <i>BMC Musculoskeletal Disorders</i> , <b>2013</b> , 14, 79	2.8	9
64	Promoting work ability in a structured national rehabilitation program in patients with musculoskeletal disorders: outcomes and predictors in a prospective cohort study. <i>BMC Musculoskeletal Disorders</i> , <b>2013</b> , 14, 57	2.8	14
63	Biological vs. conventional combination treatment and work loss in early rheumatoid arthritis: a randomized trial. <i>JAMA Internal Medicine</i> , <b>2013</b> , 173, 1407-14	11.5	28
62	Health care quality indicators on the management of rheumatoid arthritis and osteoarthritis: a literature review. <i>Rheumatology</i> , <b>2013</b> , 52, 382-90	3.9	20
61	Early increase in serum-COMP is associated with joint damage progression over the first five years in patients with rheumatoid arthritis. <i>BMC Musculoskeletal Disorders</i> , <b>2013</b> , 14, 229	2.8	33
60	Long-term work disability in patients with psoriatic arthritis treated with anti-tumour necrosis factor: a population-based regional Swedish cohort study. <i>Annals of the Rheumatic Diseases</i> , <b>2013</b> , 72, 1675-9	2.4	26
59	Addition of infliximab compared with addition of sulfasalazine and hydroxychloroquine to methotrexate in early rheumatoid arthritis: 2-year quality-of-life results of the randomised, controlled, SWEFOT trial. <i>Annals of the Rheumatic Diseases</i> , <b>2013</b> , 72, 1927-33	2.4	21
58	Healthcare consultation and sick leave before and after neck injury: a cohort study with matched population-based references. <i>BMJ Open</i> , <b>2013</b> , 3, e003172	3	10
57	Risk of sick leave and disability pension in working-age women and men with knee osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , <b>2013</b> , 72, 401-5	2.4	67
56	Work productivity in a population-based cohort of patients with spondyloarthritis. <i>Rheumatology</i> , <b>2013</b> , 52, 1708-14	3.9	32
55	Trends in the first decade of 21st century healthcare utilisation in a rheumatoid arthritis cohort compared with the general population. <i>Annals of the Rheumatic Diseases</i> , <b>2013</b> , 72, 1212-6	2.4	14
54	Disease course, outcome, and predictors of outcome in a population-based juvenile chronic arthritis cohort followed for 17 years. <i>Journal of Rheumatology</i> , <b>2013</b> , 40, 715-24	4.1	64
53	Goal-setting in multidisciplinary team care for patients with rheumatoid arthritis: an international multi-center evaluation of the contents using the International Classification of Functioning, Disability and Health as a reference. <i>Journal of Rehabilitation Medicine</i> , <b>2013</b> , 45, 888-99	3.4	15

52	Muscle strength in adolescent men and future musculoskeletal pain: a cohort study with 17 years of follow-up. <i>BMJ Open</i> , <b>2013</b> , 3,	3	9
51	Conventional combination treatment versus biological treatment in methotrexate-refractory early rheumatoid arthritis: 2 year follow-up of the randomised, non-blinded, parallel-group Swefot trial. <i>Lancet, The</i> , <b>2012</b> , 379, 1712-20	40	149
50	Differences in physical activity patterns in patients with spondylarthritis. <i>Arthritis Care and Research</i> , <b>2012</b> , 64, 1886-94	4.7	44
49	Population-based consultation patterns in patients with shoulder pain diagnoses. <i>BMC Musculoskeletal Disorders</i> , <b>2012</b> , 13, 238	2.8	57
48	Performance in physical education and health impairment 30 years later--a community based cohort study. <i>PLoS ONE</i> , <b>2012</b> , 7, e35718	3.7	1
47	Low back pain: epidemiology of consultations. <i>Arthritis Care and Research</i> , <b>2012</b> , 64, 1084-8	4.7	47
46	Sickness absence among cancer patients in the pre-diagnostic and the post-diagnostic phases of five common forms of cancer. <i>Supportive Care in Cancer</i> , <b>2012</b> , 20, 741-7	3.9	32
45	Sick leave in patients with ankylosing spondylitis before and after anti-TNF therapy: a population-based cohort study. <i>Rheumatology</i> , <b>2012</b> , 51, 243-9	3.9	30
44	Sickness absence in musculoskeletal disorders - patients' experiences of interactions with the social insurance agency and health care. A qualitative study. <i>BMC Public Health</i> , <b>2011</b> , 11, 107	4.1	24
43	Decrease in the incidence of total hip arthroplasties in patients with rheumatoid arthritis--results from a well defined population in south Sweden. <i>Arthritis Research and Therapy</i> , <b>2011</b> , 13, R67	5.7	56
42	Predictors of response to methotrexate in early DMARD naive rheumatoid arthritis: results from the initial open-label phase of the SWEFOT trial. <i>Annals of the Rheumatic Diseases</i> , <b>2011</b> , 70, 469-75	2.4	128
41	Validation of outcome measurement instruments used in a multidisciplinary rehabilitation intervention for patients with chronic inflammatory arthritis: linking of the International Classification of Functioning, Disability and Health, construct validity and responsiveness to change. <i>Journal of Rehabilitation Medicine</i> , <b>2011</b> , 43, 111-6	3.4	7
40	Patterns of background factors related to early RA patients' conceptions of the cause of their disease. <i>Clinical Rheumatology</i> , <b>2011</b> , 30, 347-52	3.9	3
39	Population-based estimates of common comorbidities and cardiovascular disease in ankylosing spondylitis. <i>Arthritis Care and Research</i> , <b>2011</b> , 63, 550-6	4.7	157
38	Incidence of physician-diagnosed carpal tunnel syndrome in the general population. <i>Archives of Internal Medicine</i> , <b>2011</b> , 171, 943-4		66
37	Prevalence of spondyloarthritis and its subtypes in southern Sweden. <i>Annals of the Rheumatic Diseases</i> , <b>2011</b> , 70, 943-8	2.4	94
36	How to develop patient-centered research: some perspectives based on surveys among people with rheumatic diseases in Scandinavia. <i>Physical Therapy</i> , <b>2010</b> , 90, 450-60	3.3	15
35	Decrease in sick leave among patients with rheumatoid arthritis in the first 12 months after start of treatment with tumour necrosis factor antagonists: a population-based controlled cohort study. <i>Annals of the Rheumatic Diseases</i> , <b>2010</b> , 69, 2131-6	2.4	30



34	Patient-reported 28 swollen and tender joint counts accurately represent RA disease activity and can be used to assess therapy responses at the group level. <i>Rheumatology</i> , <b>2010</b> , 49, 2098-103	3.9	21
33	Cost of illness from the public payers perspective in patients with ankylosing spondylitis in rheumatological care. <i>Journal of Rheumatology</i> , <b>2010</b> , 37, 2348-55	4.1	22
32	Prevalence and incidence of rheumatoid arthritis in southern Sweden 2008 and their relation to prescribed biologics. <i>Rheumatology</i> , <b>2010</b> , 49, 1563-9	3.9	96
31	Team-based rehabilitation improves long-term aerobic capacity and health-related quality of life in patients with chronic inflammatory arthritis. <i>Disability and Rehabilitation</i> , <b>2010</b> , 32, 1686-96	2.4	13
30	The grade in physical education in adolescence as predictor for musculoskeletal pain diagnoses three decades later. <i>Pain</i> , <b>2010</b> , 150, 414-419	8	8
29	Utility-based outcomes made easy: the number needed per quality-adjusted life year gained. An observational cohort study of tumor necrosis factor blockade in inflammatory arthritis from Southern Sweden. <i>Arthritis Care and Research</i> , <b>2010</b> , 62, 1399-406	4.7	16
28	Presence of peripheral arthritis and male sex predicting continuation of anti-tumor necrosis factor therapy in ankylosing spondylitis: an observational prospective cohort study from the South Swedish Arthritis Treatment Group Register. <i>Arthritis Care and Research</i> , <b>2010</b> , 62, 1362-9	4.7	69
27	Differences in longitudinal disease and treatment characteristics of patients with rheumatoid arthritis replying and not replying to a postal questionnaire. Experience from a biologics register in southern Sweden. <i>Journal of Rheumatology</i> , <b>2009</b> , 36, 1166-9	4.1	6
26	High disease activity disability burden and smoking predict severe extra-articular manifestations in early rheumatoid arthritis. <i>Rheumatology</i> , <b>2009</b> , 48, 416-20	3.9	50
25	Patients' conceptions of the cause of their rheumatoid arthritis: a qualitative study. <i>Musculoskeletal Care</i> , <b>2009</b> , 7, 243-55	1.6	7
24	Patients with ankylosing spondylitis have increased sick leave--a registry-based case-control study over 7 yrs. <i>Rheumatology</i> , <b>2009</b> , 48, 289-92	3.9	19
23	Cooperation between gatekeepers in sickness insurance - the perspective of social insurance officers. A qualitative study. <i>BMC Health Services Research</i> , <b>2008</b> , 8, 231	2.9	14
22	Association of STAT4 with rheumatoid arthritis: a replication study in three European populations. <i>Arthritis and Rheumatism</i> , <b>2008</b> , 58, 1974-80		85
21	Research in hand osteoarthritis: time for reappraisal and demand for new strategies. An opinion paper. <i>Annals of the Rheumatic Diseases</i> , <b>2007</b> , 66, 1157-61	2.4	38
20	Treatment with TNF blockers and mortality risk in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , <b>2007</b> , 66, 670-5	2.4	148
19	Serum levels of cartilage oligomeric matrix protein (COMP) increase temporarily after physical exercise in patients with knee osteoarthritis. <i>BMC Musculoskeletal Disorders</i> , <b>2006</b> , 7, 98	2.8	81
18	Association of HLA-C3 and smoking with vasculitis in patients with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , <b>2006</b> , 54, 2776-83		51
17	Analysis of IRF5 gene functional polymorphisms in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , <b>2006</b> , 54, 3815-9		46

16	How do middle-aged patients conceive exercise as a form of treatment for knee osteoarthritis? <i>Disability and Rehabilitation</i> , <b>2006</b> , 28, 51-9	2.4	44
15	A nurse-led rheumatology clinic's impact on empowering patients with rheumatoid arthritis: A qualitative study. <i>Australian Journal of Cancer Nursing</i> , <b>2006</b> , 8, 133-9	1.9	61
14	Team care. Traditions and new trends. <i>Journal of Rheumatology</i> , <b>2006</b> , 33, 1895-6	4.1	5
13	Design issues and priorities in team and nonpharmacological arthritis care research. <i>Journal of Rheumatology</i> , <b>2006</b> , 33, 1904-7	4.1	11
12	Revision in previously satisfied knee arthroplasty patients is the result of their call on the physician, not on pre-planned follow-up: a retrospective study of 181 patients who underwent revision within 2 years. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2005</b> , 76, 785-90	4.3	3
11	The impact of HLA-DRB1 genes on extra-articular disease manifestations in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , <b>2005</b> , 7, R1386-93	5.7	66
10	Evolution of team care and evaluation of effectiveness. <i>Current Opinion in Rheumatology</i> , <b>2005</b> , 17, 160-3	3.3	25
9	Six-week high-intensity exercise program for middle-aged patients with knee osteoarthritis: a randomized controlled trial [ISRCTN20244858]. <i>BMC Musculoskeletal Disorders</i> , <b>2005</b> , 6, 27	2.8	67
8	Treatment with tumor necrosis factor blockers is associated with a lower incidence of first cardiovascular events in patients with rheumatoid arthritis. <i>Journal of Rheumatology</i> , <b>2005</b> , 32, 1213-8	4.1	334
7	Health status as measured by SF-36 reflects changes and predicts outcome in chronic musculoskeletal pain: a 3-year follow up study in the general population. <i>Pain</i> , <b>2004</b> , 108, 115-23	8	110
6	Validation of the Rheumatoid and Arthritis Outcome Score (RAOS) for the lower extremity. <i>Health and Quality of Life Outcomes</i> , <b>2003</b> , 1, 55	3	36
5	Osteoarthritis of the peripheral joints. <i>Best Practice and Research in Clinical Rheumatology</i> , <b>2002</b> , 16, 741-60	5.3	62
4	Chronic widespread pain: a three year followup of pain distribution and risk factors. <i>Journal of Rheumatology</i> , <b>2002</b> , 29, 818-25	4.1	112
3	Bone scan and serum markers of bone and cartilage in patients with knee pain and osteoarthritis. <i>Osteoarthritis and Cartilage</i> , <b>1998</b> , 6, 33-9	6.2	67
2	Radiographic osteoarthritis of the knee classified by the Ahlbäck and Kellgren & Lawrence systems for the tibiofemoral joint in people aged 35-54 years with chronic knee pain. <i>Annals of the Rheumatic Diseases</i> , <b>1997</b> , 56, 493-6	2.4	180
1	Changes in serum levels of cartilage and bone markers in early osteoarthritis of the knee. <i>Acta Orthopaedica</i> , <b>1995</b> , 66, 144-145		4