Roberto Tuberosa

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1985320/publications.pdf

Version: 2024-02-01

162 16,865 papers citations

23567 58 h-index 124 g-index

175 all docs 175
docs citations

175 times ranked

11657 citing authors

| # | Article | IF | CITATIONS |
|----|--|--------------------|---------------|
| 1 | Genome Wide Association Study Uncovers the QTLome for Osmotic Adjustment and Related Drought Adaptive Traits in Durum Wheat. Genes, 2022, 13, 293. | 2.4 | 12 |
| 2 | Cloning the barley <i>nec3</i> disease lesion mimic mutant using complementation by sequencing. Plant Genome, 2022, , e20187. | 2.8 | 0 |
| 3 | Sequence-Based Marker Assisted Selection in Wheat. , 2022, , 513-538. | | 3 |
| 4 | Editorial: Proceedings of FSTP3 Congress–A Sustainable Durum Wheat Chain for Food Security and Healthy Lives. Frontiers in Plant Science, 2021, 12, 675510. | 3.6 | 0 |
| 5 | Wheat root systems as a breeding target for climate resilience. Theoretical and Applied Genetics, 2021, 134, 1645-1662. | 3.6 | 74 |
| 6 | Highâ€throughput field phenotyping reveals genetic variation in photosynthetic traits in durum wheat under drought. Plant, Cell and Environment, 2021, 44, 2858-2878. | 5.7 | 12 |
| 7 | ⟨i>ENHANCED GRAVITROPISM 2 i>encodes a STERILE ALPHA MOTIFâ€"containing protein that controls root growth angle in barley and wheat. Proceedings of the National Academy of Sciences of the United States of America, 2021, 118, . | 7.1 | 32 |
| 8 | Back Cover Image. Plant, Cell and Environment, 2021, 44, . | 5.7 | 0 |
| 9 | Breeding customâ€designed crops for improved drought adaptation. Genetics & Genomics Next, 2021, 2, e202100017. | 1.5 | 48 |
| 10 | Genome-wide association analysis unveils novel QTLs for seminal root system architecture traits in Ethiopian durum wheat. BMC Genomics, 2021, 22, 20. | 2.8 | 33 |
| 11 | Resequencing of 145 Landmark Cultivars Reveals Asymmetric Sub-genome Selection and Strong Founder Genotype Effects on Wheat Breeding in China. Molecular Plant, 2020, 13, 1733-1751. | 8.3 | 129 |
| 12 | Genomic Regions Associated with the Control of Flowering Time in Durum Wheat. Plants, 2020, 9, 1628. | 3.5 | 15 |
| 13 | The Global Durum Wheat Panel (GDP): An International Platform to Identify and Exchange Beneficial Alleles. Frontiers in Plant Science, 2020, 11, 569905. | 3.6 | 44 |
| 14 | Carbon Isotope Composition and the NDVI as Phenotyping Approaches for Drought Adaptation in Durum Wheat: Beyond Trait Selection. Agronomy, 2020, 10, 1679. | 3.0 | 4 |
| 15 | Quantitative trait loci for agronomic traits in tetraploid wheat for enhancing grain yield in Kazakhstan environments. PLoS ONE, 2020, 15, e0234863. | 2.5 | 19 |
| 16 | Genome-wide association mapping for grain shape and color traits in Ethiopian durum wheat (Triticum) Tj ETQq | 0 0 <u>0 rg</u> BT | '/Oygrlock 10 |
| 17 | Carotenoid Pigment Content in Durum Wheat (Triticum turgidum L. var durum): An Overview of Quantitative Trait Loci and Candidate Genes. Frontiers in Plant Science, 2019, 10, 1347. | 3.6 | 59 |
| 18 | Multi-Trait, Multi-Environment Genomic Prediction of Durum Wheat With Genomic Best Linear Unbiased Predictor and Deep Learning Methods. Frontiers in Plant Science, 2019, 10, 1311. | 3.6 | 47 |

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 19 | Reduced response diversity does not negatively impact wheat climate resilience. Proceedings of the National Academy of Sciences of the United States of America, 2019, 116, 10623-10624. | 7.1 | 11 |
| 20 | Marker-Assisted Breeding in Crops. , 2019, , 453-475. | | 0 |
| 21 | Durum wheat genome highlights past domestication signatures and future improvement targets. Nature Genetics, 2019, 51, 885-895. | 21.4 | 576 |
| 22 | Genomic tools for durum wheat breeding: de novo assembly of Svevo transcriptome and SNP discovery in elite germplasm. BMC Genomics, 2019, 20, 278. | 2.8 | 7 |
| 23 | Exploring and exploiting the genetic variation of Fusarium head blight resistance for genomic-assisted breeding in the elite durum wheat gene pool. Theoretical and Applied Genetics, 2019, 132, 969-988. | 3.6 | 57 |
| 24 | Asparagine synthetase genes (AsnS1 and AsnS2) in durum wheat: structural analysis and expression under nitrogen stress. Euphytica, 2018, 214, 1. | 1.2 | 21 |
| 25 | Dehydration survival of crop plants and its measurement. Journal of Experimental Botany, 2018, 69, 975-981. | 4.8 | 43 |
| 26 | Yield of chromosomally engineered durum wheat-Thinopyrum ponticum recombinant lines in a range of contrasting rain-fed environments. Field Crops Research, 2018, 228, 147-157. | 5.1 | 11 |
| 27 | Comparative Aerial and Ground Based High Throughput Phenotyping for the Genetic Dissection of NDVI as a Proxy for Drought Adaptive Traits in Durum Wheat. Frontiers in Plant Science, 2018, 9, 893. | 3.6 | 117 |
| 28 | Progress in understanding drought tolerance: from alleles to cropping systems. Journal of Experimental Botany, 2018, 69, 3175-3179. | 4.8 | 40 |
| 29 | Marker-Assisted Breeding in Crops. , 2018, , 1-24. | | 0 |
| 30 | Identification of Early Represented Gluten Proteins during Durum Wheat Grain Development. Journal of Agricultural and Food Chemistry, 2017, 65, 3242-3250. | 5.2 | 28 |
| 31 | Genome-wide association mapping for seedling and field resistance to Puccinia striiformis f. sp. tritici in elite durum wheat. Theoretical and Applied Genetics, 2017, 130, 649-667. | 3.6 | 71 |
| 32 | Wild emmer genome architecture and diversity elucidate wheat evolution and domestication. Science, 2017, 357, 93-97. | 12.6 | 781 |
| 33 | Leveraging plant genomics for better and healthier food. Current Opinion in Food Science, 2017, 16, 100-105. | 8.0 | 5 |
| 34 | <i>In vitro</i> physical mutagenesis of giant reed (<i>Arundo donax</i> L.). GCB Bioenergy, 2017, 9, 1380-1389. | 5.6 | 12 |
| 35 | Genome-wide association mapping reveals a rich genetic architecture of stripe rust resistance loci in emmer wheat (Triticum turgidum ssp. dicoccum). Theoretical and Applied Genetics, 2017, 130, 2249-2270. | 3.6 | 80 |
| 36 | Improving crop performance under drought – cross-fertilization of disciplines. Journal of Experimental Botany, 2017, 68, 1393-1398. | 4.8 | 8 |

3

| # | Article | IF | CITATIONS |
|----|--|-----------|----------------|
| 37 | Novel Sources of Stripe Rust Resistance Identified by Genome-Wide Association Mapping in Ethiopian Durum Wheat (Triticum turgidum ssp. durum). Frontiers in Plant Science, 2017, 8, 774. | 3.6 | 66 |
| 38 | Extending the Marker × Environment Interaction Model for Genomicâ€Enabled Prediction and Genomeâ€Wide Association Analysis in Durum Wheat. Crop Science, 2016, 56, 2193-2209. | 1.8 | 101 |
| 39 | A multiparental cross population for mapping <scp>QTL</scp> for agronomic traits in durum wheat (<i><scp>T</scp>riticum turgidum</i> >ssp. <i>durum</i>). Plant Biotechnology Journal, 2016, 14, 735-748. | 8.3 | 121 |
| 40 | Global agricultural intensification during climate change: a role for genomics. Plant Biotechnology Journal, 2016, 14, 1095-1098. | 8.3 | 221 |
| 41 | Genome-wide analysis of yield in Europe: allelic effects as functions of drought and heat scenarios. Plant Physiology, 2016, 172, pp.00621.2016. | 4.8 | 140 |
| 42 | Two major quantitative trait loci controlling the number of seminal roots in maize co-map with the root developmental genes <i>rtcs</i> and <i>rum1</i> . Journal of Experimental Botany, 2016, 67, 1149-1159. | 4.8 | 40 |
| 43 | Prioritizing quantitative trait loci for root system architecture in tetraploid wheat. Journal of Experimental Botany, 2016, 67, 1161-1178. | 4.8 | 206 |
| 44 | Genetic variation for aerenchyma and other root anatomical traits in durum wheat (Triticum durum) Tj ETQq0 0 0 | rgBT /Ove | erlock 10 Tf ! |
| 45 | Yield QTLome distribution correlates with gene density in maize. Plant Science, 2016, 242, 300-309. | 3.6 | 61 |
| 46 | The Genetic Basis of Composite Spike Form in Barley and â€~Miracle-Wheat'. Genetics, 2015, 201, 155-165. | 2.9 | 109 |
| 47 | Application of genomics-assisted breeding for generation of climate resilient crops: progress and prospects. Frontiers in Plant Science, 2015, 6, 563. | 3.6 | 243 |
| 48 | The crop QTLome comes of age. Current Opinion in Biotechnology, 2015, 32, 179-185. | 6.6 | 122 |
| 49 | A highâ€density, <scp>SNP</scp> â€based consensus map of tetraploid wheat as a bridge to integrate durum and bread wheat genomics and breeding. Plant Biotechnology Journal, 2015, 13, 648-663. | 8.3 | 386 |
| 50 | Genomics Approaches to Dissect the Genetic Basis of Drought Resistance in Durum Wheat., 2015,, 213-223. | | 2 |
| 51 | Association Mapping Reveals Novel Stem Rust Resistance Loci in Durum Wheat at the Seedling Stage. Plant Genome, 2014, 7, plantgenome2013.08.0026. | 2.8 | 67 |
| 52 | Two decades of InterDrought conferences: are we bridging the genotype-to-phenotype gap?. Journal of Experimental Botany, 2014, 65, 6137-6139. | 4.8 | 13 |
| 53 | A MITE Transposon Insertion Is Associated with Differential Methylation at the Maize Flowering Time QTL <i>Vgt1</i> . G3: Genes, Genomes, Genetics, 2014, 4, 805-812. | 1.8 | 93 |
| 54 | Association mapping for root architectural traits in durum wheat seedlings as related to agronomic performance. Molecular Breeding, 2014, 34, 1629-1645. | 2.1 | 115 |

| # | Article | lF | CITATIONS |
|----|---|------|-----------|
| 55 | Characterization of polyploid wheat genomic diversity using a highâ€density 90Â000 single nucleotide polymorphism array. Plant Biotechnology Journal, 2014, 12, 787-796. | 8.3 | 1,828 |
| 56 | QTL dissection of yield components and morpho-physiological traits in a durum wheat elite population tested in contrasting thermo-pluviometric conditions. Crop and Pasture Science, 2014, 65, 80. | 1.5 | 79 |
| 57 | Genomics of plant genetic resources: a gateway to a new era of global food security. Plant Genetic Resources: Characterisation and Utilisation, 2014, 12, S2-S5. | 0.8 | 4 |
| 58 | Durum wheat genomics comes of age. Molecular Breeding, 2014, 34, 1527-1530. | 2.1 | 23 |
| 59 | A consensus framework map of durum wheat (Triticum durum Desf.) suitable for linkage disequilibrium analysis and genome-wide association mapping. BMC Genomics, 2014, 15, 873. | 2.8 | 85 |
| 60 | Strategies to increase the yield and yield stability of crops under drought – are we making progress?. Functional Plant Biology, 2014, 41, 1199. | 2.1 | 32 |
| 61 | New Starch Phenotypes Produced by TILLING in Barley. PLoS ONE, 2014, 9, e107779. | 2.5 | 59 |
| 62 | Systems Responses to Progressive Water Stress in Durum Wheat. PLoS ONE, 2014, 9, e108431. | 2.5 | 52 |
| 63 | Development of COS-SNP and HRM markers for high-throughput and reliable haplotype-based detection of Lr14a in durum wheat (Triticum durum Desf.). Theoretical and Applied Genetics, 2013, 126, 1077-1101. | 3.6 | 43 |
| 64 | Searching for novel sources of field resistance to Ug99 and Ethiopian stem rust races in durum wheat via association mapping. Theoretical and Applied Genetics, 2013, 126, 1237-1256. | 3.6 | 116 |
| 65 | Sequenceâ€based SNP genotyping in durum wheat. Plant Biotechnology Journal, 2013, 11, 809-817. | 8.3 | 63 |
| 66 | Marker-Assisted breeding/breed, see also animal breeding marker-assisted Breeding breeding/breed, see also animal breeding in Crops. , 2013, , 1158-1181. | | 1 |
| 67 | Phenotyping for drought tolerance of crops in the genomics era. Frontiers in Physiology, 2012, 3, 347. | 2.8 | 448 |
| 68 | Biotechnology for enhancing plant production and food quality: IBS 2010 part III. Journal of Biotechnology, 2012, 159, 249-250. | 3.8 | 0 |
| 69 | Can genomics boost productivity of orphan crops?. Nature Biotechnology, 2012, 30, 1172-1176. | 17.5 | 248 |
| 70 | Biotechnology for a more sustainable world. Biotechnology Advances, 2012, 30, 931-932. | 11.7 | 12 |
| 71 | Genetic analysis of Soil-Borne Cereal Mosaic Virus response in durum wheat: evidence for the role of the major quantitative trait locus QSbm.ubo-2BS and of minor quantitative trait loci. Molecular Breeding, 2012, 29, 973-988. | 2.1 | 11 |
| 72 | Genomics of Root Architecture and Functions in Maize. , 2011, , 179-204. | | 14 |

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 73 | Resistance to Soil-borne cereal mosaic virus in durum wheat is controlled by a major QTL on chromosome arm 2BS and minor loci. Theoretical and Applied Genetics, 2011, 123, 527-544. | 3.6 | 25 |
| 74 | High-throughput SNP discovery and genotyping in durum wheat (Triticum durum Desf.). Theoretical and Applied Genetics, 2011, 123, 555-569. | 3.6 | 120 |
| 75 | Genetic dissection of maize phenology using an intraspecific introgression library. BMC Plant Biology, 2011, 11, 4. | 3.6 | 63 |
| 76 | Genomics of plant genetic resources: an introduction. Plant Genetic Resources: Characterisation and Utilisation, 2011, 9, 151-154. | 0.8 | 18 |
| 77 | Starch metabolism mutants in barley: A TILLING approach. Plant Genetic Resources: Characterisation and Utilisation, 2011, 9, 170-173. | 0.8 | 7 |
| 78 | Association mapping in durum wheat grown across a broad range of water regimes. Journal of Experimental Botany, 2011, 62, 409-438. | 4.8 | 270 |
| 79 | Identification of root morphology mutants in barley. Plant Genetic Resources: Characterisation and Utilisation, 2011, 9, 357-360. | 0.8 | 6 |
| 80 | Association mapping of leaf rust response in durum wheat. Molecular Breeding, 2010, 26, 189-228. | 2.1 | 86 |
| 81 | Dissection and modelling of abiotic stress tolerance in plants. Current Opinion in Plant Biology, 2010, 13, 206-212. | 7.1 | 281 |
| 82 | TILLMore: a Forward- and a Reverse-Genetics Resource for the Identification of Root Morphology-Related Mutants. Journal of Biotechnology, 2010, 150, 497-497. | 3.8 | 0 |
| 83 | Starch Metabolism Mutants in Barley: a TILLING Approach. Journal of Biotechnology, 2010, 150, 497-498. | 3.8 | 0 |
| 84 | Fine mapping approaches of two major QTLs for yield in durum wheat. Journal of Biotechnology, 2010, 150, 501-502. | 3.8 | 1 |
| 85 | 1 out of 27—European politicians score poorly in agbiotech. Nature Biotechnology, 2010, 28, 551-552. | 17.5 | 1 |
| 86 | Understanding the relationships between genetic and phenotypic structures of a collection of elite durum wheat accessions. Field Crops Research, 2010, 119, 91-105. | 5.1 | 54 |
| 87 | Virulence Phenotypes and Molecular Genotypes in Collections of Puccinia triticina from Italy. Plant Disease, 2010, 94, 420-424. | 1.4 | 26 |
| 88 | Characterization of root-yield-1.06, a major constitutive QTL for root and agronomic traits in maize across water regimes. Journal of Experimental Botany, 2010, 61, 3553-3562. | 4.8 | 79 |
| 89 | Genetic and genomic dissection of maize root development and architecture. Current Opinion in Plant Biology, 2009, 12, 172-177. | 7.1 | 230 |
| 90 | Genome studies and molecular geneticsâ€"from sequence to crops: genomics comes of age. Current Opinion in Plant Biology, 2009, 12, 103-106. | 7.1 | 23 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Genomics of Tolerance to Abiotic Stress in the Triticeae. , 2009, , 481-558. | | 8 |
| 92 | QTL for Agronomic Traits in Maize Production. , 2009, , 501-541. | | 28 |
| 93 | An integrated DArT-SSR linkage map of durum wheat. Molecular Breeding, 2008, 22, 629-648. | 2.1 | 97 |
| 94 | A major QTL for durable leaf rust resistance widely exploited in durum wheat breeding programs maps on the distal region of chromosome arm 7BL. Theoretical and Applied Genetics, 2008, 117, 1225-1240. | 3.6 | 59 |
| 95 | Vivipary as a tool to analyze late embryogenic events in maize. Heredity, 2008, 101, 465-470. | 2.6 | 23 |
| 96 | TILLMore, a resource for the discovery of chemically induced mutants in barley. Plant Biotechnology Journal, 2008, 6, 477-485. | 8.3 | 177 |
| 97 | Translational research impacting on crop productivity in drought-prone environments. Current Opinion in Plant Biology, 2008, 11, 171-179. | 7.1 | 324 |
| 98 | Principles and practices of plant penomics. Volume 1. Genome mapping. Annals of Botany, 2008, 102, 879-880. | 2.9 | 0 |
| 99 | Quantitative Trait Loci and Crop Performance under Abiotic Stress: Where Do We Stand?: Table I Plant Physiology, 2008, 147, 469-486. | 4.8 | 518 |
| 100 | Quantitative Trait Loci for Grain Yield and Adaptation of Durum Wheat (<i>Triticum durum</i> Desf.) Across a Wide Range of Water Availability. Genetics, 2008, 178, 489-511. | 2.9 | 397 |
| 101 | Conserved noncoding genomic sequences associated with a flowering-time quantitative trait locus in maize. Proceedings of the National Academy of Sciences of the United States of America, 2007, 104, 11376-11381. | 7.1 | 536 |
| 102 | Genomeâ€wide Approaches to Investigate and Improve Maize Response to Drought. Crop Science, 2007, 47, S-120. | 1.8 | 48 |
| 103 | Root system architecture: opportunities and constraints for genetic improvement of crops. Trends in Plant Science, 2007, 12, 474-481. | 8.8 | 608 |
| 104 | Cloning Qtls in Plants. , 2007, , 207-225. | | 33 |
| 105 | Relationships among durum wheat accessions. II. A comparison of molecular and pedigree information. Genome, 2007, 50, 385-399. | 2.0 | 18 |
| 106 | Relationships among durum wheat accessions. I. Comparative analysis of SSR, AFLP, and phenotypic data. Genome, 2007, 50, 373-384. | 2.0 | 37 |
| 107 | Improving water use efficiency in Mediterranean agriculture: what limits the adoption of new technologies?. Annals of Applied Biology, 2007, 150, 157-162. | 2.5 | 49 |
| 108 | Genetic dissection of seminal root architecture in elite durum wheat germplasm. Annals of Applied Biology, 2007, 151, 291-305. | 2.5 | 115 |

| # | Article | lF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Dissecting Qtls For Tolerance to Drought and Salinity. , 2007, , 381-411. | | 9 |
| 110 | Genomics-Assisted Crop Improvement: An Overview. , 2007, , 1-12. | | 18 |
| 111 | Nucleotide-binding site (NBS) profiling of genetic diversity in durum wheat. Genome, 2006, 49, 1473-1480. | 2.0 | 26 |
| 112 | A panel of elite accessions of durum wheat (Triticum durum Desf.) suitable for association mapping studies. Plant Genetic Resources: Characterisation and Utilisation, 2006, 4, 79-85. | 0.8 | 54 |
| 113 | Genomics-based approaches to improve drought tolerance of crops. Trends in Plant Science, 2006, 11, 405-412. | 8.8 | 478 |
| 114 | Validation of Quantitative Trait Loci for Multiple Disease Resistance in Barley Using Advanced Backcross Lines Developed with a Wild Barley. Crop Science, 2006, 46, 1179-1186. | 1.8 | 47 |
| 115 | Distribution of \hat{I}^2 -amylase I haplotypes among European cultivated barleys. Molecular Breeding, 2006, 18, 341-354. | 2.1 | 20 |
| 116 | Barley transcript profiles under dehydration shock and drought stress treatments: a comparative analysis. Journal of Experimental Botany, 2006, 58, 229-240. | 4.8 | 201 |
| 117 | Root-ABA1 QTL affects root lodging, grain yield, and other agronomic traits in maize grown under well-watered and water-stressed conditions. Journal of Experimental Botany, 2006, 58, 319-326. | 4.8 | 89 |
| 118 | Genetic evaluation of root complexity in maize. Acta Agronomica Hungarica: an International Multidisciplinary Journal in Agricultural Science, 2006, 54, 291-303. | 0.2 | 25 |
| 119 | A high-density genetic map of hexaploid wheat (Triticum aestivum L.) from the cross Chinese SpringÂÃ—ÂŚQ1 and its use to compare QTLs for grain yield across a range of environments. Theoretical and Applied Genetics, 2005, 110, 865-880. | 3.6 | 437 |
| 120 | Population structure and long-range linkage disequilibrium in a durum wheat elite collection. Molecular Breeding, 2005, 15, 271-290. | 2.1 | 212 |
| 121 | Validation and characterization of a major QTL affecting leaf ABA concentration in maize. Molecular Breeding, 2005, 15, 291-303. | 2.1 | 59 |
| 122 | Root-ABA1, a major constitutive QTL, affects maize root architecture and leaf ABA concentration at different water regimes. Journal of Experimental Botany, 2005, 56, 3061-3070. | 4.8 | 165 |
| 123 | To clone or not to clone plant QTLs: present and future challenges. Trends in Plant Science, 2005, 10, 297-304. | 8.8 | 487 |
| 124 | Genotype and phenotype associations with drought tolerance in barley tested in North Africa. Annals of Applied Biology, 2004, 144, 157-168. | 2.5 | 108 |
| 125 | Identification of Hordeum spontaneum QTL alleles improving field performance of barley grown under rainfed conditions. Annals of Applied Biology, 2004, 144, 309-319. | 2.5 | 101 |
| 126 | QTLs and Genes for Tolerance to Abiotic Stress in Cereals. , 2004, , 253-315. | | 9 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 127 | Extension of the Messapia x dicoccoides linkage map of Triticum turgidum (L.) Thell. Cellular and Molecular Biology Letters, 2004, 9, 529-41. | 7.0 | 18 |
| 128 | Searching for quantitative trait loci controlling root traits in maize: a critical appraisal. Plant and Soil, 2003, 255, 35-54. | 3.7 | 104 |
| 129 | Microsatellite analysis reveals a progressive widening of the genetic basis in the elite durum wheat germplasm. Theoretical and Applied Genetics, 2003, 107, 783-797. | 3.6 | 104 |
| 130 | Searching for quantitative trait loci controlling root traits in maize: a critical appraisal. , 2003, , 35-54. | | 14 |
| 131 | Unravelling the Genetic Basis of Drought Tolerance in Crops. , 2003, , 71-122. | | 11 |
| 132 | Mapping QTLs Regulating Morpho-physiological Traits and Yield: Case Studies, Shortcomings and Perspectives in Drought-stressed Maize. Annals of Botany, 2002, 89, 941-963. | 2.9 | 331 |
| 133 | Genetic diversity in cultivars and landraces of Oryza sativa subsp. indica as revealed by AFLP markers. Genome, 2002, 45, 451-459. | 2.0 | 34 |
| 134 | Cereal genomics: ushering in a brave new world. Plant Molecular Biology, 2002, 48, 445-449. | 3.9 | 37 |
| 135 | Toward positional cloning of Vgt1, a QTL controlling the transition from the vegetative to the reproductive phase in maize. Plant Molecular Biology, 2002, 48, 601-613. | 3.9 | 116 |
| 136 | Monitoring large-scale changes in transcript abundance in drought- and salt-stressed barley. Plant Molecular Biology, 2002, 48, 551-573. | 3.9 | 503 |
| 137 | Comparative genomic mapping between a 754 kb region flanking DREB1A in Arabidopsis thaliana and maize. Plant Molecular Biology, 2002, 48, 741-750. | 3.9 | 19 |
| 138 | Identification of QTLs for root characteristics in maize grown in hydroponics and analysis of their overlap with QTLs for grain yield in the field at two water regimes. Plant Molecular Biology, 2002, 48, 697-712. | 3.9 | 285 |
| 139 | Direct and Correlated Responses to Divergent Selection for Leaf Abscisic Acid Concentration in Two Maize Populations. Crop Science, 2001, 41, 335-344. | 1.8 | 32 |
| 140 | Title is missing!. Molecular Breeding, 2001, 8, 169-176. | 2.1 | 24 |
| 141 | Genetic Diversity of Japanese Barley Cultivars Based on SSR Analysis Breeding Science, 2001, 51, 215-218. | 1.9 | 7 |
| 142 | Comparative Analyses of the Genetic Diversity among European Spelt and Wheat Accessions with AFLP, RFLP and SSR Markers. Developments in Plant Breeding, 2001, , 733-738. | 0.2 | 1 |
| 143 | Integration of AFLP markers into an RFLP-based map of durum wheat. Plant Breeding, 2000, 119, 393-401. | 1.9 | 31 |
| 144 | The development and application of molecular markers for abiotic stress tolerance in barley. Journal of Experimental Botany, 2000, 51, 19-27. | 4.8 | 117 |

| # | Article | IF | CITATIONS |
|-----|---|------------|----------------|
| 145 | The development and application of molecular markers for abiotic stress tolerance in barley. Journal of Experimental Botany, 2000, 51, 19-27. | 4.8 | 29 |
| 146 | QTL analysis of drought-related traits and grain yield in relation to genetic variation for leaf abscisic acid concentration in field-grown maize. Journal of Experimental Botany, 1999, 50, 1289-1297. | 4.8 | 86 |
| 147 | RFLP mapping of quantitative trait loci controlling abscisic acid concentration in leaves of drought-stressed maize (Zea mays L.). Theoretical and Applied Genetics, 1998, 97, 744-755. | 3.6 | 105 |
| 148 | Comparative analysis of genetic relationships in barley based on RFLP and RAPD markers. Genome, 1997, 40, 607-616. | 2.0 | 29 |
| 149 | Genetic and environmental effects on abscisic acid accumulation in leaves of field-grown maize. Euphytica, 1994, 78, 81-89. | 1.2 | 11 |
| 150 | Recurrent Selection for Regeneration Capacity from Immature Embryoâ€Derived Calli in Maize. Crop Science, 1994, 34, 343-347. | 1.8 | 13 |
| 151 | Abscisic Acid Concentration in Leaf and Xylem Sap, Leaf Water Potential, and Stomatal Conductance in Maize. Crop Science, 1994, 34, 1557-1563. | 1.8 | 20 |
| 152 | Involvement of Chromosomes 5A and 5D in Cold-Induced Abscisic Acid Accumulation in and Frost Tolerance of Wheat Calli. Plant Breeding, 1993, 110, 237-242. | 1.9 | 35 |
| 153 | Effect of Abscisic Acid on Pollen Germination and Tube Growth of Maize Genotypes1. Plant Breeding, 1993, 110, 250-254. | 1.9 | 22 |
| 154 | Number of endosperm cells and endosperm abscisic acid content in relation to kernel weight in four barley genotypes. European Journal of Agronomy, 1992, 1, 125-132. | 4.1 | 14 |
| 155 | Chlorsulfuron Tolerance and Acetolactate Synthase Activity in Corn (<i>Zea mays</i> L.) Inbred Lines. Weed Science, 1991, 39, 553-557. | 1.5 | 20 |
| 156 | Comparison between responses to gametophytic and sporophytic recurrent selection in maize (Zea) Tj ETQq0 0 (| Ͻ rgβT /Ον | erlock 10 Tf 5 |
| 157 | Abscisic acid in developing grains of wheat and barley genotypes differing in grain weight. Plant Growth Regulation, 1988, 7, 3-17. | 3.4 | 26 |
| 158 | Direct and Correlated Responses to Four Cycles of Divergent Selection for Heading Date in Barley*. Plant Breeding, 1988, 101, 313-320. | 1.9 | 0 |
| 159 | Divergent Selection for Heading Date in Barley. Plant Breeding, 1986, 97, 345-351. | 1.9 | 5 |
| 160 | Registration of Gaspé Flint 1.1.1, a smallâ€size earlyâ€flowering maize inbred line. Journal of Plant Registrations, 0, , . | 0.5 | 1 |
| 161 | From QTLS to Genes Controlling Root Traits in Maize. , 0, , 15-24. | | 16 |
| 162 | Molecular Bases of Plant Adaptation to Abiotic Stress and Approaches to Enhance Tolerance to Hostile Environments. , 0 , , . | | 1 |