

# Xing Li

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/1986055/xing-li-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

40  
papers

818  
citations

15  
h-index

28  
g-index

46  
ext. papers

1,077  
ext. citations

7  
avg, IF

3.93  
L-index

#	Paper	IF	Citations
40	Angiotensin II enhances group 2 innate lymphoid cell responses via AT1a during airway inflammation.. <i>Journal of Experimental Medicine</i> , <b>2022</b> , 219,	16.6	1
39	Frequency of asthma exacerbation in children during the coronavirus disease pandemic with strict mitigative countermeasures. <i>Pediatric Pulmonology</i> , <b>2021</b> , 56, 1455-1463	3.5	3
38	Neutrophil-to-Apolipoprotein A1 Ratio Predicted Overall Survival in Hepatocellular Carcinoma Receiving Transarterial Chemoembolization. <i>Oncologist</i> , <b>2021</b> , 26, e1434-e1444	5.7	1
37	Prognostic value of serum HIF-1 $\alpha$ change following transarterial chemoembolization in hepatocellular carcinoma. <i>Clinical and Experimental Medicine</i> , <b>2021</b> , 21, 109-120	4.9	4
36	Interferon Regulatory Factor 4 Regulates the Development of Polymorphonuclear Myeloid-Derived Suppressor Cells Through the Transcription of c-Myc in Cancer. <i>Frontiers in Immunology</i> , <b>2021</b> , 12, 627072	8.4	2
35	Dexamethasone and lactoferrin induced PMN-MDSCs relieved inflammatory adverse events of anti-cancer therapy without tumor promotion. <i>Communications Biology</i> , <b>2021</b> , 4, 252	6.7	
34	Expansion of Group 2 Innate Lymphoid Cells in Patients with End-Stage Renal Disease and Their Clinical Significance. <i>Journal of Immunology</i> , <b>2020</b> , 205, 36-44	5.3	3
33	Accumulation of LOX-1 PMN-MDSCs in nasopharyngeal carcinoma survivors with chronic hepatitis B might permit immune tolerance to epstein-barr virus and relate to tumor recurrence. <i>Aging</i> , <b>2020</b> , 13, 437-449	5.6	0
32	Comparison of Face-Touching Behaviors Before and During the Coronavirus Disease 2019 Pandemic. <i>JAMA Network Open</i> , <b>2020</b> , 3, e2016924	10.4	39
31	MicroRNA-126 Inhibit Viability of Colorectal Cancer Cell by Repressing mTOR Induced Apoptosis and Autophagy. <i>OncoTargets and Therapy</i> , <b>2020</b> , 13, 2459-2468	4.4	16
30	Expansion of Monocytic Myeloid-Derived Suppressor Cells in Patients Under Hemodialysis Might Lead to Cardiovascular and Cerebrovascular Events. <i>Frontiers in Immunology</i> , <b>2020</b> , 11, 577253	8.4	1
29	Neutrophil-To-Lymphocyte Ratio Predicted Long-Term Chemotherapy Benefits In Stage IIIB-IV Non-Squamous Non-Small Cell Lung Cancer Patients Without Sensitive Mutations. <i>OncoTargets and Therapy</i> , <b>2019</b> , 12, 8779-8787	4.4	4
28	Lactoferrin-induced myeloid-derived suppressor cell therapy attenuates pathologic inflammatory conditions in newborn mice. <i>Journal of Clinical Investigation</i> , <b>2019</b> , 129, 4261-4275	15.9	29
27	IRF-7 Is a Critical Regulator of Type 2 Innate Lymphoid Cells in Allergic Airway Inflammation. <i>Cell Reports</i> , <b>2019</b> , 29, 2718-2730.e6	10.6	15
26	Transitory presence of myeloid-derived suppressor cells in neonates is critical for control of inflammation. <i>Nature Medicine</i> , <b>2018</b> , 24, 224-231	50.5	98
25	ICAM-1 controls development and function of ILC2. <i>Journal of Experimental Medicine</i> , <b>2018</b> , 215, 2157-2174	17.6	34
24	Brazilin induces FOXO3A-dependent autophagic cell death by disturbing calcium homeostasis in osteosarcoma cells. <i>Cancer Chemotherapy and Pharmacology</i> , <b>2018</b> , 82, 479-491	3.5	10

23	Identification of abnormally expressed lncRNAs induced by PM2.5 in human bronchial epithelial cells. <i>Bioscience Reports</i> , <b>2018</b> , 38,	4.1	25
22	Comparison of five staging systems in predicting the survival rate of patients with hepatocellular carcinoma undergoing trans-arterial chemoembolization therapy. <i>Oncology Letters</i> , <b>2018</b> , 15, 855-862	2.6	8
21	Endoplasmic reticulum stress induced LOX-1 CD15 polymorphonuclear myeloid-derived suppressor cells in hepatocellular carcinoma. <i>Immunology</i> , <b>2018</b> , 154, 144-155	7.8	56
20	Combination of ULK1 and LC3B improve prognosis assessment of hepatocellular carcinoma. <i>Biomedicine and Pharmacotherapy</i> , <b>2018</b> , 97, 195-202	7.5	22
19	Timing of regular egg intake for prevention of egg allergy. <i>Journal of Allergy and Clinical Immunology</i> , <b>2018</b> , 141, 459-460	11.5	2
18	Circular RNA circCDK13 suppresses cell proliferation, migration and invasion by modulating the JAK/STAT and PI3K/AKT pathways in liver cancer. <i>International Journal of Oncology</i> , <b>2018</b> , 53, 246-256	4.4	18
17	Selection of glucocorticoid-sensitive patients in interstitial lung disease secondary to connective tissue diseases population by radiomics. <i>Therapeutics and Clinical Risk Management</i> , <b>2018</b> , 14, 1975-1986 <sup>2.9</sup>		7
16	Identification of ceRNA network based on a RNA-seq shows prognostic lncRNA biomarkers in human lung adenocarcinoma. <i>Oncology Letters</i> , <b>2018</b> , 16, 5697-5708	2.6	25
15	Expansion of polymorphonuclear myeloid-derived suppressor cells in patients with end-stage renal disease may lead to infectious complications. <i>Kidney International</i> , <b>2017</b> , 91, 1236-1242	9.9	13
14	Expansion of monocytic myeloid-derived suppressor cells in endometriosis patients: A pilot study. <i>International Immunopharmacology</i> , <b>2017</b> , 47, 150-158	5.8	12
13	A 2-year step-down withdrawal from inhaled corticosteroids in asthmatic children receiving immunotherapy. <i>World Journal of Pediatrics</i> , <b>2017</b> , 13, 560-565	4.6	1
12	Different patient satisfaction levels between the first and second knee in the early stage after simultaneous bilateral total knee arthroplasty (TKA): a comparison between subjective and objective outcome assessments. <i>Journal of Orthopaedic Surgery and Research</i> , <b>2017</b> , 12, 121	2.8	5
11	Neutrophil count is associated with myeloid derived suppressor cell level and presents prognostic value of for hepatocellular carcinoma patients. <i>Oncotarget</i> , <b>2017</b> , 8, 24380-24388	3.3	29
10	Coenzyme Q10 Prevents the Interleukin-1 Beta Induced Inflammatory Response via Inhibition of MAPK Signaling Pathways in Rat Articular Chondrocytes. <i>Drug Development Research</i> , <b>2017</b> , 78, 403-410 <sup>5.1</sup>		13
9	Serum Golgi protein 73 is a prognostic rather than diagnostic marker in hepatocellular carcinoma. <i>Oncology Letters</i> , <b>2017</b> , 14, 6277-6284	2.6	6
8	miR-30b regulates chondrogenic differentiation of mouse embryo-derived stem cells by targeting SOX9. <i>Experimental and Therapeutic Medicine</i> , <b>2017</b> , 14, 6131-6137	2.1	5
7	Atorvastatin promotes the expansion of myeloid-derived suppressor cells and attenuates murine colitis. <i>Immunology</i> , <b>2016</b> , 149, 432-446	7.8	24
6	Associations of Short-Term and Long-Term Exposure to Ambient Air Pollutants With Hypertension: A Systematic Review and Meta-Analysis. <i>Hypertension</i> , <b>2016</b> , 68, 62-70	8.5	167

5	Alanine aminotransferase to hemoglobin ratio is an indicator for disease progression for hepatocellular carcinoma patients receiving transcatheter arterial chemoembolization. <i>Tumor Biology</i> , <b>2016</b> , 37, 2951-9	2.9	4
4	Hypoxemia is an independent predictor of bronchiolitis obliterans following respiratory adenoviral infection in children. <i>SpringerPlus</i> , <b>2016</b> , 5, 1622		9
3	Platelet-to-lymphocyte ratio acts as a prognostic factor for patients with advanced hepatocellular carcinoma. <i>Tumor Biology</i> , <b>2015</b> , 36, 2263-9	2.9	61
2	Aberrant expression of enhancer of zeste homologue 2, correlated with HIF-1 $\alpha$ refines relapse risk and predicts poor outcome for breast cancer. <i>Oncology Reports</i> , <b>2014</b> , 32, 1101-7	3.5	12
1	Neutrophil-to-lymphocyte ratio acts as a prognostic factor for patients with advanced hepatocellular carcinoma. <i>Tumor Biology</i> , <b>2014</b> , 35, 11057-63	2.9	34