

Carmen Lucia Queiroga

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1996998/publications.pdf>

Version: 2024-02-01

28
papers

808
citations

430874

18
h-index

501196

28
g-index

28
all docs

28
docs citations

28
times ranked

1334
citing authors

#	ARTICLE	IF	CITATIONS
1	Evaluation of the antiulcerogenic activity of friedelan-3 β -ol and friedelin isolated from <i>Maytenus ilicifolia</i> (Celastraceae). <i>Journal of Ethnopharmacology</i> , 2000, 72, 465-468.	4.1	88
2	Extracts from pitanga (<i>Eugenia uniflora</i> L.) leaves: Influence of extraction process on antioxidant properties and yield of phenolic compounds. <i>Journal of Supercritical Fluids</i> , 2011, 55, 998-1006.	3.2	85
3	Extraction of phenolic compounds from pitanga (<i>Eugenia uniflora</i> L.) leaves by sequential extraction in fixed bed extractor using supercritical CO ₂ , ethanol and water as solvents. <i>Journal of Supercritical Fluids</i> , 2014, 86, 4-14.	3.2	72
4	Extraction of phenolic compounds from pepper-rosmarin (<i>Lippia sidoides</i> Cham.) leaves by sequential extraction in fixed bed extractor using supercritical CO ₂ , ethanol and water as solvents. <i>Journal of Supercritical Fluids</i> , 2015, 99, 68-75.	3.2	59
5	Kinetics, composition and biological activity of <i>Eupatorium intermedium</i> flower extracts obtained from scCO ₂ and compressed propane. <i>Journal of Supercritical Fluids</i> , 2015, 97, 145-153.	3.2	44
6	Composition and antimalarial activity of extracts of <i>Curcuma longa</i> L. obtained by a combination of extraction processes using supercritical CO ₂ , ethanol and water as solvents. <i>Journal of Supercritical Fluids</i> , 2017, 119, 122-129.	3.2	44
7	Fractionated extraction of saponins from Brazilian ginseng by sequential process using supercritical CO ₂ , ethanol and water. <i>Journal of Supercritical Fluids</i> , 2014, 92, 272-281.	3.2	37
8	Extracts from the leaves of <i>Baccharis dracunculifolia</i> obtained by a combination of extraction processes with supercritical CO ₂ , ethanol and water. <i>Journal of Supercritical Fluids</i> , 2012, 63, 31-39.	3.2	35
9	Chemical constituents of the volatile oil from leaves of <i>Annona coriacea</i> and in vitro antiprotozoal activity. <i>Revista Brasileira De Farmacognosia</i> , 2011, 21, 0-0.	1.4	33
10	Composition of the Essential Oil of Vassoura. <i>Journal of the Brazilian Chemical Society</i> , 1990, 1, 105-109.	0.6	33
11	Three new oxygenated cadinanes from <i>Baccharis</i> species. <i>Phytochemistry</i> , 1996, 42, 1097-1103.	2.9	29
12	Linalool production from the leaves of <i>Bursera aloexylon</i> and its antimicrobial activity. <i>Farmacognosia</i> , 2007, 78, 327-328.	2.2	29
13	Comparison of volatile and polyphenolic compounds in Brazilian green propolis and its botanical origin <i>Baccharis dracunculifolia</i> . <i>Food Science and Technology</i> , 2008, 28, 178-181.	1.7	25
14	Wood typification by Venturi easy ambient sonic spray ionization mass spectrometry: the case of the endangered Mahogany tree. <i>Journal of Mass Spectrometry</i> , 2012, 47, 1-6.	1.6	25
15	Evaluation of paraffins biodegradation and biosurfactant production by <i>Bacillus subtilis</i> in the presence of crude oil. <i>Brazilian Journal of Microbiology</i> , 2003, 34, 321-324.	2.0	24
16	Extraction of <i>Campomanesia xanthocarpa</i> fruit using supercritical CO ₂ and bioactivity assessments. <i>Journal of Supercritical Fluids</i> , 2015, 98, 79-85.	3.2	24
17	Seasonal Variation of the (E)-Nerolidol and Other Volatile Compounds Within Ten Different Cultivated Populations of <i>Baccharis dracunculifolia</i> D.C. (Asteraceae). <i>Journal of Essential Oil Research</i> , 2009, 21, 308-314.	2.7	22
18	Anticholinesterase activity evaluation of alkaloids and coumarin from stems of <i>Conchocarpus fontanesianus</i> . <i>Revista Brasileira De Farmacognosia</i> , 2012, 22, 374-380.	1.4	22

#	ARTICLE	IF	CITATIONS
19	Sequential extraction of bioactive compounds from <i>Melia azedarach</i> L. in fixed bed extractor using CO ₂ , ethanol and water. <i>Journal of Supercritical Fluids</i> , 2014, 95, 355-363.	3.2	22
20	<i>In vitro</i> antiviral activity of Brazilian plants (<i>Maytenus ilicifolia</i> and <i>Aniba</i>) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 707 Td (r</i> <i>Biology</i> , 2012, 50, 1269-1275.	2.9	10
21	Production of Copaiba oleoresin particles from emulsions stabilized with modified starches. <i>Industrial Crops and Products</i> , 2017, 108, 128-139.	5.2	9
22	Comparison of the Chemical Composition of the Essential Oil and the Water Soluble Oil of <i>Baccharis dracunculifolia</i> DC. (Asteraceae). <i>Journal of Essential Oil Research</i> , 2008, 20, 111-114.	2.7	8
23	Production of copaiba (<i>Copaifera officinalis</i>) oleoresin particles by supercritical fluid extraction of emulsions. <i>Journal of Supercritical Fluids</i> , 2018, 140, 364-371.	3.2	7
24	Chemical composition and biological activity of <i>Eupatorium intermedium</i> essential oil. <i>Journal of Essential Oil Research</i> , 2017, 29, 93-100.	2.7	6
25	Solubility of oleic acid, triacylglycerol and their mixtures in supercritical carbon dioxide and thermodynamic modeling of phase equilibrium. <i>Journal of Supercritical Fluids</i> , 2019, 143, 275-285.	3.2	6
26	Study of the Variation of the Composition of the Essential Oil of Leaves and Flowers of <i>Achyrocline alata</i> (D.C.) Along a Period of the Day. <i>Journal of Essential Oil Research</i> , 2002, 14, 280-281.	2.7	5
27	High-speed countercurrent chromatography as a tool to isolate nerolidol from the <i>Baccharis dracunculifolia</i> volatile oil. <i>Journal of Essential Oil Research</i> , 2014, 26, 334-337.	2.7	4
28	Fractionation of sesquiterpenes and diterpenic acids from copaiba (<i>Copaifera officinalis</i>) oleoresin using supercritical adsorption. <i>Journal of Supercritical Fluids</i> , 2022, 184, 105565.	3.2	1