

# Marco Vecoli

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2013987/publications.pdf>

Version: 2024-02-01

64  
papers

1,610  
citations

257357

24  
h-index

330025

37  
g-index

69  
all docs

69  
docs citations

69  
times ranked

1076  
citing authors

#	ARTICLE	IF	CITATIONS
1	Low-latitude ?upper Barremianâ€“lower Aptian palynoflora and paleovegetation of the Biyadh Formation (Arabian Plate, eastern margin of northern Gondwana): evidence for a possible cold snap. <i>Cretaceous Research</i> , 2022, 129, 104995.	0.6	7
2	Angiosperm pollen assemblages from the Lower Cretaceous (Barremianâ€“lower Aptian) of offshore Saudi Arabia and their implications for early patterns of angiosperm radiation. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2022, 599, 111052.	1.0	2
3	Palynofacies, paleoenvironment and thermal maturity of early Silurian shales in Saudi Arabia (Qusaiba) Tj ETQq1 1 0,784314 rgBT /Ov	0.8	1
4	Palynological Correlation of the Late Cambrian to Middle Ordovician Saq Formation in Saudi Arabia and Equivalent Strata in Oman Manuscript Title. , 2019, , .		2
5	Application of Palynomorph Darkness Index (PDI) to assess the thermal maturity of palynomorphs: A case study from North Africa. <i>International Journal of Coal Geology</i> , 2018, 188, 64-78.	1.9	26
6	Organic matter deposition in the Ghadames Basin (Libya) during the Late Devonianâ€“A multidisciplinary approach. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018, 497, 37-51.	1.0	21
7	First Appearance Datums (FADs) of selected acritarch taxa and correlation between Lower and Middle Ordovician. <i>Lethaia</i> , 2018, 51, 228-253.	0.6	26
8	Wall ultrastructure of the oldest embryophytic spores: Implications for early land plant evolution. <i>Revue De Micropaleontologie</i> , 2017, 60, 281-288.	0.8	17
9	Correlation of Palynomorph Darkness Index and vitrinite reflectance in a submature Carboniferous well section in northern Saudi Arabia. <i>Revue De Micropaleontologie</i> , 2017, 60, 411-416.	0.8	9
10	Middle Ordovician acritarchs and problematic organic-walled microfossils from the Saq-Hanadir transitional beds in the QSIM-801 well, Saudi Arabia. <i>Revue De Micropaleontologie</i> , 2017, 60, 289-318.	0.8	14
11	A punctuated Late Ordovician and early Silurian deglaciation and transgression: Evidence from the subsurface of northern Saudi Arabia. <i>AAPG Bulletin</i> , 2017, 101, 863-886.	0.7	25
12	Middle Ordovician cryptospores from the Saq-Hanadir transitional beds in the QSIM-801 well, Saudi Arabia. <i>Revue De Micropaleontologie</i> , 2017, 60, 319-331.	0.8	8
13	New palynomorphs from the Ordovicianâ€“Silurian boundary interval: Eastern North America and Saudi Arabia. <i>Palynology</i> , 2017, 41, 106-120.	0.7	3
14	Late Ordovician (Katian) chitinozoans from northwest Saudi Arabia: Biostratigraphic and paleoenvironmental implications. <i>Revue De Micropaleontologie</i> , 2017, 60, 333-369.	0.8	9
15	Ordovician spore â€“thalliâ€™ and the evolution of the plant sporophyte. <i>Palynology</i> , 2017, 41, 57-68.	0.7	16
16	The terrestrialization process: A palaeobotanical and palynological perspective (2). <i>Review of Palaeobotany and Palynology</i> , 2016, 227, 1-3.	0.8	4
17	The terrestrialization process: A palaeobotanical and palynological perspective. <i>Review of Palaeobotany and Palynology</i> , 2016, 224, 1-3.	0.8	6
18	Plant evolution and terrestrialization during Palaeozoic timesâ€“The phylogenetic context. <i>Review of Palaeobotany and Palynology</i> , 2016, 227, 4-18.	0.8	60

#	ARTICLE	IF	CITATIONS
19	Palynology of the Ordovician Kanosh Shale at Fossil Mountain, Utah. <i>Journal of Paleontology</i> , 2015, 89, 424-447.	0.5	14
20	The Ordovician acritarch <i>Dactylofusa velifera</i> Cocchio 1982: a biostratigraphical and palaeogeographical index species. <i>Palynology</i> , 2015, 39, 125-141.	0.7	14
21	Cryptospores from the Hanadir Shale Member of the Qasim Formation, Ordovician (Darriwilian) of Saudi Arabia: taxonomy and systematics. <i>Review of Palaeobotany and Palynology</i> , 2015, 212, 97-110.	0.8	42
22	The 2.1 Ga Old Francevillian Biota: Biogenicity, Taphonomy and Biodiversity. <i>PLoS ONE</i> , 2014, 9, e99438.	1.1	53
23	The palaeobiogeographical spread of the acritarch <i>Veryhachium</i> in the Early and Middle Ordovician and its impact on biostratigraphical applications. <i>Gff</i> , 2014, 136, 234-237.	0.4	5
24	Late Tremadocian-early Floian acritarchs from graptolitic shales of the Yinzhubu and Ningkuo formations of Yiyang, South China. <i>Review of Palaeobotany and Palynology</i> , 2013, 193, 1-14.	0.8	14
25	Chapter 29 Palaeophytogeography of Ordovician-Silurian land plants. <i>Geological Society Memoir</i> , 2013, 38, 461-476.	0.9	44
26	Palaeozoic applied marine palynology. <i>Review of Palaeobotany and Palynology</i> , 2013, 198, 1.	0.8	0
27	Chitinozoan biostratigraphy across the Katian (Late Ordovician) GICE event in the Borensult-1 drillcore (Sweden). <i>Review of Palaeobotany and Palynology</i> , 2013, 198, 134-144.	0.8	3
28	Systematic occurrences of malformed (teratological) acritarchs in the run-up of Early Palaeozoic $\delta^{13}C$ isotope excursions. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2012, 367-368, 137-146.	1.0	25
29	Abnormal forms of acritarchs (phytoplankton) in the upper Hirnantian (Upper Ordovician) of Anticosti Island, Canada. <i>Review of Palaeobotany and Palynology</i> , 2012, 173, 46-56.	0.8	17
30	Biostratigraphy and paleoenvironmental characterization of the Middle Ordovician from the Sierras Subandinas (NW Argentina) based on organic-walled microfossils and sequence stratigraphy. <i>Journal of South American Earth Sciences</i> , 2011, 31, 124-138.	0.6	24
31	Aliphatic and aromatic biomarkers from Carboniferous coal deposits at Dunbar (East Lothian, UK). <i>Palaeoclimatology, Palaeoecology</i> , 2011, 309, 309-326.	1.0	46
32	Phytoplankton dynamics across the Ordovician/Silurian boundary at low palaeolatitudes: Correlations with carbon isotopic and glacial events. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2011, 312, 79-97.	1.0	47
33	New sedimentological and biostratigraphic data in the Kwahu Group (Meso- to Neo-Proterozoic), southern margin of the Volta Basin, Ghana: Stratigraphic constraints and implications on regional lithostratigraphic correlations. <i>Precambrian Research</i> , 2011, 189, 155-175.	1.2	22
34	Aliphatic and aromatic biomarkers from Gondwanan sediments of Late Ordovician to Early Devonian age: An early terrestrialization approach. <i>Organic Geochemistry</i> , 2011, 42, 605-617.	0.9	40
35	Palynology and isotope geochronology of the Upper Ordovician-Silurian successions (Ghelli and) stratigraphic and palaeogeographic implications. <i>Review of Palaeobotany and Palynology</i> , 2011, 164, 251-271.	0.8	50
36	Cryptospore assemblages from Upper Ordovician (Katian-Hirnantian) strata of Anticosti Island, Quebec, Canada, and Estonia: Palaeophytogeographic and palaeoclimatic implications. <i>Review of Palaeobotany and Palynology</i> , 2011, 166, 76-93.	0.8	35

#	ARTICLE	IF	CITATIONS
37	Acritarchs from the Ordovician–Silurian boundary beds of the Valga-10 drill core, southern Estonia (Baltica) and their stratigraphical and palaeobiogeographical implications. <i>Palynology</i> , 2011, 35, 4-45.	0.7	21
38	Laser Raman micro-spectroscopy of Proterozoic and Palaeozoic organic-walled microfossils (acritarchs and prasinophytes) from the Ghadamis Basin, Libya and Volta Basin, Ghana. <i>Spectroscopy</i> , 2010, 24, 207-212.	0.8	5
39	Occurrence of retene in upper Silurian–lower Devonian sediments from North Africa: Origin and implications. <i>Organic Geochemistry</i> , 2010, 41, 302-306.	0.9	40
40	The effects of terrestrialization on marine ecosystems: the fall of CO <sub>2</sub> . <i>Geological Society Special Publication</i> , 2010, 339, 37-48.	0.8	9
41	Morphology and wall ultrastructure of the megaspore <i>Lagenicula (Triletes) mixta</i> (Winslow 1962) comb. nov. from the Carboniferous (Early Mississippian: mid Tournaisian) of Ohio, USA. <i>Review of Palaeobotany and Palynology</i> , 2009, 156, 51-61.	0.8	5
42	Palynology, organic geochemistry and carbon isotope analysis of a latest Ordovician through Silurian clastic succession from borehole Tt1, Ghadamis Basin, southern Tunisia, North Africa: Palaeoenvironmental interpretation. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2009, 273, 378-394.	1.0	74
43	Palynostratigraphy and vegetational changes in the Siluro-Devonian of the Ghadamis Basin, North Africa. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2009, 282, 1-18.	1.0	31
44	Cambrian Acritarchs from the Col di Foglia (Agordo) southalpine metamorphic basement, Italian Eastern Alps: the oldest biostratigraphic record in the alps. <i>Rendiconti Lincei</i> , 2008, 19, 45-55.	1.0	6
45	The Ordovician Biodiversification: revolution in the oceanic trophic chain. <i>Lethaia</i> , 2008, 41, 99-109.	0.6	175
46	The Ordovician acritarch genus <i>Coryphidium</i> . <i>Revue De Micropaleontologie</i> , 2008, 51, 97-120.	0.8	19
47	Fossil microphytoplankton dynamics across the Ordovician–Silurian boundary. <i>Review of Palaeobotany and Palynology</i> , 2008, 148, 91-107.	0.8	24
48	First biostratigraphic (palynological) dating of Middle and Late Cambrian strata in the subsurface of northwestern Algeria, North Africa: Implications for regional stratigraphy. <i>Review of Palaeobotany and Palynology</i> , 2008, 149, 57-62.	0.8	11
49	Palynostratigraphy of Middle Cambrian to lowermost Ordovician stratal sequences in the High Zagros Mountains, southern Iran: Regional stratigraphic implications, and palaeobiogeographic significance. <i>Review of Palaeobotany and Palynology</i> , 2008, 150, 97-114.	0.8	27
50	The acritarch genus <i>Veryhachium</i> Deunff 1954: Taxonomic evaluation and first appearance. <i>Palynology</i> , 2007, 31, 191-203.	0.7	46
51	A review of the Late Cambrian (Furongian) palaeogeography in the western Mediterranean region, NW Gondwana. <i>Earth-Science Reviews</i> , 2007, 85, 47-81.	4.0	54
52	Ordovician chitinozoans and acritarchs from southern and southeastern Turkey. <i>Revue De Micropaleontologie</i> , 2007, 50, 81-107.	0.8	57
53	Environmental changes reflected by palynomorphs in the early Middle Ordovician Hanadir Member of the Qasim Formation, Saudi Arabia. <i>Revue De Micropaleontologie</i> , 2007, 50, 3-16.	0.8	40
54	Latest Ordovician–early Silurian chitinozoans from the eastern Alborz Mountain Range, Kopet–Dagh region, northeastern Iran: biostratigraphy and palaeobiogeography. <i>Review of Palaeobotany and Palynology</i> , 2007, 145, 173-192.	0.8	28

#	ARTICLE	IF	CITATIONS
55	32. Acritarchs. , 2004, , 348-360.		17
56	Billigen (Lower Arenig/Lower Ordovician) acritarchs from the East European Platform and their palaeobiogeographic significance. <i>Lethaia</i> , 2004, 37, 97-111.	0.6	13
57	<i>Rhopaliophora? asymmetrica</i> , a new acritarch species from the lowermost <i>Arenig</i> ™ of Baltica and its biostratigraphic potential. <i>Review of Palaeobotany and Palynology</i> , 2003, 126, 39-48.	0.8	4
58	Timing of the Avalonia-Baltica plate convergence as inferred from palaeogeographic and stratigraphic data of chitinozoan assemblages in west Pomerania, northern Poland. <i>Geological Society Special Publication</i> , 2002, 201, 95-113.	0.8	15
59	Quantitative evaluation of microplankton palaeobiogeography in the Ordovician–Early Silurian of the northern Trans European Suture Zone: implications for the timing of the Avalonia–Baltica collision. <i>Review of Palaeobotany and Palynology</i> , 2001, 115, 43-68.	0.8	33
60	Ordovician palynomorphs from the subsurface of R <sup>1</sup> / <sub>4</sub> gen (NE-Germany): review and perspectives. <i>Neues Jahrbuch Fur Geologie Und Palaontologie - Abhandlungen</i> , 2001, 222, 123-139.	0.2	11
61	Ordovician-Silurian palynostratigraphy (Chitinozoa and acritarchs) of the G14-1/86 borehole, southern Baltic Sea. <i>Neues Jahrbuch Fur Geologie Und Palaontologie - Abhandlungen</i> , 2001, 222, 91-122.	0.2	12
62	Palaeoenvironmental interpretation of microphytoplankton diversity trends in the Cambrian–Ordovician of the northern Sahara Platform. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2000, 160, 329-346.	1.0	55
63	Stratigraphically significant acritarchs in uppermost Cambrian to basal Ordovician strata of Northwestern Algeria. <i>Grana</i> , 1997, 36, 17-28.	0.4	10
64	Ordovician. , 0, , 203-248.		9