## Kenichi Meguro

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2017353/publications.pdf

Version: 2024-02-01

135 papers 6,593 citations

236612 25 h-index 79 g-index

147 all docs

147 docs citations

147 times ranked 8203 citing authors

#	Article	IF	CITATIONS
1	Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS–ADRDA criteria. Lancet Neurology, The, 2007, 6, 734-746.	4.9	3,755
2	Motoric cognitive risk syndrome. Neurology, 2014, 83, 718-726.	1.5	345
3	Prevalence of Possible Idiopathic Normal-Pressure Hydrocephalus in Japan: The Osaki-Tajiri Project. Neuroepidemiology, 2009, 32, 171-175.	1.1	149
4	Cerebral circulation and oxygen metabolism associated with subclinical periventricular hyperintensity as shown by magnetic resonance imaging. Annals of Neurology, 1990, 28, 378-383.	2.8	142
5	Prevalence of Dementia and Dementing Diseases in Japan. Archives of Neurology, 2002, 59, 1109.	4.9	126
6	Effect of age and education on the Trail Making Test and determination of normative data for Japanese elderly people: The Tajiri Project. Psychiatry and Clinical Neurosciences, 2006, 60, 422-428.	1.0	89
7	Determinants of cognitive performance and decline in 20 diverse ethno-regional groups: A COSMIC collaboration cohort study. PLoS Medicine, 2019, 16, e1002853.	3.9	86
8	Prevalence of dementia and dementing diseases in the oldâ€old population in Japan: the Kurihara Project. Implications for Longâ€Term Care Insurance data. Psychogeriatrics, 2012, 12, 226-234.	0.6	82
9	Trends in the Prevalence of Dementia in Japan. International Journal of Alzheimer's Disease, 2012, 2012, 1-11.	1.1	73
10	Prevalence and Cognitive Performances of Clinical Dementia Rating 0.5 and Mild Cognitive Impairment in Japan. Alzheimer Disease and Associated Disorders, 2004, 18, 3-10.	0.6	70
11	Decreased striatal D2 receptor density associated with severe behavioral abnormality in Alzheimer's disease. Annals of Nuclear Medicine, 2003, 17, 567-573.	1.2	68
12	Incidence of dementia and associated risk factors in Japan: The Osaki-Tajiri Project. Journal of the Neurological Sciences, 2007, 260, 175-182.	0.3	67
13	Cognitive function and frontal lobe atrophy in normal elderly adults: Implications for dementia not as aging-related disorders and the reserve hypothesis. Psychiatry and Clinical Neurosciences, 2001, 55, 565-572.	1.0	60
14	Mild cognitive impairment, poor episodic memory, and late-life depression are associated with cerebral cortical thinning and increased white matter hyperintensities. Frontiers in Aging Neuroscience, 2014, 6, 306.	1.7	59
15	Risperidone is Effective for Wandering and Disturbed Sleep/Wake Patterns in Alzheimer's Disease. Journal of Geriatric Psychiatry and Neurology, 2004, 17, 61-67.	1.2	54
16	Chronic Kidney Disease: A Risk Factor for Dementia Onset: A Populationâ€Based Study. The Osakiâ€Tajiri Project. Journal of the American Geriatrics Society, 2011, 59, 1175-1181.	1.3	53
17	Non-verbal learning is impaired in very mild Alzheimer's disease (CDR 0.5): Normative data from the learning version of the Rey-Osterrieth Complex Figure Test. Psychiatry and Clinical Neurosciences, 2006, 60, 139-146.	1.0	49
18	Combined Memory and Executive Function Tests Can Screen Mild Cognitive Impairment and Converters to Dementia in a Community: The Osaki-Tajiri Project. Neuroepidemiology, 2009, 33, 103-110.	1.1	42

#	Article	IF	Citations
19	Therapeutic Psychosocial Intervention for Elderly Subjects With Very Mild Alzheimer Disease in a Community: The Tajiri Project. Alzheimer Disease and Associated Disorders, 2002, 16, 261-269.	0.6	39
20	Corpus Callosum Atrophy, White Matter Lesions, and Frontal Executive Dysfunction in Normal Aging and Alzheimer's Disease. A Community-Based Study: The Tajiri Project. International Psychogeriatrics, 2003, 15, 9-25.	0.6	35
21	Improved social interaction and increased anterior cingulate metabolism after group reminiscence with reality orientation approach for vascular dementia. Psychiatry Research - Neuroimaging, 2011, 192, 183-187.	0.9	35
22	A Randomized Controlled Trial of the Group Reminiscence Approach in Patients with Vascular Dementia. Dementia and Geriatric Cognitive Disorders, 2007, 24, 48-54.	0.7	34
23	Donepezil and life expectancy in Alzheimer's disease: A retrospective analysis in the Tajiri Project. BMC Neurology, 2014, 14, 83.	0.8	33
24	Comprehensive approach of donepezil and psychosocial interventions on cognitive function and quality of life for Alzheimer's disease: the Osaki-Tajiri Project. Age and Ageing, 2008, 37, 469-473.	0.7	31
25	Disorders of "taste cognition―are associated with insular involvement in patients with Alzheimer's disease and vascular dementia: "Memory of food is impaired in dementia and responsible for poor diet― International Psychogeriatrics, 2014, 26, 1127-1138.	0.6	28
26	Language deterioration in four Japanese-Portuguese bilingual patients with Alzheimer's disease: a trans-cultural study of Japanese elderly immigrants in Brazil. Psychogeriatrics, 2003, 3, 63-68.	0.6	26
27	Necker cube copying ability in normal elderly and Alzheimer's disease. A community-based study: The Tajiri project. Psychogeriatrics, 2006, 6, 4-9.	0.6	26
28	Cortical glucose metabolism in psychiatric wandering patients with vascular dementia. Psychiatry Research - Neuroimaging, 1996, 67, 71-80.	0.9	25
29	Depressive symptoms and associated factors in a cognitively normal elderly population: the Tajiri Project. International Journal of Geriatric Psychiatry, 2001, 16, 780-788.	1.3	25
30	Cross-cultural comparisons of the Mini-mental State Examination between Japanese and U.S. cohorts. International Psychogeriatrics, 2009, 21, 113.	0.6	25
31	Screening for Very Mild Subcortical Vascular Dementia Patients Aged 75 and Above Using the Montreal Cognitive Assessment and Mini-Mental State Examination in a Community: The Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2012, 2, 503-515.	0.6	25
32	Prevalence of senile dementia in a rural community in Japan: the Tajiri project. Archives of Gerontology and Geriatrics, 2000, 29, 249-265.	1.4	23
33	Executive dysfunction can explain word-list learning disability in very mild Alzheimer's disease: The Tajiri Project. Psychiatry and Clinical Neurosciences, 2004, 58, 54-60.	1.0	23
34	Decreased cortical glucose metabolism in converters from CDR 0.5 to Alzheimer's disease in a community: the Osaki-Tajiri Project. International Psychogeriatrics, 2009, 21, 148.	0.6	22
35	A Cluster Randomized Controlled Trial of Nonpharmacological Interventions for Old-Old Subjects with a Clinical Dementia Rating of 0.5: The Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2015, 5, 221-232.	0.6	22
36	Neuroepidemiologic and Neurobehavioral Characteristics of Motoric Cognitive Risk Syndrome in an Old-Old Population: The Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2016, 6, 176-182.	0.6	22

3

#	Article	IF	CITATIONS
37	Silent Infarction or White Matter Hyperintensity and Impaired Attention Task Scores in a Nondemented Population: The Osaki-Tajiri Project. Journal of Stroke and Cerebrovascular Diseases, 2012, 21, 275-282.	0.7	21
38	Does parity matter in women's risk of dementia? A COSMIC collaboration cohort study. BMC Medicine, 2020, 18, 210.	2.3	21
39	Impaired verbal description ability assessed by the Picture Description Task in Alzheimer's disease. Archives of Gerontology and Geriatrics, 1998, 27, 57-65.	1.4	19
40	Neuropsychosocial Features of Very Mild Alzheimer's Disease (CDR 0.5) and Progression to Dementia in a Community: The Tajiri Project. Journal of Geriatric Psychiatry and Neurology, 2004, 17, 183-189.	1.2	18
41	Different MRI findings for normal elderly and very mild Alzheimer's disease in a community: Implications for clinical practice. Archives of Gerontology and Geriatrics, 2006, 42, 59-71.	1.4	18
42	Elderly Japanese emigrants to Brazil before World War II: II. prevalence of senile dementia. International Journal of Geriatric Psychiatry, 2001, 16, 775-779.	1.3	17
43	Content of delusional thoughts in Alzheimer's disease and assessment of content-specific brain dysfunctions with BEHAVE-AD-FW and SPECT. International Psychogeriatrics, 2013, 25, 939-948.	0.6	16
44	Fear of Falling and Cognitive Impairments in Elderly People with Hip Fractures. Dementia and Geriatric Cognitive Disorders Extra, 2018, 7, 386-394.	0.6	16
45	Decreased Behavioral Abnormalities After Treatment with Combined Donepezil and Yokukansankachimpihange in Alzheimer Disease: An Observational Study. The Osaki-Tajiri Project. Neurology and Therapy, 2018, 7, 333-340.	1.4	16
46	Impaired memory and executive function associated with decreased medial temporal and prefrontal blood flow in Clinical Dementia Rating 0.5 status: the Osaki†ajiri project. Psychogeriatrics, 2012, 12, 27-33.	0.6	15
47	Impaired instrumental activities of daily living affect conversion from mild cognitive impairment to dementia: the Osakiâ€√ajiri Project. Psychogeriatrics, 2012, 12, 34-42.	0.6	15
48	Apathy is more severe in vascular than amnestic mild cognitive impairment in a community: The <scp>K</scp> urihara <scp>P</scp> roject. Psychiatry and Clinical Neurosciences, 2013, 67, 517-525.	1.0	15
49	The effects of group work therapy in patients with Alzheimer's disease. International Journal of Geriatric Psychiatry, 2000, 15, 532-534.	1.3	14
50	Vascular dementia with left thalamic infarction: Neuropsychological and behavioral implications suggested by involvement of the thalamic nucleus and the remote effect on cerebral cortex. The Osaki–Tajiri project. Psychiatry Research - Neuroimaging, 2013, 213, 56-62.	0.9	14
51	Clinical utility of the <scp>F</scp> unctional <scp>I</scp> ndependence <scp>M</scp> easure for assessment of patients with <scp>A</scp> lzheimer's disease and vascular dementia. Psychogeriatrics, 2013, 13, 199-205.	0.6	14
52	Speaking one more language in early life has only minor effects on cognition in Taiwanese with low education level: the Taishan Project. Psychogeriatrics, 2017, 17, 256-261.	0.6	14
53	Vascular Mild Cognitive Impairment: Identifying Disease in Community-Dwelling Older Adults, Reducing Risk Factors, and Providing Support. The Osaki-Tajiri and Kurihara Projects. Journal of Alzheimer's Disease, 2019, 70, S293-S302.	1.2	14
54	Costs and Resource Use Associated with Community-Dwelling Patients with Alzheimer's Disease in Japan: Baseline Results from the Prospective Observational GERAS-J Study. Journal of Alzheimer's Disease, 2020, 74, 127-138.	1.2	14

#	Article	IF	Citations
55	Disturbances in Everyday Life Activities and Sequence Disabilities in Tool use for Alzheimer Disease and Vascular Dementia. Cognitive and Behavioral Neurology, 2009, 22, 215-221.	0.5	13
56	Health economic effect of donepezil treatment for CDR 0.5 converters to Alzheimer's disease as shown by the Markov model. Archives of Gerontology and Geriatrics, 2010, 50, 295-299.	1.4	13
57	Singing therapy can be effective for a patient with severe nonfluent aphasia. International Journal of Rehabilitation Research, 2012, 35, 78-81.	0.7	13
58	Prognosis of Vascular Mild Cognitive Impairment Includes Vascular Dementia Onset and Death by Cardiovascular Disease: Reanalysis From the Osaki-Tajiri Project. Journal of Stroke and Cerebrovascular Diseases, 2012, 21, 607-611.	0.7	13
59	Singing can improve speech function in aphasics associated with intact right basal ganglia and preserve right temporal glucose metabolism: Implications for singing therapy indication. International Journal of Neuroscience, 2016, 126, 39-45.	0.8	13
60	Normative data on Benton Visual Form Discrimination Test for older adults and impaired scores in Clinical Dementia Rating 0.5 participants: Communityâ€based study. The Osaki–Tajiri Project. Psychiatry and Clinical Neurosciences, 2009, 63, 9-16.	1.0	12
61	Decreased Physical Activity Associated with Executive Dysfunction Correlates with Cognitive Impairment among Older Adults in the Community: A Retrospective Analysis from the Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2016, 6, 350-360.	0.6	12
62	Striatal dopaminergic transmission and neocortical glucose utilization in Alzheimer's disease: a triple-tracer positron emission tomography study. Archives of Gerontology and Geriatrics, 2000, 31, 147-158.	1.4	11
63	Elderly Japanese emigrants to Brazil before World War II: I. clinical profiles based on specific historical background. International Journal of Geriatric Psychiatry, 2001, 16, 768-774.	1.3	11
64	Incidence of dementia and cause of death in elderly Japanese emigrants to Brazil before World War II. Archives of Gerontology and Geriatrics, 2011, 52, 75-78.	1.4	11
65	Alzheimer's disease patients institutionalized in group homes run by long-term care insurance exhibit fewer symptoms of behavioural problems as evaluated by the Behavioural Pathology in Alzheimer's Disease Rating Scale. Psychogeriatrics, 2015, 15, 102-108.	0.6	11
66	Early symptoms of <scp>A</scp> lzheimer's disease in <scp>J</scp> apan and <scp>T</scp> aiwan. Geriatrics and Gerontology International, 2016, 16, 797-803.	0.7	11
67	Depression and Dementia in Old-Old Population: History of Depression May Be Associated with Dementia Onset. The Tome Project. Frontiers in Aging Neuroscience, 2017, 9, 335.	1.7	11
68	Clinical Features of Mild Cognitive Impairment and Dementia in a Community: An update of the Osaki-Tajiri Project. Tohoku Journal of Experimental Medicine, 2008, 215, 125-131.	0.5	10
69	Greater Responsiveness to Donepezil in Alzheimer Patients With Higher Levels of Acetylcholinesterase Based on Attention Task Scores and a Donepezil PET Study. Alzheimer Disease and Associated Disorders, 2012, 26, 113-118.	0.6	10
70	Disturbed social recognition and impaired risk judgement in older residents with mild cognitive impairment after the Great East Japan Earthquake of 2011: the Tome Project. Psychogeriatrics, 2016, 16, 349-354.	0.6	10
71	Impaired Eating and Swallowing Function in Older Adults in the Community: The Kurihara Project. International Journal of Environmental Research and Public Health, 2019, 16, 4040.	1.2	10
72	International Report: Local response following the Great East Japan Earthquake 2011. Neurology, 2011, 77, e12-5.	1.5	9

#	Article	IF	CITATIONS
73	Vascular lesions in mixed dementia, vascular dementia, and Alzheimer disease with cerebrovascular disease: The Kurihara Project. Journal of the Neurological Sciences, 2012, 322, 157-160.	0.3	9
74	Overestimation of Self-reported Activities of Daily Living in Vascular Dementia Patients with a Right Hemisphere Lesion. Journal of Stroke and Cerebrovascular Diseases, 2013, 22, 9-14.	0.7	9
75	Normal Hearing Ability but Impaired Auditory Selective Attention Associated with Prediction of Response to Donepezil in Patients with Alzheimer's Disease. Behavioural Neurology, 2015, 2015, 1-6.	1.1	9
76	Sleep disturbance is associated with decreased daily activity and impaired nocturnal reduction of blood pressure in dementia patients. Archives of Gerontology and Geriatrics, 2011, 53, 323-327.	1.4	8
77	Donepezil can improve daily activities and promote rehabilitation for severe Alzheimer's patients in long-term care health facilities. BMC Neurology, 2014, 14, 243.	0.8	8
78	Collage technique may provide new perspectives for Alzheimer patients by exploring messages from their inner world. Dementia E Neuropsychologia, 2009, 3, 299-302.	0.3	7
79	Activated thalamic glucose metabolism after combined donepezil and psychosocial intervention. British Journal of Neuroscience Nursing, 2010, 6, 176-180.	0.1	7
80	The Group Reminiscence Approach Can Increase Self-Awareness ofÂMemory Deficits and Evoke a Life Review in People With Mild Cognitive Impairment: The Kurihara Project Data. Journal of the American Medical Directors Association, 2016, 17, 501-507.	1.2	7
81	Impaired attention function based on the Montréal Cognitive Assessment in vascular dementia patients with frontal hypoperfusion: The Osaki-Tajiri project. Journal of Clinical Neuroscience, 2016, 28, 128-132.	0.8	7
82	Cholinesterase inhibitors are compatible with psychosocial intervention for Alzheimer disease patients suggested by neuroimaging findings. Psychiatry Research - Neuroimaging, 2017, 259, 29-33.	0.9	7
83	Self-Reported Cognitive Decline Based on the Ascertain Dementia 8 Questionnaire May Be Effective for Detecting Mild Cognitive Impairment and Dementia in the Community: The Wakuya Project. Dementia and Geriatric Cognitive Disorders Extra, 2021, 11, 45-50.	0.6	7
84	Global intellectual deterioration in Alzheimer $\hat{a} \in \mathbb{N}$ s disease and a reverse model of intellectual development: An applicability of the Binet scale. Psychiatry and Clinical Neurosciences, 2001, 55, 559-563.	1.0	6
85	Correlation between functional assessment staging and the 'Basic Age' by the Binet scale supports the retrogenesis model of Alzheimer's disease: a preliminary study. Psychogeriatrics, 2003, 3, 82-87.	0.6	6
86	Alzheimer's patients do not show left unilateral spatial neglect but exhibit peripheral inattention and simplification. Dementia E Neuropsychologia, 2007, 1, 374-380.	0.3	6
87	†Residence Is Not Home' Is a Particular Type of Delusion Associated with Cognitive Decline of Alzheimer's Disease. Dementia and Geriatric Cognitive Disorders, 2014, 38, 46-54.	0.7	6
88	Education and the moderating roles of age, sex, ethnicity and apolipoprotein epsilon 4 on the risk of cognitive impairment. Archives of Gerontology and Geriatrics, 2020, 91, 104112.	1.4	6
89	An Environmental Change Does Not Affect Dementia Prevalence but Affects Depressive State and Physical Activity: A Trans-Cultural Study of Japanese Elderly Subjects and Japanese Elderly Immigrants in Brazil. Psychogeriatrics, 2001, 1, 295-302.	0.6	5
90	Behavioral improvement of dementia residents in a group home with an increased number of residents after the Great East Japan Earthquake 2011. International Psychogeriatrics, 2014, 26, 871-872.	0.6	5

#	Article	IF	Citations
91	Costs and resource use of community-dwelling patients with Alzheimer's disease in Japan: 18-month results from the GERAS-J study. Current Medical Research and Opinion, 2021, 37, 1331-1339.	0.9	5
92	Community based measures for managing mild cognitive impairment: The Osaki?Tajiri Project. Psychogeriatrics, 2007, 7, 132-136.	0.6	4
93	Recurrent delusional ideas due to left caudate head infarction, without dementia. Psychogeriatrics, 2012, 12, 58-61.	0.6	4
94	Life concerns of elderly people living at home determined as by <scp>C</scp> ommunity <scp>G</scp> eneral <scp>S</scp> upport <scp>C</scp> enter staff: Implications for organizing a more effective integrated community care system. The <scp>K</scp> urihara <scp>P</scp> roject. Psychogeriatrics, 2014, 14, 188-195.	0.6	4
95	Spontaneous writing in very mild Alzheimer's disease: Analyses of MMSE by older residents in a community. Higher Brain Function Research, 2004, 24, 360-367.	0.0	4
96	Estimated Quality-Adjusted Life-Year Associated with the Degree of Activities of Daily Living in Patients with Alzheimer's Disease. Dementia and Geriatric Cognitive Disorders Extra, 2013, 3, 482-488.	0.6	3
97	Lifetime expectancy and quality-adjusted life-year in Alzheimer's disease with and without cerebrovascular disease: effects of nursing home replacement and donepezil administration – a retrospective analysis in the Tajiri Project. BMC Neurology, 2015, 15, 227.	0.8	3
98	Qualitative Assessment of Instrumental Activities of Daily Living in Older Persons with Very Mild Dementia: The Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2016, 6, 374-381.	0.6	3
99	Prevalence and prognosis of prodromal <scp>A</scp>  zheimer's disease as assessed by magnetic resonance imaging and <sup>18</sup> <scp>F</scp> â€fluorodeoxyglucoseâ€positron emission tomography in a community: reanalysis from the <scp>O</scp> sakiâ€ <scp>T</scp> ajiri <scp>P</scp> roject. Psvchogeriatrics. 2016. 16. 116-120.	0.6	3
100	Improved learning of sequential behaviour during cane gait training or stair climbing after femoral neck fracture: an implication for donepezil for very mild Alzheimer's disease. Psychogeriatrics, 2017, 17, 144-145.	0.6	3
101	Monitoring of drug treatment and psychosocial intervention with SPECT in Alzheimer patients Implications for neurologically appropriate psychosocial interventions. An observational study. The Osaki-Tajiri Project. Dementia E Neuropsychologia, 2018, 12, 380-387.	0.3	3
102	Patients With Very Mild Dementia May Confuse Objective Cognitive Impairments With Subjective Physical Health of Quality of Life: The Tome City Project in Japan. Frontiers in Psychology, 2018, 9, 533.	1.1	3
103	Patient-Reported Outcome Is Important in Psychosocial Intervention for Dementia: A Secondary Analysis of a Randomized Controlled Trial of Group Reminiscence Approach Data. Dementia and Geriatric Cognitive Disorders Extra, 2013, 3, 37-38.	0.6	2
104	Improved functional status by comprehensive physical and psychosocial approach through right insula activation in poststroke vascular dementia. International Journal of Neuroscience, 2013, 123, 698-704.	0.8	2
105	Difficulty with learning of exercise instructions associated with 'working memory' dysfunction and frontal glucose hypometabolism in a patient with very mild subcortical vascular dementia with knee osteoarthritis. BMJ Case Reports, 2013, 2013, bcr2013008577-bcr2013008577.	0.2	2
106	Semantic Dementia Shows both Storage and Access Disorders of Semantic Memory. Behavioural Neurology, 2014, 2014, 1-4.	1.1	2
107	Measures to prevent <scp>A</scp> Izheimer's patients from discontinuing the use of cholinesterase inhibitors. Psychogeriatrics, 2014, 14, 210-211.	0.6	2
108	Necker cube copying may not be appropriate as an examination of dementia: reanalysis from the <scp>T</scp> ajiri Project. Psychogeriatrics, 2016, 16, 298-304.	0.6	2

7

#	Article	IF	CITATIONS
109	Impaired Picture Arrangement subscores (WAIS-III) associated with decreased place orientation and frontal/occipital blood flow in Alzheimer's disease: Implications for social judgment dysfunction. The Osaki-Tajiri Project. Psychiatry Research - Neuroimaging, 2016, 256, 65-69.	0.9	2
110	Language background in early life may be related to neuropsychiatry symptoms in patients with Alzheimer disease. BMC Geriatrics, 2017, 17, 50.	1.1	2
111	Lifetime Expectancy in Dementia with Lewy Bodies: Effects of Donepezil Administration and Special Nursing Home Replacement. A Retrospective Analysis in the Tajiri Project. , 2018, 08, .		2
112	Decreased Time on the Nondominant One-Leg Standing Test Associated with Repeated Falls in Older Residents with Healthy Aging, Mild Cognitive Impairment, and Dementia: The Wakuya Project. Dementia and Geriatric Cognitive Disorders Extra, 2021, 11, 122-128.	0.6	2
113	Depressive symptoms and associated factors in a cognitively normal elderly population: the Tajiri Project., 2001, 16, 780.		2
114	More Delusions May Be Observed in Low-Proficient Multilingual Alzheimer's Disease Patients. PLoS ONE, 2015, 10, e0140714.	1.1	2
115	Subjective memory complaints are not sine qua non as diagnostic criteria for MCI: the Tajiri project. Acta Neurologica Taiwanica, 2006, 15, 55-7.	0.3	2
116	Behavioral neurology in language and aphasia: from basic studies to clinical applications. Acta Medica Indonesiana, 2012, 44, 327-34.	0.9	2
117	Application of Traditional Medical Ideas to Geriatric Syndrome. Advances in Geriatrics, 2014, 2014, 1-20.	1.6	1
118	P4-186: DONEPEZIL CAN IMPROVE DAILY ACTIVITIES AND PROMOTE REHABILITATION FOR SEVERE ALZHEIMER'S PATIENTS IN LONG-TERM CARE HEALTH FACILITIES. , 2014, 10, P857-P857.		1
119	Cognitive Impairment Affects Mortality in Older People in Japan. The Tajiri Project 1991–2005. Journal of the American Geriatrics Society, 2014, 62, 1793-1795.	1.3	1
120	Observational assessment of communication disorders in vascular dementia patients with right hemisphere damage. Psychogeriatrics, 2014, 14, 143-151.	0.6	1
121	Gender's Effects to the Early Symptoms of Alzheimer's Disease in 5 Asian Countries. American Journal of Alzheimer's Disease and Other Dementias, 2017, 32, 194-199.	0.9	1
122	[P1â€"337]: EFFECT OF DRUG TREATMENT AND NURSING HOME RESIDENCY ON LIFE EXPECTANCY IN DEMENTIA (DLB VS AD): A RETROSPECTIVE ANALYSIS IN THE TAJIRI PROJECT. Alzheimer's and Dementia, 2017, 13, P385.	0.4	1
123	Music intonation therapy is effective for speech output in a patient with nonâ€fluent aphasia in a chronic stage. Psychogeriatrics, 2021, 21, 430-433.	0.6	1
124	Humans can only be cared for by humans: Opposition against a recent misunderstanding of the digital society as a future society and describing the importance of †the second person†< scp>QOL. Psychogeriatrics, 2021, 21, 948-949.	0.6	1
125	Kanji and Kana agraphia in mild cognitive impairment and dementia: a trans-cultural comparison of elderly Japanese subjects living in Japan and Brazil. Higher Brain Function Research, 2011, 31, 393-400.	0.0	1
126	P2-189: PREVALENCE AND PROGNOSIS OF "PRODROMAL AD―(DUBOIS ET AL.) ASSESSED WITH MRI AND FDG-PET IN A COMMUNITY: REANALYSIS FROM THE OSAKI-TAJIRI PROJECT. , 2014, 10, P540-P540.		0

#	Article	IF	CITATIONS
127	P3â€019: Decreased Behavioral Abnormalities After Treatment of Combined Donepezil and Yokukansankachimpihange in Alzheimer's Disease: The Osakiâ€Tajiri Project. Alzheimer's and Dementia, 2016, 12, P824.	0.4	0
128	[P2â€"419]: MONITORING OF DRUG TREATMENT AND PSYCHOSOCIAL INTERVENTION WITH THE EZIS SYSTEM ( <sup>99M</sup> TCâ€ECD SPECT) FOR AD PATIENTS: THE OSAKIâ€TAJIRI PROJECT. Alzheimer's and Dementia, 2017, 13, P794.	0.4	0
129	Can dementia patients live in a community? A case of a patient in trouble with the law. Psychogeriatrics, 2019, 19, 605-608.	0.6	0
130	Alzheimer's deterioration in intellectual and neurobiological staging supports the retrogenesis model: a double dissociation between verbal/nonâ€verbal judgments and the left/right parietoâ€temporal glucose metabolism. A retrospective data analysis from the Tajiri Project. Psychogeriatrics, 2020, 20, 149-155.	0.6	0
131	Word Processing Is Faster than Picture Processing in Alzheimer's Disease. Behavioural Neurology, 2020, 2020, 1-8.	1.1	0
132	Mental and Physical Self-Awareness of Alzheimer Patients: Decreased Awareness of Amnesia and Increased Fear of Falling Compared to Views of Families: The Tajiri and Wakuya Projects. Dementia and Geriatric Cognitive Disorders, 2021, 50, 96-102.	0.7	0
133	Reversion to the original concept of clinical practice for patients with dementia: towards an appropriate intervention for a favourable prognosis. Psychogeriatrics, 2021, 21, 447-448.	0.6	0
134	Comprehensive intervention for dementia: Possibility of cognitive rehabilitation. Higher Brain Function Research, 2004, 24, 176-183.	0.0	0
135	Introduction of Induction Heating is too late for older residents with difficulty in handling fire. Dementia E Neuropsychologia, 2022, 16, 115-120.	0.3	0