
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2017353/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Introduction of Induction Heating is too late for older residents with difficulty in handling fire. Dementia E Neuropsychologia, 2022, 16, 115-120.	0.3	Ο
2	Mental and Physical Self-Awareness of Alzheimer Patients: Decreased Awareness of Amnesia and Increased Fear of Falling Compared to Views of Families: The Tajiri and Wakuya Projects. Dementia and Geriatric Cognitive Disorders, 2021, 50, 96-102.	0.7	0
3	Self-Reported Cognitive Decline Based on the Ascertain Dementia 8 Questionnaire May Be Effective for Detecting Mild Cognitive Impairment and Dementia in the Community: The Wakuya Project. Dementia and Geriatric Cognitive Disorders Extra, 2021, 11, 45-50.	0.6	7
4	Music intonation therapy is effective for speech output in a patient with nonâ€fluent aphasia in a chronic stage. Psychogeriatrics, 2021, 21, 430-433.	0.6	1
5	Reversion to the original concept of clinical practice for patients with dementia: towards an appropriate intervention for a favourable prognosis. Psychogeriatrics, 2021, 21, 447-448.	0.6	0
6	Costs and resource use of community-dwelling patients with Alzheimer's disease in Japan: 18-month results from the GERAS-J study. Current Medical Research and Opinion, 2021, 37, 1331-1339.	0.9	5
7	Decreased Time on the Nondominant One-Leg Standing Test Associated with Repeated Falls in Older Residents with Healthy Aging, Mild Cognitive Impairment, and Dementia: The Wakuya Project. Dementia and Geriatric Cognitive Disorders Extra, 2021, 11, 122-128.	0.6	2
8	Humans can only be cared for by humans: Opposition against a recent misunderstanding of the digital society as a future society and describing the importance of †the second person' <scp>QOL</scp> . Psychogeriatrics, 2021, 21, 948-949.	0.6	1
9	Alzheimer's deterioration in intellectual and neurobiological staging supports the retrogenesis model: a double dissociation between verbal/nonâ€verbal judgments and the left/right parietoâ€temporal glucose metabolism. A retrospective data analysis from the Tajiri Project. Psychogeriatrics, 2020, 20, 149-155.	0.6	0
10	Education and the moderating roles of age, sex, ethnicity and apolipoprotein epsilon 4 on the risk of cognitive impairment. Archives of Gerontology and Geriatrics, 2020, 91, 104112.	1.4	6
11	Does parity matter in women's risk of dementia? A COSMIC collaboration cohort study. BMC Medicine, 2020, 18, 210.	2.3	21
12	Word Processing Is Faster than Picture Processing in Alzheimer's Disease. Behavioural Neurology, 2020, 2020, 1-8.	1.1	0
13	Costs and Resource Use Associated with Community-Dwelling Patients with Alzheimer's Disease in Japan: Baseline Results from the Prospective Observational GERAS-J Study. Journal of Alzheimer's Disease, 2020, 74, 127-138.	1.2	14
14	Determinants of cognitive performance and decline in 20 diverse ethno-regional groups: A COSMIC collaboration cohort study. PLoS Medicine, 2019, 16, e1002853.	3.9	86
15	Impaired Eating and Swallowing Function in Older Adults in the Community: The Kurihara Project. International Journal of Environmental Research and Public Health, 2019, 16, 4040.	1.2	10
16	Can dementia patients live in a community? A case of a patient in trouble with the law. Psychogeriatrics, 2019, 19, 605-608.	0.6	0
17	Vascular Mild Cognitive Impairment: Identifying Disease in Community-Dwelling Older Adults, Reducing Risk Factors, and Providing Support. The Osaki-Tajiri and Kurihara Projects. Journal of Alzheimer's Disease, 2019, 70, S293-S302.	1.2	14
18	Fear of Falling and Cognitive Impairments in Elderly People with Hip Fractures. Dementia and Geriatric Cognitive Disorders Extra. 2018, 7, 386-394.	0.6	16

#	Article	IF	CITATIONS
19	Monitoring of drug treatment and psychosocial intervention with SPECT in Alzheimer patients Implications for neurologically appropriate psychosocial interventions. An observational study. The Osaki-Tajiri Project. Dementia E Neuropsychologia, 2018, 12, 380-387.	0.3	3
20	Lifetime Expectancy in Dementia with Lewy Bodies: Effects of Donepezil Administration and Special Nursing Home Replacement. A Retrospective Analysis in the Tajiri Project. , 2018, 08, .		2
21	Decreased Behavioral Abnormalities After Treatment with Combined Donepezil and Yokukansankachimpihange in Alzheimer Disease: An Observational Study. The Osaki-Tajiri Project. Neurology and Therapy, 2018, 7, 333-340.	1.4	16
22	Patients With Very Mild Dementia May Confuse Objective Cognitive Impairments With Subjective Physical Health of Quality of Life: The Tome City Project in Japan. Frontiers in Psychology, 2018, 9, 533.	1.1	3
23	Speaking one more language in early life has only minor effects on cognition in Taiwanese with low education level: the Taishan Project. Psychogeriatrics, 2017, 17, 256-261.	0.6	14
24	Language background in early life may be related to neuropsychiatry symptoms in patients with Alzheimer disease. BMC Geriatrics, 2017, 17, 50.	1.1	2
25	Gender's Effects to the Early Symptoms of Alzheimer's Disease in 5 Asian Countries. American Journal of Alzheimer's Disease and Other Dementias, 2017, 32, 194-199.	0.9	1
26	Cholinesterase inhibitors are compatible with psychosocial intervention for Alzheimer disease patients suggested by neuroimaging findings. Psychiatry Research - Neuroimaging, 2017, 259, 29-33.	0.9	7
27	Improved learning of sequential behaviour during cane gait training or stair climbing after femoral neck fracture: an implication for donepezil for very mild Alzheimer's disease. Psychogeriatrics, 2017, 17, 144-145.	0.6	3
28	[P2–419]: MONITORING OF DRUG TREATMENT AND PSYCHOSOCIAL INTERVENTION WITH THE EZIS SYSTEM ( <sup>99M</sup> TCâ€ECD SPECT) FOR AD PATIENTS: THE OSAKIâ€TAJIRI PROJECT. Alzheimer's and Dementia, 2017, 13, P794.	0.4	0
29	[P1â€"337]: EFFECT OF DRUG TREATMENT AND NURSING HOME RESIDENCY ON LIFE EXPECTANCY IN DEMENTIA (DLB VS AD): A RETROSPECTIVE ANALYSIS IN THE TAJIRI PROJECT. Alzheimer's and Dementia, 2017, 13, P385.	0.4	1
30	Depression and Dementia in Old-Old Population: History of Depression May Be Associated with Dementia Onset. The Tome Project. Frontiers in Aging Neuroscience, 2017, 9, 335.	1.7	11
31	Necker cube copying may not be appropriate as an examination of dementia: reanalysis from the <scp>T</scp> ajiri Project. Psychogeriatrics, 2016, 16, 298-304.	0.6	2
32	Early symptoms of <scp>A</scp> lzheimer's disease in <scp>J</scp> apan and <scp>T</scp> aiwan. Geriatrics and Gerontology International, 2016, 16, 797-803.	0.7	11
33	Disturbed social recognition and impaired risk judgement in older residents with mild cognitive impairment after the Great East Japan Earthquake of 2011: the Tome Project. Psychogeriatrics, 2016, 16, 349-354.	0.6	10
34	P3â€019: Decreased Behavioral Abnormalities After Treatment of Combined Donepezil and Yokukansankachimpihange in Alzheimer's Disease: The Osakiâ€Tajiri Project. Alzheimer's and Dementia, 2016, 12, P824.	0.4	0
35	The Group Reminiscence Approach Can Increase Self-Awareness ofÂMemory Deficits and Evoke a Life Review in People With Mild Cognitive Impairment: The Kurihara Project Data. Journal of the American Medical Directors Association, 2016, 17, 501-507.	1.2	7
36	Impaired Picture Arrangement subscores (WAIS-III) associated with decreased place orientation and frontal/occipital blood flow in Alzheimer's disease: Implications for social judgment dysfunction. The Osaki-Tajiri Project. Psychiatry Research - Neuroimaging, 2016, 256, 65-69.	0.9	2

#	Article	IF	CITATIONS
37	Qualitative Assessment of Instrumental Activities of Daily Living in Older Persons with Very Mild Dementia: The Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2016, 6, 374-381.	0.6	3
38	Neuroepidemiologic and Neurobehavioral Characteristics of Motoric Cognitive Risk Syndrome in an Old-Old Population: The Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2016, 6, 176-182.	0.6	22
39	Decreased Physical Activity Associated with Executive Dysfunction Correlates with Cognitive Impairment among Older Adults in the Community: A Retrospective Analysis from the Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2016, 6, 350-360.	0.6	12
40	Prevalence and prognosis of prodromal <scp>A</scp> lzheimer's disease as assessed by magnetic resonance imaging and <sup>18</sup> <scp>F</scp> â€fluorodeoxyglucoseâ€positron emission tomography in a community: reanalysis from the <scp>O</scp> sakiâ€ <scp>T</scp> ajiri <scp>P</scp> roject. Psychogeriatrics, 2016, 16, 116-120.	0.6	3
41	Singing can improve speech function in aphasics associated with intact right basal ganglia and preserve right temporal glucose metabolism: Implications for singing therapy indication. International Journal of Neuroscience, 2016, 126, 39-45.	0.8	13
42	Impaired attention function based on the Montréal Cognitive Assessment in vascular dementia patients with frontal hypoperfusion: The Osaki-Tajiri project. Journal of Clinical Neuroscience, 2016, 28, 128-132.	0.8	7
43	Lifetime expectancy and quality-adjusted life-year in Alzheimer's disease with and without cerebrovascular disease: effects of nursing home replacement and donepezil administration – a retrospective analysis in the Tajiri Project. BMC Neurology, 2015, 15, 227.	0.8	3
44	Alzheimer's disease patients institutionalized in group homes run by long-term care insurance exhibit fewer symptoms of behavioural problems as evaluated by the Behavioural Pathology in Alzheimer's Disease Rating Scale. Psychogeriatrics, 2015, 15, 102-108.	0.6	11
45	Normal Hearing Ability but Impaired Auditory Selective Attention Associated with Prediction of Response to Donepezil in Patients with Alzheimer's Disease. Behavioural Neurology, 2015, 2015, 1-6.	1.1	9
46	A Cluster Randomized Controlled Trial of Nonpharmacological Interventions for Old-Old Subjects with a Clinical Dementia Rating of 0.5: The Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2015, 5, 221-232.	0.6	22
47	More Delusions May Be Observed in Low-Proficient Multilingual Alzheimer's Disease Patients. PLoS ONE, 2015, 10, e0140714.	1.1	2
48	Application of Traditional Medical Ideas to Geriatric Syndrome. Advances in Geriatrics, 2014, 2014, 1-20.	1.6	1
49	Mild cognitive impairment, poor episodic memory, and late-life depression are associated with cerebral cortical thinning and increased white matter hyperintensities. Frontiers in Aging Neuroscience, 2014, 6, 306.	1.7	59
50	Semantic Dementia Shows both Storage and Access Disorders of Semantic Memory. Behavioural Neurology, 2014, 2014, 1-4.	1.1	2
51	P4-186: DONEPEZIL CAN IMPROVE DAILY ACTIVITIES AND PROMOTE REHABILITATION FOR SEVERE ALZHEIMER'S PATIENTS IN LONG-TERM CARE HEALTH FACILITIES. , 2014, 10, P857-P857.		1
52	Cognitive Impairment Affects Mortality in Older People in Japan. The Tajiri Project 1991–2005. Journal of the American Geriatrics Society, 2014, 62, 1793-1795.	1.3	1
53	Observational assessment of communication disorders in vascular dementia patients with right hemisphere damage. Psychogeriatrics, 2014, 14, 143-151.	0.6	1
54	Life concerns of elderly people living at home determined as by <scp>C</scp> ommunity <scp>G</scp> eneral <scp>S</scp> upport <scp>C</scp> enter staff: Implications for organizing a more effective integrated community care system. The <scp>K</scp> urihara <scp>P</scp> roject. Psychogeriatrics, 2014, 14, 188-195.	0.6	4

#	Article	IF	CITATIONS
55	Motoric cognitive risk syndrome. Neurology, 2014, 83, 718-726.	1.5	345
56	Disorders of "taste cognition―are associated with insular involvement in patients with Alzheimer's disease and vascular dementia: "Memory of food is impaired in dementia and responsible for poor diet― International Psychogeriatrics, 2014, 26, 1127-1138.	0.6	28
57	Donepezil can improve daily activities and promote rehabilitation for severe Alzheimer's patients in long-term care health facilities. BMC Neurology, 2014, 14, 243.	0.8	8
58	Behavioral improvement of dementia residents in a group home with an increased number of residents after the Great East Japan Earthquake 2011. International Psychogeriatrics, 2014, 26, 871-872.	0.6	5
59	â€~Residence Is Not Home' Is a Particular Type of Delusion Associated with Cognitive Decline of Alzheimer's Disease. Dementia and Geriatric Cognitive Disorders, 2014, 38, 46-54.	0.7	6
60	Measures to prevent <scp>A</scp> lzheimer's patients from discontinuing the use of cholinesterase inhibitors. Psychogeriatrics, 2014, 14, 210-211.	0.6	2
61	Donepezil and life expectancy in Alzheimer's disease: A retrospective analysis in the Tajiri Project. BMC Neurology, 2014, 14, 83.	0.8	33
62	P2-189: PREVALENCE AND PROGNOSIS OF "PRODROMAL AD―(DUBOIS ET AL.) ASSESSED WITH MRI AND FDG-PET IN A COMMUNITY: REANALYSIS FROM THE OSAKI-TAJIRI PROJECT. , 2014, 10, P540-P540.		0
63	Overestimation of Self-reported Activities of Daily Living in Vascular Dementia Patients with a Right Hemisphere Lesion. Journal of Stroke and Cerebrovascular Diseases, 2013, 22, 9-14.	0.7	9
64	Vascular dementia with left thalamic infarction: Neuropsychological and behavioral implications suggested by involvement of the thalamic nucleus and the remote effect on cerebral cortex. The Osaki–Tajiri project. Psychiatry Research - Neuroimaging, 2013, 213, 56-62.	0.9	14
65	Patient-Reported Outcome Is Important in Psychosocial Intervention for Dementia: A Secondary Analysis of a Randomized Controlled Trial of Group Reminiscence Approach Data. Dementia and Geriatric Cognitive Disorders Extra, 2013, 3, 37-38.	0.6	2
66	Improved functional status by comprehensive physical and psychosocial approach through right insula activation in poststroke vascular dementia. International Journal of Neuroscience, 2013, 123, 698-704.	0.8	2
67	Clinical utility of the <scp>F</scp> unctional <scp>I</scp> ndependence <scp>M</scp> easure for assessment of patients with <scp>A</scp> lzheimer's disease and vascular dementia. Psychogeriatrics, 2013, 13, 199-205.	0.6	14
68	Apathy is more severe in vascular than amnestic mild cognitive impairment in a community: The <scp>K</scp> urihara <scp>P</scp> roject. Psychiatry and Clinical Neurosciences, 2013, 67, 517-525.	1.0	15
69	Content of delusional thoughts in Alzheimer's disease and assessment of content-specific brain dysfunctions with BEHAVE-AD-FW and SPECT. International Psychogeriatrics, 2013, 25, 939-948.	0.6	16
70	Estimated Quality-Adjusted Life-Year Associated with the Degree of Activities of Daily Living in Patients with Alzheimer's Disease. Dementia and Geriatric Cognitive Disorders Extra, 2013, 3, 482-488.	0.6	3
71	Difficulty with learning of exercise instructions associated with 'working memory' dysfunction and frontal glucose hypometabolism in a patient with very mild subcortical vascular dementia with knee osteoarthritis. BMJ Case Reports, 2013, 2013, bcr2013008577-bcr2013008577.	0.2	2
72	Greater Responsiveness to Donepezil in Alzheimer Patients With Higher Levels of Acetylcholinesterase Based on Attention Task Scores and a Donepezil PET Study. Alzheimer Disease and Associated Disorders, 2012, 26, 113-118.	0.6	10

#	Article	IF	CITATIONS
73	Singing therapy can be effective for a patient with severe nonfluent aphasia. International Journal of Rehabilitation Research, 2012, 35, 78-81.	0.7	13
74	Screening for Very Mild Subcortical Vascular Dementia Patients Aged 75 and Above Using the Montreal Cognitive Assessment and Mini-Mental State Examination in a Community: The Kurihara Project. Dementia and Geriatric Cognitive Disorders Extra, 2012, 2, 503-515.	0.6	25
75	Prognosis of Vascular Mild Cognitive Impairment Includes Vascular Dementia Onset and Death by Cardiovascular Disease: Reanalysis From the Osaki-Tajiri Project. Journal of Stroke and Cerebrovascular Diseases, 2012, 21, 607-611.	0.7	13
76	Impaired memory and executive function associated with decreased medial temporal and prefrontal blood flow in Clinical Dementia Rating 0.5 status: the Osakiâ€īajiri project. Psychogeriatrics, 2012, 12, 27-33.	0.6	15
77	Recurrent delusional ideas due to left caudate head infarction, without dementia. Psychogeriatrics, 2012, 12, 58-61.	0.6	4
78	Impaired instrumental activities of daily living affect conversion from mild cognitive impairment to dementia: the Osakiâ€Tajiri Project. Psychogeriatrics, 2012, 12, 34-42.	0.6	15
79	Silent Infarction or White Matter Hyperintensity and Impaired Attention Task Scores in a Nondemented Population: The Osaki-Tajiri Project. Journal of Stroke and Cerebrovascular Diseases, 2012, 21, 275-282.	0.7	21
80	Prevalence of dementia and dementing diseases in the oldâ€old population in Japan: the Kurihara Project. Implications for Longâ€Term Care Insurance data. Psychogeriatrics, 2012, 12, 226-234.	0.6	82
81	Vascular lesions in mixed dementia, vascular dementia, and Alzheimer disease with cerebrovascular disease: The Kurihara Project. Journal of the Neurological Sciences, 2012, 322, 157-160.	0.3	9
82	Trends in the Prevalence of Dementia in Japan. International Journal of Alzheimer's Disease, 2012, 2012, 1-11.	1.1	73
83	Behavioral neurology in language and aphasia: from basic studies to clinical applications. Acta Medica Indonesiana, 2012, 44, 327-34.	0.9	2
84	Chronic Kidney Disease: A Risk Factor for Dementia Onset: A Populationâ€Based Study. The Osakiâ€Tajiri Project. Journal of the American Geriatrics Society, 2011, 59, 1175-1181.	1.3	53
85	Improved social interaction and increased anterior cingulate metabolism after group reminiscence with reality orientation approach for vascular dementia. Psychiatry Research - Neuroimaging, 2011, 192, 183-187.	0.9	35
86	Sleep disturbance is associated with decreased daily activity and impaired nocturnal reduction of blood pressure in dementia patients. Archives of Gerontology and Geriatrics, 2011, 53, 323-327.	1.4	8
87	Incidence of dementia and cause of death in elderly Japanese emigrants to Brazil before World War II. Archives of Gerontology and Geriatrics, 2011, 52, 75-78.	1.4	11
88	International Report: Local response following the Great East Japan Earthquake 2011. Neurology, 2011, 77, e12-5.	1.5	9
89	Kanji and Kana agraphia in mild cognitive impairment and dementia : a trans-cultural comparison of elderly Japanese subjects living in Japan and Brazil. Higher Brain Function Research, 2011, 31, 393-400.	0.0	1
90	Health economic effect of donepezil treatment for CDR 0.5 converters to Alzheimer's disease as shown by the Markov model. Archives of Gerontology and Geriatrics, 2010, 50, 295-299.	1.4	13

#	Article	IF	CITATIONS
91	Activated thalamic glucose metabolism after combined donepezil and psychosocial intervention. British Journal of Neuroscience Nursing, 2010, 6, 176-180.	0.1	7
92	Collage technique may provide new perspectives for Alzheimer patients by exploring messages from their inner world. Dementia E Neuropsychologia, 2009, 3, 299-302.	0.3	7
93	Prevalence of Possible Idiopathic Normal-Pressure Hydrocephalus in Japan: The Osaki-Tajiri Project. Neuroepidemiology, 2009, 32, 171-175.	1.1	149
94	Combined Memory and Executive Function Tests Can Screen Mild Cognitive Impairment and Converters to Dementia in a Community: The Osaki-Tajiri Project. Neuroepidemiology, 2009, 33, 103-110.	1.1	42
95	Normative data on Benton Visual Form Discrimination Test for older adults and impaired scores in Clinical Dementia Rating 0.5 participants: Communityâ€based study. The Osaki–Tajiri Project. Psychiatry and Clinical Neurosciences, 2009, 63, 9-16.	1.0	12
96	Disturbances in Everyday Life Activities and Sequence Disabilities in Tool use for Alzheimer Disease and Vascular Dementia. Cognitive and Behavioral Neurology, 2009, 22, 215-221.	0.5	13
97	Cross-cultural comparisons of the Mini-mental State Examination between Japanese and U.S. cohorts. International Psychogeriatrics, 2009, 21, 113.	0.6	25
98	Decreased cortical glucose metabolism in converters from CDR 0.5 to Alzheimer's disease in a community: the Osaki-Tajiri Project. International Psychogeriatrics, 2009, 21, 148.	0.6	22
99	Comprehensive approach of donepezil and psychosocial interventions on cognitive function and quality of life for Alzheimer's disease: the Osaki-Tajiri Project. Age and Ageing, 2008, 37, 469-473.	0.7	31
100	Clinical Features of Mild Cognitive Impairment and Dementia in a Community: An update of the Osaki-Tajiri Project. Tohoku Journal of Experimental Medicine, 2008, 215, 125-131.	0.5	10
101	A Randomized Controlled Trial of the Group Reminiscence Approach in Patients with Vascular Dementia. Dementia and Geriatric Cognitive Disorders, 2007, 24, 48-54.	0.7	34
102	Incidence of dementia and associated risk factors in Japan: The Osaki-Tajiri Project. Journal of the Neurological Sciences, 2007, 260, 175-182.	0.3	67
103	Alzheimer's patients do not show left unilateral spatial neglect but exhibit peripheral inattention and simplification. Dementia E Neuropsychologia, 2007, 1, 374-380.	0.3	6
104	Community based measures for managing mild cognitive impairment: The Osaki?Tajiri Project. Psychogeriatrics, 2007, 7, 132-136.	0.6	4
105	Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS–ADRDA criteria. Lancet Neurology, The, 2007, 6, 734-746.	4.9	3,755
106	Non-verbal learning is impaired in very mild Alzheimer's disease (CDR 0.5): Normative data from the learning version of the Rey-Osterrieth Complex Figure Test. Psychiatry and Clinical Neurosciences, 2006, 60, 139-146.	1.0	49
107	Effect of age and education on the Trail Making Test and determination of normative data for Japanese elderly people: The Tajiri Project. Psychiatry and Clinical Neurosciences, 2006, 60, 422-428.	1.0	89
108	Necker cube copying ability in normal elderly and Alzheimer's disease. A community-based study: The Tajiri project. Psychogeriatrics, 2006, 6, 4-9.	0.6	26

#	Article	IF	CITATIONS
109	Different MRI findings for normal elderly and very mild Alzheimer's disease in a community: Implications for clinical practice. Archives of Gerontology and Geriatrics, 2006, 42, 59-71.	1.4	18
110	Subjective memory complaints are not sine qua non as diagnostic criteria for MCI: the Tajiri project. Acta Neurologica Taiwanica, 2006, 15, 55-7.	0.3	2
111	Neuropsychosocial Features of Very Mild Alzheimer's Disease (CDR 0.5) and Progression to Dementia in a Community: The Tajiri Project. Journal of Geriatric Psychiatry and Neurology, 2004, 17, 183-189.	1.2	18
112	Risperidone is Effective for Wandering and Disturbed Sleep/Wake Patterns in Alzheimer's Disease. Journal of Geriatric Psychiatry and Neurology, 2004, 17, 61-67.	1.2	54
113	Executive dysfunction can explain word-list learning disability in very mild Alzheimer's disease: The Tajiri Project. Psychiatry and Clinical Neurosciences, 2004, 58, 54-60.	1.0	23
114	Prevalence and Cognitive Performances of Clinical Dementia Rating 0.5 and Mild Cognitive Impairment in Japan. Alzheimer Disease and Associated Disorders, 2004, 18, 3-10.	0.6	70
115	Spontaneous writing in very mild Alzheimer's disease: Analyses of MMSE by older residents in a community. Higher Brain Function Research, 2004, 24, 360-367.	0.0	4
116	Comprehensive intervention for dementia: Possibility of cognitive rehabilitation. Higher Brain Function Research, 2004, 24, 176-183.	0.0	0
117	Decreased striatal D2 receptor density associated with severe behavioral abnormality in Alzheimer's disease. Annals of Nuclear Medicine, 2003, 17, 567-573.	1.2	68
118	Language deterioration in four Japanese-Portuguese bilingual patients with Alzheimer's disease: a trans-cultural study of Japanese elderly immigrants in Brazil. Psychogeriatrics, 2003, 3, 63-68.	0.6	26
119	Correlation between functional assessment staging and the 'Basic Age' by the Binet scale supports the retrogenesis model of Alzheimer's disease: a preliminary study. Psychogeriatrics, 2003, 3, 82-87.	0.6	6
120	Corpus Callosum Atrophy, White Matter Lesions, and Frontal Executive Dysfunction in Normal Aging and Alzheimer's Disease. A Community-Based Study: The Tajiri Project. International Psychogeriatrics, 2003, 15, 9-25.	0.6	35
121	Prevalence of Dementia and Dementing Diseases in Japan. Archives of Neurology, 2002, 59, 1109.	4.9	126
122	Therapeutic Psychosocial Intervention for Elderly Subjects With Very Mild Alzheimer Disease in a Community: The Tajiri Project. Alzheimer Disease and Associated Disorders, 2002, 16, 261-269.	0.6	39
123	Global intellectual deterioration in Alzheimer's disease and a reverse model of intellectual development: An applicability of the Binet scale. Psychiatry and Clinical Neurosciences, 2001, 55, 559-563.	1.0	6
124	Cognitive function and frontal lobe atrophy in normal elderly adults: Implications for dementia not as aging-related disorders and the reserve hypothesis. Psychiatry and Clinical Neurosciences, 2001, 55, 565-572.	1.0	60
125	Elderly Japanese emigrants to Brazil before World War II: I. clinical profiles based on specific historical background. International Journal of Geriatric Psychiatry, 2001, 16, 768-774.	1.3	11
126	Elderly Japanese emigrants to Brazil before World War II: II. prevalence of senile dementia. International Journal of Geriatric Psychiatry, 2001, 16, 775-779.	1.3	17

#	Article	IF	CITATIONS
127	Depressive symptoms and associated factors in a cognitively normal elderly population: the Tajiri Project. International Journal of Geriatric Psychiatry, 2001, 16, 780-788.	1.3	25
128	An Environmental Change Does Not Affect Dementia Prevalence but Affects Depressive State and Physical Activity: A Trans-Cultural Study of Japanese Elderly Subjects and Japanese Elderly Immigrants in Brazil. Psychogeriatrics, 2001, 1, 295-302.	0.6	5
129	Depressive symptoms and associated factors in a cognitively normal elderly population: the Tajiri Project. , 2001, 16, 780.		2
130	The effects of group work therapy in patients with Alzheimer's disease. International Journal of Geriatric Psychiatry, 2000, 15, 532-534.	1.3	14
131	Striatal dopaminergic transmission and neocortical glucose utilization in Alzheimer's disease: a triple-tracer positron emission tomography study. Archives of Gerontology and Geriatrics, 2000, 31, 147-158.	1.4	11
132	Prevalence of senile dementia in a rural community in Japan: the Tajiri project. Archives of Gerontology and Geriatrics, 2000, 29, 249-265.	1.4	23
133	Impaired verbal description ability assessed by the Picture Description Task in Alzheimer's disease. Archives of Gerontology and Geriatrics, 1998, 27, 57-65.	1.4	19
134	Cortical glucose metabolism in psychiatric wandering patients with vascular dementia. Psychiatry Research - Neuroimaging, 1996, 67, 71-80.	0.9	25
135	Cerebral circulation and oxygen metabolism associated with subclinical periventricular hyperintensity as shown by magnetic resonance imaging. Annals of Neurology, 1990, 28, 378-383.	2.8	142