

Ana Carolina J Vasques

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2022713/publications.pdf>

Version: 2024-02-01

35
papers

1,035
citations

687220

13
h-index

434063

31
g-index

36
all docs

36
docs citations

36
times ranked

1915
citing authors

#	ARTICLE	IF	CITATIONS
1	TyG index performs better than HOMA in a Brazilian population: A hyperglycemic clamp validated study. <i>Diabetes Research and Clinical Practice</i> , 2011, 93, e98-e100.	1.1	380
2	Neck circumference as a simple tool for identifying the metabolic syndrome and insulin resistance: results from the Brazilian Metabolic Syndrome Study. <i>Clinical Endocrinology</i> , 2013, 78, 874-881.	1.2	157
3	Índice de massa corporal e circunferência abdominal: associação com fatores de risco cardiovascular. <i>Arquivos Brasileiros De Cardiologia</i> , 2006, 87, 728-34.	0.3	77
4	Obesity and Excess Protein and Carbohydrate Consumption Are Risk Factors for Thyroid Cancer. <i>Nutrition and Cancer</i> , 2012, 64, 1190-1195.	0.9	49
5	The HOMA-Adiponectin (HOMA-AD) Closely Mirrors the HOMA-IR Index in the Screening of Insulin Resistance in the Brazilian Metabolic Syndrome Study (BRAMS). <i>PLoS ONE</i> , 2016, 11, e0158751.	1.1	36
6	Association of Sleep Deprivation With Reduction in Insulin Sensitivity as Assessed by the Hyperglycemic Clamp Technique in Adolescents. <i>JAMA Pediatrics</i> , 2016, 170, 487.	3.3	35
7	Impaired incretin secretion and pancreatic dysfunction with older age and diabetes. <i>Metabolism: Clinical and Experimental</i> , 2014, 63, 922-929.	1.5	32
8	Utilização de medidas antropométricas para a avaliação do acúmulo de gordura visceral. <i>Revista De Nutricao</i> , 2010, 23, 107-118.	0.4	29
9	Changes in serum levels of lipopolysaccharides and CD26 in patients with Crohn's disease. <i>Intestinal Research</i> , 2017, 15, 352.	1.0	28
10	Sagittal Abdominal Diameter as a Surrogate Marker of Insulin Resistance in an Admixed Population – Brazilian Metabolic Syndrome Study (BRAMS). <i>PLoS ONE</i> , 2015, 10, e0125365.	1.1	22
11	Influence of selenium supplementation on patients with inflammation: A pilot double blind randomized study. <i>Nutrition</i> , 2017, 41, 32-36.	1.1	17
12	Diferentes aferições do diâmetro abdominal sagital e do perímetro da cintura na predição do HOMA-IR. <i>Arquivos Brasileiros De Cardiologia</i> , 2009, 93, 511-518.	0.3	15
13	Prevalence of hepatitis B and hepatitis C among diabetes mellitus type 2 individuals. <i>PLoS ONE</i> , 2019, 14, e0211193.	1.1	14
14	Glucose Metabolism Parameters and Post-Prandial GLP-1 and GLP-2 Release Largely Vary in Several Distinct Situations: a Controlled Comparison Among Individuals with Crohn's Disease and Individuals with Obesity Before and After Bariatric Surgery. <i>Obesity Surgery</i> , 2018, 28, 378-388.	1.1	12
15	Homeostatic model assessment of adiponectin (HOMA-Adiponectin) as a surrogate measure of insulin resistance in adolescents: Comparison with the hyperglycaemic clamp and homeostatic model assessment of insulin resistance. <i>PLoS ONE</i> , 2019, 14, e0214081.	1.1	12
16	β-Cell Function Improvements in Grade I/II Obese Subjects With Type 2 Diabetes 1 Month After Biliopancreatic Diversion: Results from modeling analyses of oral glucose tolerance tests and hyperglycemic clamp studies. <i>Diabetes Care</i> , 2013, 36, 4117-4124.	4.3	10
17	Epicardial and Pericardial Fat in Type 2 Diabetes: Favourable Effects of Biliopancreatic Diversion. <i>Obesity Surgery</i> , 2015, 25, 477-485.	1.1	10
18	Hypertriglyceridemic Waist Phenotype Indicates Insulin Resistance in Adolescents According to the Clamp Technique in the BRAMS Study. <i>Childhood Obesity</i> , 2016, 12, 446-454.	0.8	10

#	ARTICLE	IF	CITATIONS
19	Accuracy of predictive equations versus indirect calorimetry for the evaluation of energy expenditure in cancer patients with solid tumors – An integrative systematic review study. <i>Clinical Nutrition ESPEN</i> , 2020, 35, 12-19.	0.5	10
20	Sagittal abdominal diameter as a marker for epicardial adipose tissue in premenopausal women. <i>Metabolism: Clinical and Experimental</i> , 2013, 62, 1032-1036.	1.5	9
21	Serum levels of retinol binding protein 4 in women with different levels of adiposity and glucose tolerance. <i>Arquivos Brasileiros De Endocrinologia E Metabologia</i> , 2014, 58, 709-714.	1.3	8
22	Sagittal abdominal diameter resembles waist circumference as a surrogate marker of insulin resistance in adolescents-Brazilian Metabolic Syndrome Study. <i>Pediatric Diabetes</i> , 2018, 19, 882-891.	1.2	8
23	Irritable bowel syndrome: associations between FODMAPS intake, problematic foods, adiposity, and gastrointestinal symptoms. <i>European Journal of Clinical Nutrition</i> , 2019, 73, 637-641.	1.3	7
24	Long-Term Outcomes of Biliopancreatic Diversion on Glycemic Control, Insulin Sensitivity and Beta Cell Function. <i>Obesity Surgery</i> , 2016, 26, 2572-2580.	1.1	5
25	Transthyretin levels: Potential biomarker for monitoring nutritional support efficacy and clinical complications risk in patients receiving parenteral nutrition. <i>Clinical Nutrition ESPEN</i> , 2018, 24, 134-139.	0.5	5
26	PREVALENCE OF HEPATIC STEATOSIS AMONG CHILDREN AND ADOLESCENTS WITH CYSTIC FIBROSIS AND ITS ASSOCIATION WITH NUTRITIONAL STATUS. <i>Revista Paulista De Pediatria</i> , 2019, 37, 435-441.	0.4	5
27	Maternal and paternal obesity are associated with offspring obestatin levels in the Nutritionists™ Health Study. <i>Nutrition</i> , 2021, 83, 111067.	1.1	5
28	Effect of biliopancreatic diversion on sleep quality and daytime sleepiness in patients with obesity and type 2 diabetes. <i>Archives of Endocrinology and Metabolism</i> , 2017, 61, 623-627.	0.3	4
29	Inflammatory process of patients receiving parenteral nutrition is not exclusively responsible for low selenium and glutathione peroxidase levels. <i>Nutrition</i> , 2019, 61, 202-207.	1.1	4
30	FODMAP project: Development, validation and reproducibility of a short food frequency questionnaire to estimate consumption of fermentable carbohydrates. <i>Clinical Nutrition</i> , 2021, 40, 3409-3420.	2.3	4
31	Adiposity and family history of type 2 diabetes in an admixed population of adolescents: Associations with insulin sensitivity, beta-cell function, and hepatic insulin extraction in BRAMS study. <i>Diabetes Research and Clinical Practice</i> , 2018, 137, 72-82.	1.1	3
32	NUTRITIONAL STATUS, QUALITY OF LIFE AND LIFE HABITS OF WOMEN WITH IRRITABLE BOWEL SYNDROME: A CASE-CONTROL STUDY. <i>Arquivos De Gastroenterologia</i> , 2020, 57, 114-120.	0.3	3
33	Validade de equações preditivas para estimativa do gasto energético de repouso em mulheres com diferentes fenótipos metabólicos e de adiposidade participantes do nutritionists health study - nutrihs. , 0, , .		2
34	Energy Expenditure in 21-Hydroxylase Congenital Adrenal Hyperplasia Patients and Comparison with Predictive Equations. <i>Endocrine Practice</i> , 2020, 26, 388-398.	1.1	0
35	Reply – Letter to the editor. <i>Clinical Nutrition ESPEN</i> , 2020, 38, 286-287.	0.5	0