

Ji-Xin Shi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2030767/publications.pdf>

Version: 2024-02-01

12
papers

474
citations

933447

10
h-index

1199594

12
g-index

12
all docs

12
docs citations

12
times ranked

772
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Biochanin A Reduces Inflammatory Injury and Neuronal Apoptosis following Subarachnoid Hemorrhage via Suppression of the TLRs/TIRAP/MyD88/NF- κ B Pathway. <i>Behavioural Neurology</i> , 2018, 2018, 1-10. | 2.1 | 41 |
| 2 | Roles of Pannexin-1 Channels in Inflammatory Response through the TLRs/NF-Kappa B Signaling Pathway Following Experimental Subarachnoid Hemorrhage in Rats. <i>Frontiers in Molecular Neuroscience</i> , 2017, 10, 175. | 2.9 | 46 |
| 3 | Fisetin alleviates early brain injury following experimental subarachnoid hemorrhage in rats possibly by suppressing TLR 4/NF- κ B signaling pathway. <i>Brain Research</i> , 2015, 1629, 250-259. | 2.2 | 40 |
| 4 | Decreased progranulin levels in patients and rats with subarachnoid hemorrhage: a potential role in inhibiting inflammation by suppressing neutrophil recruitment. <i>Journal of Neuroinflammation</i> , 2015, 12, 200. | 7.2 | 30 |
| 5 | Constriction and dysfunction of pial arterioles after regional hemorrhage in the subarachnoid space. <i>Brain Research</i> , 2015, 1601, 85-91. | 2.2 | 3 |
| 6 | Baincalein alleviates early brain injury after experimental subarachnoid hemorrhage in rats: Possible involvement of TLR4/NF- κ B-mediated inflammatory pathway. <i>Brain Research</i> , 2015, 1594, 245-255. | 2.2 | 46 |
| 7 | Activation of the Protein Kinase B (Akt) Reduces Nur77-induced Apoptosis During Early Brain Injury after Experimental Subarachnoid Hemorrhage in Rat. <i>Annals of Clinical and Laboratory Science</i> , 2015, 45, 615-22. | 0.2 | 7 |
| 8 | Astaxanthin Alleviates Early Brain Injury Following Subarachnoid Hemorrhage in Rats: Possible Involvement of Akt/Bad Signaling. <i>Marine Drugs</i> , 2014, 12, 4291-4310. | 4.6 | 68 |
| 9 | Ghrelin alleviates early brain injury after subarachnoid hemorrhage via the PI3K/Akt signaling pathway. <i>Brain Research</i> , 2014, 1587, 15-22. | 2.2 | 29 |
| 10 | Cyclosporin A ameliorates early brain injury after subarachnoid hemorrhage through inhibition of a Nur77 dependent apoptosis pathway. <i>Brain Research</i> , 2014, 1556, 67-76. | 2.2 | 17 |
| 11 | Resveratrol prevents neuronal apoptosis in an early brain injury model. <i>Journal of Surgical Research</i> , 2014, 189, 159-165. | 1.6 | 44 |
| 12 | Astaxanthin offers neuroprotection and reduces neuroinflammation in experimental subarachnoid hemorrhage. <i>Journal of Surgical Research</i> , 2014, 192, 206-213. | 1.6 | 103 |