

Ewa M Roos

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/203178/publications.pdf>

Version: 2024-02-01

281
papers

31,828
citations

7561

77
h-index

4427

172
g-index

287
all docs

287
docs citations

287
times ranked

16925
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Knee Injury and Osteoarthritis Outcome Score (KOOS) – Development of a Self-Administered Outcome Measure. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 1998, 28, 88-96. | 1.7 | 2,936 |
| 2 | OARSI guidelines for the non-surgical management of knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2014, 22, 363-388. | 0.6 | 2,298 |
| 3 | The Long-term Consequence of Anterior Cruciate Ligament and Meniscus Injuries. <i>American Journal of Sports Medicine</i> , 2007, 35, 1756-1769. | 1.9 | 1,871 |
| 4 | The Knee injury and Osteoarthritis Outcome Score (KOOS): from joint injury to osteoarthritis. <i>Health and Quality of Life Outcomes</i> , 2003, 1, 64. | 1.0 | 1,594 |
| 5 | Measures of knee function: International Knee Documentation Committee (IKDC) Subjective Knee Evaluation Form, Knee Injury and Osteoarthritis Outcome Score (KOOS), Knee Injury and Osteoarthritis Outcome Score Physical Function Short Form (KOOS-PS), Knee Outcome Survey Activities of Daily Living Scale (KOS-ADL), Lysholm Knee Scoring Scale, Oxford Knee Score (OKS), Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), Activity Rating Scale (ARS), and Tegner Activity Score (TAS). <i>Arthritis Care and Research</i> , 2011, 63, S208-28. | 1.5 | 897 |
| 6 | Hip disability and osteoarthritis outcome score (HOOS) – validity and responsiveness in total hip replacement. <i>BMC Musculoskeletal Disorders</i> , 2003, 4, 10. | 0.8 | 772 |
| 7 | Knee injury and Osteoarthritis Outcome Score (KOOS) - validation and comparison to the WOMAC in total knee replacement. <i>Health and Quality of Life Outcomes</i> , 2003, 1, 17. | 1.0 | 769 |
| 8 | High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes. <i>Annals of the Rheumatic Diseases</i> , 2004, 63, 269-273. | 0.5 | 692 |
| 9 | A Randomized Trial of Treatment for Acute Anterior Cruciate Ligament Tears. <i>New England Journal of Medicine</i> , 2010, 363, 331-342. | 13.9 | 686 |
| 10 | Knee osteoarthritis after meniscectomy: Prevalence of radiographic changes after twenty-one years, compared with matched controls. <i>Arthritis and Rheumatism</i> , 1998, 41, 687-693. | 6.7 | 631 |
| 11 | Validation of the Foot and Ankle Outcome Score for Ankle Ligament Reconstruction. <i>Foot and Ankle International</i> , 2001, 22, 788-794. | 1.1 | 612 |
| 12 | Non-contact ACL injuries in female athletes: an International Olympic Committee current concepts statement. <i>British Journal of Sports Medicine</i> , 2008, 42, 394-412. | 3.1 | 582 |
| 13 | OARSI recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2013, 21, 1042-1052. | 0.6 | 545 |
| 14 | A Randomized, Controlled Trial of Total Knee Replacement. <i>New England Journal of Medicine</i> , 2015, 373, 1597-1606. | 13.9 | 498 |
| 15 | Knee injury and Osteoarthritis Outcome Score (KOOS) – validation of a Swedish version. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 1998, 8, 439-448. | 1.3 | 485 |
| 16 | Knee Injury and Osteoarthritis Outcome Score (KOOS): systematic review and meta-analysis of measurement properties. <i>Osteoarthritis and Cartilage</i> , 2016, 24, 1317-1329. | 0.6 | 412 |
| 17 | Impact of Exercise Type and Dose on Pain and Disability in Knee Osteoarthritis: A Systematic Review and Meta-Regression Analysis of Randomized Controlled Trials. <i>Arthritis and Rheumatology</i> , 2014, 66, 622-636. | 2.9 | 398 |
| 18 | The Copenhagen Hip and Groin Outcome Score (HAGOS): development and validation according to the COSMIN checklist. <i>British Journal of Sports Medicine</i> , 2011, 45, 478-491. | 3.1 | 396 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Positive effects of moderate exercise on glycosaminoglycan content in knee cartilage: A four-month, randomized, controlled trial in patients at risk of osteoarthritis. <i>Arthritis and Rheumatism</i> , 2005, 52, 3507-3514. | 6.7 | 392 |
| 20 | Psychometric Properties of Patient-Reported Outcome Measures for Hip Arthroscopic Surgery. <i>American Journal of Sports Medicine</i> , 2013, 41, 2065-2073. | 1.9 | 389 |
| 21 | Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. <i>BMJ, The</i> , 2013, 346, f232-f232. | 3.0 | 369 |
| 22 | Joint injury causes knee osteoarthritis in young adults. <i>Current Opinion in Rheumatology</i> , 2005, 17, 195-200. | 2.0 | 353 |
| 23 | Strategies for the prevention of knee osteoarthritis. <i>Nature Reviews Rheumatology</i> , 2016, 12, 92-101. | 3.5 | 340 |
| 24 | Knee arthroplasty: are patients' expectations fulfilled?. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2009, 80, 55-61. | 1.2 | 281 |
| 25 | Clinical improvement after 6 weeks of eccentric exercise in patients with mid-portion Achilles tendinopathy - a randomized trial with 1-year follow-up. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2004, 14, 286-295. | 1.3 | 264 |
| 26 | Knee complaints vary with age and gender in the adult population. Population-based reference data for the Knee injury and Osteoarthritis Outcome Score (KOOS). <i>BMC Musculoskeletal Disorders</i> , 2006, 7, 38. | 0.8 | 262 |
| 27 | Good Life with osteoArthritis in Denmark (GLA:D [®]): evidence-based education and supervised neuromuscular exercise delivered by certified physiotherapists nationwide. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 72. | 0.8 | 249 |
| 28 | Patient-relevant outcomes fourteen years after meniscectomy: influence of type of meniscal tear and size of resection. <i>Rheumatology</i> , 2001, 40, 631-639. | 0.9 | 235 |
| 29 | Predictors of patient relevant outcome after total hip replacement for osteoarthritis: a prospective study. <i>Annals of the Rheumatic Diseases</i> , 2003, 62, 923-930. | 0.5 | 225 |
| 30 | The development of a short measure of physical function for knee OA KOOS-Physical Function Shortform (KOOS-PS) – an OARSI/OMERACT initiative. <i>Osteoarthritis and Cartilage</i> , 2008, 16, 542-550. | 0.6 | 219 |
| 31 | Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle aged patients: randomised controlled trial with two year follow-up. <i>BMJ, The</i> , 2016, 354, i3740. | 3.0 | 215 |
| 32 | Measurement properties of performance-based measures to assess physical function in hip and knee osteoarthritis: a systematic review. <i>Osteoarthritis and Cartilage</i> , 2012, 20, 1548-1562. | 0.6 | 209 |
| 33 | Long-term outcome of meniscectomy: symptoms, function, and performance tests in patients with or without radiographic osteoarthritis compared to matched controls. <i>Osteoarthritis and Cartilage</i> , 2001, 9, 316-324. | 0.6 | 192 |
| 34 | Sex Differences in Patient-Reported Outcomes After Anterior Cruciate Ligament Reconstruction. <i>American Journal of Sports Medicine</i> , 2010, 38, 1334-1342. | 1.9 | 189 |
| 35 | Isokinetic knee extensor strength and functional performance in healthy female soccer players. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 1998, 8, 257-264. | 1.3 | 167 |
| 36 | The role of pain and functional impairment in the decision to recommend total joint replacement in hip and knee osteoarthritis: an international cross-sectional study of 1909 patients. Report of the OARSI-OMERACT Task Force on total joint replacement. <i>Osteoarthritis and Cartilage</i> , 2011, 19, 147-154. | 0.6 | 162 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Validity, reliability and responsiveness of patient-reported outcome questionnaires when assessing hip and groin disability: a systematic review. <i>British Journal of Sports Medicine</i> , 2010, 44, 1186-1196. | 3.1 | 158 |
| 38 | Knee osteoarthritis risk is increased 4-6 fold after knee injury – a systematic review and meta-analysis. <i>British Journal of Sports Medicine</i> , 2019, 53, 1454-1463. | 3.1 | 158 |
| 39 | Muscle weakness, afferent sensory dysfunction and exercise in knee osteoarthritis. <i>Nature Reviews Rheumatology</i> , 2011, 7, 57-63. | 3.5 | 157 |
| 40 | Is Knee Pain During Adolescence a Self-limiting Condition?. <i>American Journal of Sports Medicine</i> , 2016, 44, 1165-1171. | 1.9 | 157 |
| 41 | Cross-cultural adaptation and validation of the French version of the Knee injury and Osteoarthritis Outcome Score (KOOS) in knee osteoarthritis patients. <i>Osteoarthritis and Cartilage</i> , 2008, 16, 423-428. | 0.6 | 156 |
| 42 | The development of a short measure of physical function for hip OA HOOS-Physical Function Shortform (HOOS-PS): an OARSI/OMERACT initiative. <i>Osteoarthritis and Cartilage</i> , 2008, 16, 551-559. | 0.6 | 156 |
| 43 | Reduced functional performance in the lower extremity predicted radiographic knee osteoarthritis five years later. <i>Annals of the Rheumatic Diseases</i> , 2004, 63, 402-407. | 0.5 | 147 |
| 44 | Validation of the Dutch version of the Hip disability and Osteoarthritis Outcome Score. <i>Osteoarthritis and Cartilage</i> , 2007, 15, 104-109. | 0.6 | 147 |
| 45 | Feasibility of neuromuscular training in patients with severe hip or knee OA: The individualized goal-based NEMEX-TJR training program. <i>BMC Musculoskeletal Disorders</i> , 2010, 11, 126. | 0.8 | 144 |
| 46 | Validity and inter-rater reliability of medio-lateral knee motion observed during a single-limb mini squat. <i>BMC Musculoskeletal Disorders</i> , 2010, 11, 265. | 0.8 | 143 |
| 47 | Neuromuscular Versus Quadriceps Strengthening Exercise in Patients With Medial Knee Osteoarthritis and Varus Malalignment: A Randomized Controlled Trial. <i>Arthritis and Rheumatology</i> , 2014, 66, 950-959. | 2.9 | 138 |
| 48 | WOMAC Osteoarthritis Index – additional dimensions for use in subjects with post-traumatic osteoarthritis of the knee. <i>Osteoarthritis and Cartilage</i> , 1999, 7, 216-221. | 0.6 | 137 |
| 49 | Magnitude and meaningfulness of change in SF-36 scores in four types of orthopedic surgery. <i>Health and Quality of Life Outcomes</i> , 2008, 6, 55. | 1.0 | 135 |
| 50 | Comparative, validity and responsiveness of the HOOS-PS and KOOS-PS to the WOMAC physical function subscale in total joint replacement for Osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2009, 17, 843-847. | 0.6 | 135 |
| 51 | Minimal clinically important improvement (MCII) and patient-acceptable symptom state (PASS) in total hip arthroplasty (THA) patients 1 year postoperatively. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2014, 85, 39-48. | 1.2 | 134 |
| 52 | Muscle strength and functional performance in patients with anterior cruciate ligament injury treated with training and surgical reconstruction or training only: A two to five-year followup. <i>Arthritis and Rheumatism</i> , 2008, 59, 1773-1779. | 6.7 | 133 |
| 53 | The acutely ACL injured knee assessed by MRI: changes in joint fluid, bone marrow lesions, and cartilage during the first year. <i>Osteoarthritis and Cartilage</i> , 2009, 17, 161-167. | 0.6 | 133 |
| 54 | A 5 year prospective study of patient-relevant outcomes after total knee replacement. <i>Osteoarthritis and Cartilage</i> , 2009, 17, 601-606. | 0.6 | 133 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Patient-Reported Outcome (PRO) questionnaires for young to middle-aged adults with hip and groin disability: a systematic review of the clinimetric evidence. <i>British Journal of Sports Medicine</i> , 2015, 49, 812-812. | 3.1 | 133 |
| 56 | Osteoarthritis: Models for appropriate care across the disease continuum. <i>Best Practice and Research in Clinical Rheumatology</i> , 2016, 30, 503-535. | 1.4 | 123 |
| 57 | Patient-Reported Outcomes for Total Hip and Knee Arthroplasty. <i>Clinics in Geriatric Medicine</i> , 2012, 28, 367-394. | 1.0 | 121 |
| 58 | Effectiveness and practice variation of rehabilitation after joint replacement. <i>Current Opinion in Rheumatology</i> , 2003, 15, 160-162. | 2.0 | 114 |
| 59 | ICRS Recommendation Document. <i>Cartilage</i> , 2011, 2, 122-136. | 1.4 | 114 |
| 60 | Clinical update: treating osteoarthritis. <i>Lancet, The</i> , 2007, 370, 2082-2084. | 6.3 | 113 |
| 61 | Exercise during school hours when added to patient education improves outcome for 2-3 years in adolescent patellofemoral pain: a cluster randomised trial. <i>British Journal of Sports Medicine</i> , 2015, 49, 406-412. | 3.1 | 113 |
| 62 | High prevalence of daily and multi-site pain "a cross-sectional population-based study among 3000 Danish adolescents. <i>BMC Pediatrics</i> , 2013, 13, 191. | 0.7 | 106 |
| 63 | Total knee replacement and non-surgical treatment of knee osteoarthritis: 2-year outcome from two parallel randomized controlled trials. <i>Osteoarthritis and Cartilage</i> , 2018, 26, 1170-1180. | 0.6 | 106 |
| 64 | Knee extension and flexion muscle power after anterior cruciate ligament reconstruction with patellar tendon graft or hamstring tendons graft: a cross-sectional comparison 3 years post surgery. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2009, 17, 162-169. | 2.3 | 105 |
| 65 | Foot Orthoses for the Treatment of Plantar Fasciitis. <i>Foot and Ankle International</i> , 2006, 27, 606-611. | 1.1 | 104 |
| 66 | The acutely ACL injured knee assessed by MRI: are large volume traumatic bone marrow lesions a sign of severe compression injury?. <i>Osteoarthritis and Cartilage</i> , 2008, 16, 829-836. | 0.6 | 98 |
| 67 | Serum levels of Cartilage Oligomeric Matrix Protein (COMP) increase temporarily after physical exercise in patients with knee osteoarthritis. <i>BMC Musculoskeletal Disorders</i> , 2006, 7, 98. | 0.8 | 94 |
| 68 | Muscle strength, functional performance, and self-reported outcomes four years after arthroscopic partial meniscectomy in middle-aged patients. <i>Arthritis and Rheumatism</i> , 2006, 55, 946-952. | 6.7 | 93 |
| 69 | Lower Mechanical Pressure Pain Thresholds in Female Adolescents With Patellofemoral Pain Syndrome. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2013, 43, 414-421. | 1.7 | 92 |
| 70 | Substantial disability 3 months after arthroscopic partial meniscectomy: A prospective study of patient-relevant outcomes. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2000, 16, 619-626. | 1.3 | 90 |
| 71 | Establishing outcome measures in early knee osteoarthritis. <i>Nature Reviews Rheumatology</i> , 2019, 15, 438-448. | 3.5 | 88 |
| 72 | Proportion of Patients Reporting Acceptable Symptoms or Treatment Failure and Their Associated KOOS Values at 6 to 24 Months After Anterior Cruciate Ligament Reconstruction. <i>American Journal of Sports Medicine</i> , 2015, 43, 1902-1907. | 1.9 | 87 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | The effect of eight weeks of exercise on knee adduction moment in early knee osteoarthritis – a pilot study. <i>Osteoarthritis and Cartilage</i> , 2007, 15, 1163-1170. | 0.6 | 86 |
| 74 | Neuromuscular Exercise as Treatment of Degenerative Knee Disease. <i>Exercise and Sport Sciences Reviews</i> , 2015, 43, 14-22. | 1.6 | 84 |
| 75 | Development of the Knee Injury and Osteoarthritis Outcome Score for Children (KOOS-Child). <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2012, 83, 666-673. | 1.2 | 83 |
| 76 | Six-week high-intensity exercise program for middle-aged patients with knee osteoarthritis: a randomized controlled trial [ISRCTN20244858]. <i>BMC Musculoskeletal Disorders</i> , 2005, 6, 27. | 0.8 | 82 |
| 77 | The OMERACT-OARSI Core Domain Set for Measurement in Clinical Trials of Hip and/or Knee Osteoarthritis. <i>Journal of Rheumatology</i> , 2019, 46, 981-989. | 1.0 | 82 |
| 78 | Cross-cultural adaptation and validation of Singapore English and Chinese versions of the Knee injury and Osteoarthritis Outcome Score (KOOS) in Asians with knee osteoarthritis in Singapore. <i>Osteoarthritis and Cartilage</i> , 2006, 14, 1098-1103. | 0.6 | 80 |
| 79 | The patellofemoral pain and osteoarthritis subscale of the KOOS (KOOS-PF): development and validation using the COSMIN checklist. <i>British Journal of Sports Medicine</i> , 2018, 52, 1130-1136. | 3.1 | 80 |
| 80 | Postoperative effects of neuromuscular exercise prior to hip or knee arthroplasty: a randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 1130-1137. | 0.5 | 77 |
| 81 | Comparative responsiveness of measures of pain and function after total hip replacement. <i>Arthritis and Rheumatism</i> , 2001, 45, 258-262. | 6.7 | 76 |
| 82 | OMERACT/OARSI initiative to define states of severity and indication for joint replacement in hip and knee osteoarthritis. <i>Journal of Rheumatology</i> , 2007, 34, 1432-5. | 1.0 | 74 |
| 83 | Cross-cultural adaptation and validation of the French version of the Hip disability and Osteoarthritis Outcome Score (HOOS) in hip osteoarthritis patients. <i>Osteoarthritis and Cartilage</i> , 2010, 18, 522-529. | 0.6 | 73 |
| 84 | Immediate Efficacy of Neuromuscular Exercise in Patients with Severe Osteoarthritis of the Hip or Knee: A Secondary Analysis from a Randomized Controlled Trial. <i>Journal of Rheumatology</i> , 2014, 41, 1385-1394. | 1.0 | 72 |
| 85 | Meaningful Change Scores in the Knee Injury and Osteoarthritis Outcome Score in Patients Undergoing Anterior Cruciate Ligament Reconstruction. <i>American Journal of Sports Medicine</i> , 2018, 46, 1120-1128. | 1.9 | 72 |
| 86 | Validity and reliability of functional performance tests in meniscectomized patients with or without knee osteoarthritis. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2006, 17, 061120070736056-??? | 1.3 | 71 |
| 87 | The efficacy of 12 weeks non-surgical treatment for patients not eligible for total knee replacement: a randomized controlled trial with 1-year follow-up. <i>Osteoarthritis and Cartilage</i> , 2015, 23, 1465-1475. | 0.6 | 70 |
| 88 | Osteoarthritis 2012 year in review: rehabilitation and outcomes. <i>Osteoarthritis and Cartilage</i> , 2012, 20, 1477-1483. | 0.6 | 68 |
| 89 | Impact of exercise on articular cartilage in people at risk of, or with established, knee osteoarthritis: a systematic review of randomised controlled trials. <i>British Journal of Sports Medicine</i> , 2019, 53, 940-947. | 3.1 | 67 |
| 90 | Effects of neuromuscular training (NEMEX-TJR) on patient-reported outcomes and physical function in severe primary hip or knee osteoarthritis: a controlled before-and-after study. <i>BMC Musculoskeletal Disorders</i> , 2013, 14, 232. | 0.8 | 66 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Delaying ACL reconstruction and treating with exercise therapy alone may alter prognostic factors for 5-year outcome: an exploratory analysis of the KANON trial. <i>British Journal of Sports Medicine</i> , 2017, 51, 1622-1629. | 3.1 | 64 |
| 92 | A 12-item short form of the Knee injury and Osteoarthritis Outcome Score (KOOS-12): tests of reliability, validity and responsiveness. <i>Osteoarthritis and Cartilage</i> , 2019, 27, 762-770. | 0.6 | 64 |
| 93 | Variation in age and physical status prior to total knee and hip replacement surgery: A comparison of centers in Australia and Europe. <i>Arthritis and Rheumatism</i> , 2009, 61, 166-173. | 6.7 | 61 |
| 94 | Patterns of musculoskeletal pain in the population: A latent class analysis using a nationally representative interviewer-based survey of 4817 Danes. <i>European Journal of Pain</i> , 2013, 17, 452-460. | 1.4 | 61 |
| 95 | Minimal important change values for the Oxford Knee Score and the Forgotten Joint Score at 1 year after total knee replacement. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018, 89, 541-547. | 1.2 | 59 |
| 96 | The challenge of recruiting patients with anterior cruciate ligament injury of the knee into a randomized clinical trial comparing surgical and non-surgical treatment. <i>Contemporary Clinical Trials</i> , 2007, 28, 295-302. | 0.8 | 58 |
| 97 | Increased pain sensitivity but normal function of exercise induced analgesia in hip and knee osteoarthritis – treatment effects of neuromuscular exercise and total joint replacement. <i>Osteoarthritis and Cartilage</i> , 2013, 21, 1299-1307. | 0.6 | 58 |
| 98 | Lower extremity performance following ACL rehabilitation in the KANON-trial: impact of reconstruction and predictive value at 2 and 5 years. <i>British Journal of Sports Medicine</i> , 2013, 47, 980-985. | 3.1 | 58 |
| 99 | Hip and Knee Strength Is Not Affected in 12-16 Year Old Adolescents with Patellofemoral Pain - A Cross-Sectional Population-Based Study. <i>PLoS ONE</i> , 2013, 8, e79153. | 1.1 | 58 |
| 100 | Measurement properties of performance-based outcome measures to assess physical function in young and middle-aged people known to be at high risk of hip and/or knee osteoarthritis: a systematic review. <i>Osteoarthritis and Cartilage</i> , 2014, 22, 26-39. | 0.6 | 58 |
| 101 | It is time to move beyond “body region silos” to manage musculoskeletal pain: five actions to change clinical practice. <i>British Journal of Sports Medicine</i> , 2020, 54, 438-439. | 3.1 | 58 |
| 102 | The effect of patient characteristics on variability in pain and function over two years in early knee osteoarthritis. <i>Health and Quality of Life Outcomes</i> , 2005, 3, 59. | 1.0 | 56 |
| 103 | How do middle-aged patients conceive exercise as a form of treatment for knee osteoarthritis?. <i>Disability and Rehabilitation</i> , 2006, 28, 51-59. | 0.9 | 53 |
| 104 | Psychometric properties of the Knee injury and Osteoarthritis Outcome Score for Children (KOOS-Child) in children with knee disorders. <i>British Journal of Sports Medicine</i> , 2014, 48, 1437-1446. | 3.1 | 52 |
| 105 | Effect of Exercise Therapy Compared with Arthroscopic Surgery on Knee Muscle Strength and Functional Performance in Middle-Aged Patients with Degenerative Meniscus Tears. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2015, 94, 460-473. | 0.7 | 51 |
| 106 | Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. <i>British Journal of Sports Medicine</i> , 2015, 49, 700-700. | 3.1 | 51 |
| 107 | Pain trajectory and exercise-induced pain flares during 8 weeks of neuromuscular exercise in individuals with knee and hip pain. <i>Osteoarthritis and Cartilage</i> , 2016, 24, 589-592. | 0.6 | 51 |
| 108 | Change in self-reported outcomes and objective physical function over 7 years in middle-aged subjects with or at high risk of knee osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2008, 67, 505-510. | 0.5 | 50 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Care-seeking behaviour of adolescents with knee pain: a population-based study among 504 adolescents. <i>BMC Musculoskeletal Disorders</i> , 2013, 14, 225. | 0.8 | 50 |
| 110 | Comparison of neuromuscular and quadriceps strengthening exercise in the treatment of varus malaligned knees with medial knee osteoarthritis: a randomised controlled trial protocol. <i>BMC Musculoskeletal Disorders</i> , 2011, 12, 276. | 0.8 | 47 |
| 111 | A 12-Week Exercise Therapy Program in Middle-Aged Patients With Degenerative Meniscus Tears: A Case Series With 1-Year Follow-up. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2012, 42, 919-931. | 1.7 | 47 |
| 112 | A 12-item short form of the Hip disability and Osteoarthritis Outcome Score (HOOS-12): tests of reliability, validity and responsiveness. <i>Osteoarthritis and Cartilage</i> , 2019, 27, 754-761. | 0.6 | 47 |
| 113 | A pragmatic approach to prevent post-traumatic osteoarthritis after sport or exercise-related joint injury. <i>Best Practice and Research in Clinical Rheumatology</i> , 2019, 33, 158-171. | 1.4 | 46 |
| 114 | Choosing surgery: patients' preferences within a trial of treatments for anterior cruciate ligament injury. A qualitative study. <i>BMC Musculoskeletal Disorders</i> , 2009, 10, 100. | 0.8 | 45 |
| 115 | Effectiveness and safety of strengthening, aerobic, and coordination exercises for patients with osteoarthritis. <i>Current Opinion in Rheumatology</i> , 2003, 15, 141-144. | 2.0 | 44 |
| 116 | Effects of functional exercise training on performance and muscle strength after meniscectomy: a randomized trial. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2009, 19, 156-165. | 1.3 | 43 |
| 117 | Effect of leisure time physical activity on severe knee or hip osteoarthritis leading to total joint replacement: a population-based prospective cohort study. <i>BMC Musculoskeletal Disorders</i> , 2012, 13, 73. | 0.8 | 43 |
| 118 | Neuromuscular Activity and Knee Kinematics in Adolescents with Patellofemoral Pain. <i>Medicine and Science in Sports and Exercise</i> , 2013, 45, 1730-1739. | 0.2 | 43 |
| 119 | Arthroscopic partial meniscectomy in middle-aged patients with mild or no knee osteoarthritis: a protocol for a double-blind, randomized sham-controlled multi-centre trial. <i>BMC Musculoskeletal Disorders</i> , 2013, 14, 71. | 0.8 | 42 |
| 120 | Five-year prognosis and impact of adolescent knee pain: a prospective population-based cohort study of 504 adolescents in Denmark. <i>BMJ Open</i> , 2019, 9, e024113. | 0.8 | 42 |
| 121 | It is good to feel better, but better to feel good: whether a patient finds treatment "successful" or not depends on the questions researchers ask. <i>British Journal of Sports Medicine</i> , 2019, 53, 1474-1478. | 3.1 | 42 |
| 122 | Validation of the Rheumatoid and Arthritis Outcome Score (RAOS) for the lower extremity. <i>Health and Quality of Life Outcomes</i> , 2003, 1, 55. | 1.0 | 41 |
| 123 | Examining the Minimal Important Difference of Patient-reported Outcome Measures for Individuals with Knee Osteoarthritis: A Model Using the Knee Injury and Osteoarthritis Outcome Score. <i>Journal of Rheumatology</i> , 2016, 43, 395-404. | 1.0 | 41 |
| 124 | Cross-cultural translation and measurement properties of the Polish version of the Knee injury and Osteoarthritis Outcome Score (KOOS) following anterior cruciate ligament reconstruction. <i>Health and Quality of Life Outcomes</i> , 2013, 11, 107. | 1.0 | 40 |
| 125 | GLA:D to have a high-value option for patients with knee and hip arthritis across four continents: Good Life with osteoArthritis from Denmark. <i>British Journal of Sports Medicine</i> , 2018, 52, 1544-1545. | 3.1 | 40 |
| 126 | GLA:D® Back group-based patient education integrated with exercises to support self-management of back pain - development, theories and scientific evidence. <i>BMC Musculoskeletal Disorders</i> , 2018, 19, 418. | 0.8 | 40 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | A 2-year prospective study of patient-relevant outcomes in patients operated on for knee osteoarthritis with tibial osteotomy. <i>BMC Musculoskeletal Disorders</i> , 2005, 6, 18. | 0.8 | 39 |
| 128 | Item selection for 12-item short forms of the Knee injury and Osteoarthritis Outcome Score (KOOS-12) and Hip disability and Osteoarthritis Outcome Score (HOOS-12). <i>Osteoarthritis and Cartilage</i> , 2019, 27, 746-753. | 0.6 | 39 |
| 129 | Over-optimistic patient expectations of recovery and leisure activities after arthroscopic meniscus surgery. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2016, 87, 615-621. | 1.2 | 37 |
| 130 | Supervised neuromuscular exercise prior to hip and knee replacement: 12-month clinical effect and cost-utility analysis alongside a randomised controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 5. | 0.8 | 37 |
| 131 | Three steps to changing the narrative about knee osteoarthritis care: a call to action. <i>British Journal of Sports Medicine</i> , 2020, 54, 256-258. | 3.1 | 37 |
| 132 | Relationships between postural orientation and self reported function, hop performance and muscle power in subjects with anterior cruciate ligament injury. <i>BMC Musculoskeletal Disorders</i> , 2010, 11, 143. | 0.8 | 35 |
| 133 | Influence of Biomechanical Characteristics on Pain and Function Outcomes From Exercise in Medial Knee Osteoarthritis and Varus Malalignment: Exploratory Analyses From a Randomized Controlled Trial. <i>Arthritis Care and Research</i> , 2015, 67, 1281-1288. | 1.5 | 35 |
| 134 | Better outcome from arthroscopic partial meniscectomy than skin incisions only? A sham-controlled randomised trial in patients aged 35-55 years with knee pain and an MRI-verified meniscal tear. <i>BMJ Open</i> , 2018, 8, e019461. | 0.8 | 35 |
| 135 | Feasibility of 4 patient-reported outcome measures in a registry setting. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2012, 83, 321-327. | 1.2 | 34 |
| 136 | Agreement and Reliability of Functional Performance and Muscle Power in Patients with Advanced Osteoarthritis of the Hip or Knee. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2012, 91, 401-410. | 0.7 | 34 |
| 137 | Early intervention for adolescents with Patellofemoral Pain Syndrome - a pragmatic cluster randomised controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2012, 13, 9. | 0.8 | 34 |
| 138 | Patient education with or without manual therapy compared to a control group in patients with osteoarthritis of the hip. A proof-of-principle three-arm parallel group randomized clinical trial. <i>Osteoarthritis and Cartilage</i> , 2013, 21, 1494-1503. | 0.6 | 34 |
| 139 | Effect of preoperative neuromuscular training (NEMEX-TJR) on functional outcome after total knee replacement: an assessor-blinded randomized controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2015, 16, 101. | 0.8 | 34 |
| 140 | The impact of physical activity level on the short- and long-term pain relief from supervised exercise therapy and education: a study of 12,796 Danish patients with knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2018, 26, 1474-1478. | 0.6 | 34 |
| 141 | Impact of a daily exercise dose on knee joint cartilage - a systematic review and meta-analysis of randomized controlled trials in healthy animals. <i>Osteoarthritis and Cartilage</i> , 2017, 25, 1223-1237. | 0.6 | 33 |
| 142 | Impact of Exercise Therapy on Molecular Biomarkers Related to Cartilage and Inflammation in Individuals at Risk of, or With Established, Knee Osteoarthritis: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. <i>Arthritis Care and Research</i> , 2019, 71, 1504-1515. | 1.5 | 33 |
| 143 | Psychometric properties of the French translation of the reduced KOOS and HOOS (KOOS-PS and HOOS-PS). <i>Osteoarthritis and Cartilage</i> , 2017, 25, 1238-1247. | 0.6 | 32 |
| 144 | Self-reported musculoskeletal pain predicts long-term increase in general health care use: A population-based cohort study with 20-year follow-up. <i>Scandinavian Journal of Public Health</i> , 2014, 42, 698-704. | 1.2 | 32 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | The evidence base for orthopaedics and sports medicine. <i>BMJ</i> , The, 2015, 350, g7835-g7835. | 3.0 | 32 |
| 146 | Young females with long-standing patellofemoral pain display impaired conditioned pain modulation, increased temporal summation of pain, and widespread hyperalgesia. <i>Pain</i> , 2018, 159, 2530-2537. | 2.0 | 32 |
| 147 | Comparison of three sets of clinical classification criteria for knee osteoarthritis: a cross-sectional study of 13,459 patients treated in primary care. <i>Osteoarthritis and Cartilage</i> , 2020, 28, 167-172. | 0.6 | 32 |
| 148 | Psychometric properties of the Neck OutCome Score, Neck Disability Index, and Short Form-36 were evaluated in patients with neck pain. <i>Journal of Clinical Epidemiology</i> , 2016, 79, 31-40. | 2.4 | 31 |
| 149 | Half of 12-15-year-olds with knee pain still have pain after one year. <i>Danish Medical Journal</i> , 2013, 60, A4725. | 0.5 | 30 |
| 150 | Knee kinematics and kinetics in former soccer players with a 16-year-old ACL injury – the effects of twelve weeks of knee-specific training. <i>BMC Musculoskeletal Disorders</i> , 2007, 8, 35. | 0.8 | 29 |
| 151 | Criteria used when deciding on eligibility for total knee arthroplasty – Between thinking and doing. <i>Knee</i> , 2016, 23, 300-305. | 0.8 | 29 |
| 152 | Thigh muscle strength, functional capacity, and self-reported function in patients at high risk of knee osteoarthritis compared with controls. <i>Arthritis Care and Research</i> , 2010, 62, 1244-1251. | 1.5 | 28 |
| 153 | Reliability and reference values of two clinical measurements of dynamic and static knee position in healthy children. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2011, 19, 2060-2066. | 2.3 | 28 |
| 154 | Clinical Outcomes Assessment in Clinical Trials to Assess Treatment of Femoroacetabular Impingement: Use of Patient-reported Outcome Measures. <i>Journal of the American Academy of Orthopaedic Surgeons</i> , The, 2013, 21, S39-S46. | 1.1 | 28 |
| 155 | Knee function and knee muscle strength in middle-aged patients with degenerative meniscal tears eligible for arthroscopic partial meniscectomy. <i>British Journal of Sports Medicine</i> , 2014, 48, 784-788. | 3.1 | 28 |
| 156 | Muscle function is associated with future patient-reported outcomes in young adults with ACL injury. <i>BMJ Open Sport and Exercise Medicine</i> , 2016, 2, e000154. | 1.4 | 28 |
| 157 | Patients report inferior quality of care for knee osteoarthritis prior to assessment for knee replacement surgery – a cross-sectional study of 517 patients in Denmark. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2020, 91, 82-87. | 1.2 | 27 |
| 158 | Program evaluation of GLA:D® Australia: Physiotherapist training outcomes and effectiveness of implementation for people with knee osteoarthritis. <i>Osteoarthritis and Cartilage Open</i> , 2021, 3, 100175. | 0.9 | 27 |
| 159 | Is exercise therapy for femoroacetabular impingement in or out of FASHIoN? We need to talk about current best practice for the non-surgical management of FAI syndrome. <i>British Journal of Sports Medicine</i> , 2019, 53, 1204-1205. | 3.1 | 26 |
| 160 | Similar group mean scores, but large individual variations, in patient-relevant outcomes over 2 years in meniscectomized subjects with and without radiographic knee osteoarthritis. <i>Health and Quality of Life Outcomes</i> , 2004, 2, 38. | 1.0 | 25 |
| 161 | The challenge of recruiting patients into a placebo-controlled surgical trial. <i>Trials</i> , 2014, 15, 167. | 0.7 | 25 |
| 162 | Uptake of the OMERACT-OARSI Hip and Knee Osteoarthritis Core Outcome Set: Review of Randomized Controlled Trials from 1997 to 2017. <i>Journal of Rheumatology</i> , 2019, 46, 976-980. | 1.0 | 25 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 163 | Rehabilitation after anterior cruciate ligament and meniscal injuries: a best-evidence synthesis of systematic reviews for the OPTIKNEE consensus. <i>British Journal of Sports Medicine</i> , 2022, 56, 1445-1453. | 3.1 | 25 |
| 164 | Group education and exercise is feasible in knee and hip osteoarthritis. <i>Danish Medical Journal</i> , 2012, 59, A4554. | 0.5 | 24 |
| 165 | Efficacy of multimodal, systematic non-surgical treatment of knee osteoarthritis for patients not eligible for a total knee replacement: a study protocol of a randomised controlled trial. <i>BMJ Open</i> , 2012, 2, e002168. | 0.8 | 23 |
| 166 | Muscle strength and functional performance in patients at high risk of knee osteoarthritis: a follow-up study. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2012, 20, 1110-1117. | 2.3 | 23 |
| 167 | Middle-aged patients with an MRI-verified medial meniscal tear report symptoms commonly associated with knee osteoarthritis. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2017, 88, 664-669. | 1.2 | 23 |
| 168 | Factors associated with the orthopaedic surgeon's decision to recommend total joint replacement in hip and knee osteoarthritis: an international cross-sectional study of 1905 patients. <i>Osteoarthritis and Cartilage</i> , 2018, 26, 1311-1318. | 0.6 | 23 |
| 169 | The effect on knee-joint load of instruction in analgesic use compared with neuromuscular exercise in patients with knee osteoarthritis: study protocol for a randomized, single-blind, controlled trial (the EXERPHARMA trial). <i>Trials</i> , 2014, 15, 444. | 0.7 | 22 |
| 170 | OARSI Clinical Trials Recommendations: Design and conduct of clinical trials for primary prevention of osteoarthritis by joint injury prevention in sport and recreation. <i>Osteoarthritis and Cartilage</i> , 2015, 23, 815-825. | 0.6 | 22 |
| 171 | A Randomized, Controlled Trial of Total Knee Replacement. <i>New England Journal of Medicine</i> , 2016, 374, 691-692. | 13.9 | 22 |
| 172 | A Pragmatic Approach to the Implementation of Osteoarthritis Guidelines Has Fewer Potential Barriers Than Recommended Implementation Frameworks. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2019, 49, 1-4. | 1.7 | 22 |
| 173 | The efficacy of non-surgical treatment on pain and sensitization in patients with knee osteoarthritis: a pre-defined ancillary analysis from a randomized controlled trial. <i>Osteoarthritis and Cartilage</i> , 2016, 24, 108-116. | 0.6 | 21 |
| 174 | Targeted prevention in primary care aimed at lifestyle-related diseases: a study protocol for a non-randomised pilot study. <i>BMC Family Practice</i> , 2018, 19, 124. | 2.9 | 21 |
| 175 | Development of osteoarthritis in patients with degenerative meniscal tears treated with exercise therapy or surgery: a randomized controlled trial. <i>Osteoarthritis and Cartilage</i> , 2020, 28, 897-906. | 0.6 | 21 |
| 176 | Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle aged patients: randomised controlled trial with two year follow-up. <i>British Journal of Sports Medicine</i> , 2016, 50, 1473-1480. | 3.1 | 20 |
| 177 | The Association Between Knee Confidence and Muscle Power, Hop Performance, and Postural Orientation in People With Anterior Cruciate Ligament Injury. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2016, 46, 477-482. | 1.7 | 20 |
| 178 | Routine knee arthroscopic surgery for the painful knee in middle-aged and old patientsâ€”time to abandon ship. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2016, 87, 2-4. | 1.2 | 20 |
| 179 | Patients use fewer analgesics following supervised exercise therapy and patient education: an observational study of 16â€”499 patients with knee or hip osteoarthritis. <i>British Journal of Sports Medicine</i> , 2021, 55, 670-675. | 3.1 | 19 |
| 180 | Total knee replacement plus physical and medical therapy or treatment with physical and medical therapy alone: a randomised controlled trial in patients with knee osteoarthritis (the MEDIC-study). <i>BMC Musculoskeletal Disorders</i> , 2012, 13, 67. | 0.8 | 18 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 181 | Republished research: Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. <i>British Journal of Sports Medicine</i> , 2013, 47, 373-373. | 3.1 | 17 |
| 182 | Effect of exercise therapy on neuromuscular activity and knee strength in female adolescents with patellofemoral pain—An ancillary analysis of a cluster randomized trial. <i>Clinical Biomechanics</i> , 2016, 34, 22-29. | 0.5 | 17 |
| 183 | Complex Tears, Extrusion, and Larger Excision Are Prognostic Factors for Worse Outcomes 1 and 2 Years After Arthroscopic Partial Meniscectomy for Degenerative Meniscal Tears: A Secondary Explorative Study of the Surgically Treated Group From the Odense-Oslo Meniscectomy Versus Exercise (OMEX) Trial. <i>American Journal of Sports Medicine</i> , 2019, 47, 2402-2411. | 1.9 | 17 |
| 184 | Association between thigh muscle strength four years after partial meniscectomy and radiographic features of osteoarthritis 11 years later. <i>BMC Musculoskeletal Disorders</i> , 2019, 20, 512. | 0.8 | 17 |
| 185 | Implementing a national first-line management program for moderate-severe knee osteoarthritis in Australia: A budget impact analysis focusing on knee replacement avoidance. <i>Osteoarthritis and Cartilage Open</i> , 2020, 2, 100070. | 0.9 | 17 |
| 186 | Reliability and validity of videotaped functional performance tests in ACL-injured subjects. <i>Physiotherapy Research International</i> , 2008, 13, 119-130. | 0.7 | 16 |
| 187 | Development of a patient-reported outcome: The Neck Outcome Score (NOOS) – Content and construct validity. <i>Journal of Rehabilitation Medicine</i> , 2015, 47, 844-853. | 0.8 | 16 |
| 188 | Bridging Disciplines as a pathway to Finding New Solutions for Osteoarthritis a collaborative program presented at the 2019 Orthopaedic Research Society and the Osteoarthritis Research Society International. <i>Osteoarthritis and Cartilage Open</i> , 2020, 2, 100026. | 0.9 | 16 |
| 189 | Associations between comorbidities and immediate and one-year outcomes following supervised exercise therapy and patient education – A cohort study of 24,513 individuals with knee or hip osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2021, 29, 39-49. | 0.6 | 16 |
| 190 | Validity and responsiveness of the Clubfoot Assessment Protocol (CAP). A methodological study. <i>BMC Musculoskeletal Disorders</i> , 2006, 7, 28. | 0.8 | 15 |
| 191 | Response to Letter to the Editor entitled “Comments on OARSI guidelines for the non-surgical management of knee osteoarthritis”. <i>Osteoarthritis and Cartilage</i> , 2014, 22, 890-891. | 0.6 | 15 |
| 192 | Self-reported Recovery is Associated With Improvement in Localized Hyperalgesia Among Adolescent Females With Patellofemoral Pain. <i>Clinical Journal of Pain</i> , 2016, 32, 428-434. | 0.8 | 15 |
| 193 | Association of osteoarthritis risk factors with knee and hip pain in a population-based sample of 29-59-year olds in Denmark: a cross-sectional analysis. <i>BMC Musculoskeletal Disorders</i> , 2018, 19, 300. | 0.8 | 15 |
| 194 | Predictors of long-term effect from education and exercise in patients with knee and hip pain. <i>Danish Medical Journal</i> , 2014, 61, A4867. | 0.5 | 15 |
| 195 | Association between knee alignment and knee pain in patients surgically treated for medial knee osteoarthritis by high tibial osteotomy. A one year follow-up study. <i>BMC Musculoskeletal Disorders</i> , 2009, 10, 154. | 0.8 | 14 |
| 196 | The iHOT-33: How Valid Is It?. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2012, 28, 1194-1195. | 1.3 | 14 |
| 197 | Neuromuscular Exercise post Partial Medial Meniscectomy. <i>Medicine and Science in Sports and Exercise</i> , 2015, 47, 1557-1566. | 0.2 | 14 |
| 198 | An 8-Week Neuromuscular Exercise Program for Patients With Mild to Moderate Knee Osteoarthritis: A Case Series Drawn From a Registered Clinical Trial. <i>Journal of Athletic Training</i> , 2017, 52, 592-605. | 0.9 | 14 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 199 | 3 steps to improve reporting and interpretation of patient-reported outcome scores in orthopedic studies. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018, 89, 1-2. | 1.2 | 14 |
| 200 | Cost-effectiveness of total knee replacement in addition to non-surgical treatment: a 2-year outcome from a randomised trial in secondary care in Denmark. <i>BMJ Open</i> , 2020, 10, e033495. | 0.8 | 14 |
| 201 | Vibratory perception threshold in young and middle-aged patients at high risk of knee osteoarthritis compared to controls. <i>Arthritis Care and Research</i> , 2012, 64, 144-148. | 1.5 | 13 |
| 202 | Knee Confidence as It Relates to Self-reported and Objective Correlates of Knee Osteoarthritis: A Cross-sectional Study of 220 Patients. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2015, 45, 765-771. | 1.7 | 13 |
| 203 | Non-surgical treatment of hip osteoarthritis. Hip school, with or without the addition of manual therapy, in comparison to a minimal control intervention: Protocol for a three-armed randomized clinical trial. <i>BMC Musculoskeletal Disorders</i> , 2011, 12, 88. | 0.8 | 12 |
| 204 | What fooled us in the knee may trip us up in the hip: lessons from arthroscopy. <i>British Journal of Sports Medicine</i> , 2014, 48, 1200-1201. | 3.1 | 12 |
| 205 | Surgical or non-surgical treatment of traumatic skeletal fractures in adults: systematic review and meta-analysis of benefits and harms. <i>Systematic Reviews</i> , 2020, 9, 179. | 2.5 | 12 |
| 206 | Individualized predictions of changes in knee pain, quality of life and walking speed following patient education and exercise therapy in patients with knee osteoarthritis â€” a prognostic model study. <i>Osteoarthritis and Cartilage</i> , 2020, 28, 1191-1201. | 0.6 | 12 |
| 207 | Cutoff Values to Interpret Short-term Treatment Outcomes After Arthroscopic Meniscal Surgery, Measured With the Knee Injury and Osteoarthritis Outcome Score. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2021, 51, 281-288. | 1.7 | 12 |
| 208 | Clinical Outcomes Assessment in Clinical Trials to Assess Treatment of Femoroacetabular Impingement: Use of Patientreported Outcome Measures. <i>Journal of the American Academy of Orthopaedic Surgeons</i> , The, 2013, 21, S39-S46. | 1.1 | 12 |
| 209 | Knee and hip osteoarthritis are more alike than different in baseline characteristics and outcomes: a longitudinal study of 32,599 patients participating in supervised education and exercise therapy. <i>Osteoarthritis and Cartilage</i> , 2022, 30, 681-688. | 0.6 | 12 |
| 210 | Neuromuscular Function during a Forward Lunge in Meniscectomized Patients. <i>Medicine and Science in Sports and Exercise</i> , 2012, 44, 1358-1365. | 0.2 | 11 |
| 211 | The effects of neuromuscular exercise on medial knee joint load post-arthroscopic partial medial meniscectomy: â€”SCOPEXâ€” a randomised control trial protocol. <i>BMC Musculoskeletal Disorders</i> , 2012, 13, 233. | 0.8 | 11 |
| 212 | Effect of pre-operative neuromuscular training on functional outcome after total knee replacement: a randomized-controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2013, 14, 157. | 0.8 | 11 |
| 213 | Crossâ€”Cultural and Construct Validity of the Animated Activity Questionnaire. <i>Arthritis Care and Research</i> , 2017, 69, 1349-1359. | 1.5 | 11 |
| 214 | Disease modification in OA â€” will we ever get there?. <i>Nature Reviews Rheumatology</i> , 2019, 15, 133-135. | 3.5 | 11 |
| 215 | Mapping EQ-5D-3L from the Knee Injury and Osteoarthritis Outcome Score (KOOS). <i>Quality of Life Research</i> , 2020, 29, 265-274. | 1.5 | 11 |
| 216 | Assessing baseline dependency of anchor-based minimal important change (MIC): donâ€™t stratify on the baseline score!. <i>Quality of Life Research</i> , 2021, 30, 2773-2782. | 1.5 | 11 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 217 | Prevalence of multimorbid degenerative lumbar spinal stenosis with knee or hip osteoarthritis: a systematic review and meta-analysis. <i>BMC Musculoskeletal Disorders</i> , 2022, 23, 177. | 0.8 | 11 |
| 218 | The evidence base for orthopaedics and sports medicine: scandalously poor in parts. <i>British Journal of Sports Medicine</i> , 2016, 50, 564-565. | 3.1 | 10 |
| 219 | Impact of treatment strategy and physical performance on future knee-related self-efficacy in individuals with ACL injury. <i>BMC Musculoskeletal Disorders</i> , 2018, 19, 50. | 0.8 | 10 |
| 220 | On a Trajectory for Success—9 in Every 10 People With a Degenerative Meniscus Tear Have Improved Knee Function Within 2 Years After Treatment: A Secondary Exploratory Analysis of a Randomized Controlled Trial. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2021, 51, 289-297. | 1.7 | 10 |
| 221 | Physical Capacity in Female Soccer Players - Does Age Make a Difference?. <i>Advances in Physiotherapy</i> , 2000, 2, 39-48. | 0.2 | 9 |
| 222 | Exercise-induced hypoalgesia in young adult females with long-standing patellofemoral pain – A randomized crossover study. <i>European Journal of Pain</i> , 2019, 23, 1780-1789. | 1.4 | 9 |
| 223 | Room for improvement: a randomised controlled trial with nested qualitative interviews on space, place and treatment delivery. <i>British Journal of Sports Medicine</i> , 2019, 53, 359-367. | 3.1 | 9 |
| 224 | <p></p>The Danish Hip Arthroscopy Registry: Registration Completeness and Patient Characteristics Between Responders and Non-Responders</p>. <i>Clinical Epidemiology</i> , 2020, Volume 12, 825-833. | 1.5 | 9 |
| 225 | Cost-effectiveness of an 8-week supervised education and exercise therapy programme for knee and hip osteoarthritis: a pre-post analysis of 16%255 patients participating in Good Life with osteoArthritis in Denmark (GLA:D). <i>BMJ Open</i> , 2021, 11, e049541. | 0.8 | 9 |
| 226 | Principles of brain plasticity in improving sensorimotor function of the knee and leg in healthy subjects: A double-blind randomized exploratory trial. <i>BMC Musculoskeletal Disorders</i> , 2009, 10, 99. | 0.8 | 8 |
| 227 | Neuromuscular Function during Stair Descent in Meniscectomized Patients and Controls. <i>Medicine and Science in Sports and Exercise</i> , 2011, 43, 1272-1279. | 0.2 | 8 |
| 228 | Immediate effect of valgus bracing on knee joint moments in meniscectomised patients: An exploratory study. <i>Journal of Science and Medicine in Sport</i> , 2016, 19, 964-969. | 0.6 | 8 |
| 229 | It is time to stop meniscectomy. <i>British Journal of Sports Medicine</i> , 2017, 51, 490-491. | 3.1 | 8 |
| 230 | Patients with osteoarthritis are least likely to receive lifestyle advice compared with patients with diabetes and hypertension: A national health survey study from Denmark. <i>Osteoarthritis and Cartilage</i> Open, 2020, 2, 100067. | 0.9 | 8 |
| 231 | Efficacy of the SOAR knee health program: protocol for a two-arm stepped-wedge randomized delayed-controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2022, 23, 85. | 0.8 | 8 |
| 232 | Principles of brain plasticity in improving sensorimotor function of the knee and leg in patients with anterior cruciate ligament injury: a double-blind randomized exploratory trial. <i>BMC Musculoskeletal Disorders</i> , 2012, 13, 68. | 0.8 | 7 |
| 233 | Prevalence, Severity, and Correlates of Pain Flares in Response to a Repeated Sit-to-Stand Activity: A Cross-sectional Study of 14 902 Patients With Knee and Hip Osteoarthritis in Primary Care. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2020, 50, 309-318. | 1.7 | 7 |
| 234 | What tests should be used to assess functional performance in youth and young adults following anterior cruciate ligament or meniscal injury? A systematic review of measurement properties for the OPTIKNEE consensus. <i>British Journal of Sports Medicine</i> , 2022, 56, 1454-1464. | 3.1 | 7 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 235 | Letter to the Editor. American Journal of Sports Medicine, 2009, 37, 1042-1044. | 1.9 | 6 |
| 236 | The effect of education and supervised exercise vs. education alone on the time to total hip replacement in patients with severe hip osteoarthritis. A randomized clinical trial protocol. BMC Musculoskeletal Disorders, 2013, 14, 21. | 0.8 | 6 |
| 237 | Exploring the effect of space and place on response to exercise therapy for knee and hip pain—a protocol for a double-blind randomised controlled clinical trial: the CONEX trial. BMJ Open, 2015, 5, e007701-e007701. | 0.8 | 6 |
| 238 | The 6-m timed hop test is a prognostic factor for outcomes in patients with meniscal tears treated with exercise therapy or arthroscopic partial meniscectomy: a secondary, exploratory analysis of the Odense “Oslo meniscectomy versus exercise (OMEX) trial. Knee Surgery, Sports Traumatology, Arthroscopy, 2019, 27, 2478-2487. | 2.3 | 6 |
| 239 | Infographic. Therapeutic exercise relieves pain and does not harm knee cartilage nor trigger inflammation. British Journal of Sports Medicine, 2020, 54, 118-119. | 3.1 | 6 |
| 240 | Prevalence of multimorbid degenerative lumbar spinal stenosis with knee and/or hip osteoarthritis: protocol for a systematic review and meta-analysis. Systematic Reviews, 2020, 9, 232. | 2.5 | 6 |
| 241 | Translation, Cross-Cultural Adaptation, and Validation of the Activity Rating Scale for Disorders of the Knee. Orthopaedic Journal of Sports Medicine, 2017, 5, 232596711772936. | 0.8 | 6 |
| 242 | Prognosis and transition of multi-site pain during the course of 5 years: Results of knee pain and function from a prospective cohort study among 756 adolescents. PLoS ONE, 2021, 16, e0250415. | 1.1 | 5 |
| 243 | Prognostic factors for health outcomes after exercise therapy and education in people with knee and hip osteoarthritis with or without comorbidities: a study of 37,576 patients treated in primary care. Arthritis Care and Research, 2021, , . | 1.5 | 5 |
| 244 | Agreement Between Self-Reported Information and Administrative Data on Comorbidities, Imaging and Treatment in Denmark “ A Validation Study of 38,745 Patients with Knee or Hip Osteoarthritis. Clinical Epidemiology, 2021, Volume 13, 779-790. | 1.5 | 5 |
| 245 | Analgesic use in adolescents with patellofemoral pain or Osgood “Schlatter Disease: a secondary cross-sectional analysis of 323 subjects. Scandinavian Journal of Pain, 2022, 22, 543-551. | 0.5 | 5 |
| 246 | On the art of measuring. Acta Orthopaedica, 1997, 68, 3-5. | 1.4 | 4 |
| 247 | Impact of educational level and employment status on short-term and long-term pain relief from supervised exercise therapy and education: an observational study of 22 “ 588 patients with knee and hip osteoarthritis. BMJ Open, 2021, 11, e045156. | 0.8 | 4 |
| 248 | Muscle Strength and Osteoarthritis Progression After Surgery or Exercise for Degenerative Meniscal Tears: Secondary Analyses of a Randomized Trial. Arthritis Care and Research, 2022, 74, 70-78. | 1.5 | 4 |
| 249 | Prognostic factors of total hip replacement during a 2-year period in participants enrolled in supervised education and exercise therapy: a prognostic study of 3657 participants with hip osteoarthritis. Arthritis Research and Therapy, 2021, 23, 235. | 1.6 | 4 |
| 250 | Feasibility of the SOAR (Stop OsteoARthritis) program. Osteoarthritis and Cartilage Open, 2022, 4, 100239. | 0.9 | 4 |
| 251 | Reliability, validity and responsiveness of a new leisure index: The Patient “ Specific Leisure Scale (PSLS). Musculoskeletal Care, 2009, 7, 178-193. | 0.6 | 3 |
| 252 | Association between sensory function and hop performance and self-reported outcomes in patients with anterior cruciate ligament injury. Open Access Journal of Sports Medicine, 2017, Volume 8, 1-8. | 0.6 | 3 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 253 | Proportion of patients with hip osteoarthritis in primary care identified by differing clinical criteria: a cross-sectional study of 4699 patients. <i>Osteoarthritis and Cartilage Open</i> , 2020, 2, 100111. | 0.9 | 3 |
| 254 | Arthroscopic meniscectomy versus non-surgical or sham treatment in patients with MRI confirmed degenerative meniscus lesions: a protocol for an individual participant data meta-analysis. <i>BMJ Open</i> , 2020, 10, e031864. | 0.8 | 3 |
| 255 | Infographic. Roadmap to managing a person with musculoskeletal pain irrespective of body region. <i>British Journal of Sports Medicine</i> , 2020, 54, 554-555. | 3.1 | 3 |
| 256 | Percentile curves for the knee injury and osteoarthritis outcome score in the middle-aged Dutch population. <i>Osteoarthritis and Cartilage</i> , 2020, 28, 1046-1054. | 0.6 | 3 |
| 257 | Factors influencing pain response following GLA:DÂ® patient education and supervised exercise in males and females with hip osteoarthritis. <i>Arthritis Care and Research</i> , 2022, , . | 1.5 | 3 |
| 258 | Implementation of clinical guidelines for osteoarthritis together (IMPACT): protocol for a participatory health research approach to implementing high value care. <i>BMC Musculoskeletal Disorders</i> , 2022, 23, . | 0.8 | 3 |
| 259 | Recommendations for publication of cross-cultural validation studies of patient-reported outcomes (PROs) in Osteoarthritis and Cartilage. <i>Osteoarthritis and Cartilage</i> , 2012, 20, 4-5. | 0.6 | 2 |
| 260 | Authors' response to editorial by Levy and colleagues on treating ACL injuries in young moderately active adults. <i>BMJ, The</i> , 2013, 346, f2082-f2082. | 3.0 | 2 |
| 261 | Infographic. Risk profile for sport-related post-traumatic knee osteoarthritis. <i>British Journal of Sports Medicine</i> , 2019, 54, bjsports-2019-100877. | 3.1 | 2 |
| 262 | Association between received treatment elements and satisfaction with care for patients with knee osteoarthritis seen in general practice in Denmark. <i>Scandinavian Journal of Primary Health Care</i> , 2021, 39, 257-264. | 0.6 | 2 |
| 263 | Knee osteoarthritis after meniscectomy: Prevalence of radiographic changes after twenty-one years, compared with matched controls. , 1998, 41, 687. | | 2 |
| 264 | Exercise Therapy as Treatment for Patients with Osteoarthritis of the Knee. , 2013, , 25-33. | | 1 |
| 265 | Comments on: Validation Study of an Electronic Method of Condensed Outcomes Tools Reporting in Orthopaedics (<i>J Knee Surg</i> 2013;26:445â€“452). <i>Journal of Knee Surgery</i> , 2014, 27, 165-166. | 0.9 | 1 |
| 266 | Authorsâ€™ reply to Joshi. <i>BMJ, The</i> , 2016, 354, i4623. | 3.0 | 1 |
| 267 | Comparative responsiveness of measures of pain and function after total hip replacement. <i>Arthritis and Rheumatism</i> , 2001, 45, 258-262. | 6.7 | 1 |
| 268 | Care-seeking behaviour of adolescents with patellofemoral pain: a retrospective cohort study. <i>F1000Research</i> , 0, 11, 161. | 0.8 | 1 |
| 269 | Prevention and early treatment, a future focus for OA research. <i>Osteoarthritis and Cartilage</i> , 2021, 29, 1627-1629. | 0.6 | 1 |
| 270 | Expert Panels: Can They Be Trusted?. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2013, 29, 1127-1128. | 1.3 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 271 | Authors' reply to Chitnavis. BMJ, The, 2015, 350, h431-h431. | 3.0 | 0 |
| 272 | A more correct interpretation. Cmaj, 2015, 187, 358.1-358. | 0.9 | 0 |
| 273 | Muscle Function at 3 Years Following ACL Injury Affects 5 Year Patient-Reported Outcomes. Medicine and Science in Sports and Exercise, 2016, 48, 301. | 0.2 | 0 |
| 274 | Calculate Patient-Reported Scores as Recommended so as Not to Threaten the Validity of the Measure: Letter to the Editor. Orthopaedic Journal of Sports Medicine, 2018, 6, 232596711881698. | 0.8 | 0 |
| 275 | Response to Letter to Editor: ~2-year outcome from two parallel randomized controlled trials. Reporting considerations™. Osteoarthritis and Cartilage, 2019, 27, e1-e2. | 0.6 | 0 |
| 276 | Single postoperative infusion of zoledronic acid to improve patient-reported outcome after hip or knee replacement: study protocol for a randomised, controlled, double-blinded clinical trial. BMJ Open, 2020, 10, e040985. | 0.8 | 0 |
| 277 | Response to Letter to the Editor: ~Osteoarthritis progression after exercise therapy or meniscectomy in patients with degenerative meniscal tears~. Osteoarthritis and Cartilage, 2020, 28, 1148-1149. | 0.6 | 0 |
| 278 | Erratum. Journal of Bone and Joint Surgery - Series A, 2006, 88, 448. | 1.4 | 0 |
| 279 | PROMs for Osteoarthritis. , 2016, , 249-285. | | 0 |
| 280 | Does Pain Medication Use Influence the Outcome of 8 Weeks of Education and Exercise Therapy in Patients with Knee or Hip Osteoarthritis? An Observational Study. Pain Medicine, 2022, , . | 0.9 | 0 |
| 281 | Changes in medial-to-lateral knee joint loading patterns assessed by Dual-Energy X-ray Absorptiometry following supervised neuromuscular exercise therapy and patient education in patients with knee osteoarthritis: an exploratory cohort study. Physiotherapy Theory and Practice, 2022, , 1-10. | 0.6 | 0 |