

Ewa M Roos

List of Publications by Citations

Source: <https://exaly.com/author-pdf/203178/ewa-m-roos-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

261
papers

24,488
citations

70
h-index

154
g-index

287
ext. papers

28,084
ext. citations

4.7
avg. IF

7.19
L-index

#	Paper	IF	Citations
261	Knee Injury and Osteoarthritis Outcome Score (KOOS)--development of a self-administered outcome measure. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 1998 , 28, 88-96	4.2	2318
260	OARSI guidelines for the non-surgical management of knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2014 , 22, 363-88	6.2	1808
259	The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. <i>American Journal of Sports Medicine</i> , 2007 , 35, 1756-69	6.8	1548
258	The Knee injury and Osteoarthritis Outcome Score (KOOS): from joint injury to osteoarthritis. <i>Health and Quality of Life Outcomes</i> , 2003 , 1, 64	3	1173
257	Measures of knee function: International Knee Documentation Committee (IKDC) Subjective Knee Evaluation Form, Knee Injury and Osteoarthritis Outcome Score (KOOS), Knee Injury and Osteoarthritis Outcome Score Physical Function Short Form (KOOS-PS), Knee Outcome Survey Activities of Daily Living Scale (KOS-ADL), Lysholm Knee Scoring Scale, Oxford Knee Score (OKS).	4.7	661
256	Knee injury and Osteoarthritis Outcome Score (KOOS) - validation and comparison to the WOMAC in total knee replacement. <i>Health and Quality of Life Outcomes</i> , 2003 , 1, 17 (Suppl 1), S208-28	3	623
255	Hip disability and osteoarthritis outcome score (HOOS)--validity and responsiveness in total hip replacement. <i>BMC Musculoskeletal Disorders</i> , 2003 , 4, 10	2.8	613
254	High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes. <i>Annals of the Rheumatic Diseases</i> , 2004 , 63, 269-73	2.4	598
253	Knee osteoarthritis after meniscectomy: prevalence of radiographic changes after twenty-one years, compared with matched controls. <i>Arthritis and Rheumatism</i> , 1998 , 41, 687-93		557
252	A randomized trial of treatment for acute anterior cruciate ligament tears. <i>New England Journal of Medicine</i> , 2010 , 363, 331-42	59.2	532
251	Validation of the foot and ankle outcome score for ankle ligament reconstruction. <i>Foot and Ankle International</i> , 2001 , 22, 788-94	3.3	501
250	Non-contact ACL injuries in female athletes: an International Olympic Committee current concepts statement. <i>British Journal of Sports Medicine</i> , 2008 , 42, 394-412	10.3	442
249	Knee injury and Osteoarthritis Outcome Score (KOOS)--validation of a Swedish version. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 1998 , 8, 439-48	4.6	417
248	OARSI recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2013 , 21, 1042-52	6.2	388
247	Positive effects of moderate exercise on glycosaminoglycan content in knee cartilage: a four-month, randomized, controlled trial in patients at risk of osteoarthritis. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3507-14		338
246	A Randomized, Controlled Trial of Total Knee Replacement. <i>New England Journal of Medicine</i> , 2015 , 373, 1597-606	59.2	336
245	The Copenhagen Hip and Groin Outcome Score (HAGOS): development and validation according to the COSMIN checklist. <i>British Journal of Sports Medicine</i> , 2011 , 45, 478-91	10.3	328

244	Psychometric properties of patient-reported outcome measures for hip arthroscopic surgery. <i>American Journal of Sports Medicine</i> , 2013 , 41, 2065-73	6.8	321
243	Joint injury causes knee osteoarthritis in young adults. <i>Current Opinion in Rheumatology</i> , 2005 , 17, 195-200	9.0	306
242	Impact of exercise type and dose on pain and disability in knee osteoarthritis: a systematic review and meta-regression analysis of randomized controlled trials. <i>Arthritis and Rheumatology</i> , 2014 , 66, 622-36	9.5	302
241	Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. <i>BMJ, The</i> , 2013 , 346, f232	5.9	282
240	Knee Injury and Osteoarthritis Outcome Score (KOOS): systematic review and meta-analysis of measurement properties. <i>Osteoarthritis and Cartilage</i> , 2016 , 24, 1317-29	6.2	260
239	Knee arthroplasty: are patients' expectations fulfilled? A prospective study of pain and function in 102 patients with 5-year follow-up. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2009 , 80, 55-61	4.3	231
238	Clinical improvement after 6 weeks of eccentric exercise in patients with mid-portion Achilles tendinopathy -- a randomized trial with 1-year follow-up. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2004 , 14, 286-95	4.6	227
237	Knee complaints vary with age and gender in the adult population. Population-based reference data for the Knee injury and Osteoarthritis Outcome Score (KOOS). <i>BMC Musculoskeletal Disorders</i> , 2006 , 7, 38	2.8	222
236	Patient-relevant outcomes fourteen years after meniscectomy: influence of type of meniscal tear and size of resection. <i>Rheumatology</i> , 2001 , 40, 631-9	3.9	210
235	Predictors of patient relevant outcome after total hip replacement for osteoarthritis: a prospective study. <i>Annals of the Rheumatic Diseases</i> , 2003 , 62, 923-30	2.4	195
234	Strategies for the prevention of knee osteoarthritis. <i>Nature Reviews Rheumatology</i> , 2016 , 12, 92-101	8.1	182
233	Measurement properties of performance-based measures to assess physical function in hip and knee osteoarthritis: a systematic review. <i>Osteoarthritis and Cartilage</i> , 2012 , 20, 1548-62	6.2	177
232	Long-term outcome of meniscectomy: symptoms, function, and performance tests in patients with or without radiographic osteoarthritis compared to matched controls. <i>Osteoarthritis and Cartilage</i> , 2001 , 9, 316-24	6.2	172
231	Sex differences in patient-reported outcomes after anterior cruciate ligament reconstruction: data from the Swedish knee ligament register. <i>American Journal of Sports Medicine</i> , 2010 , 38, 1334-42	6.8	153
230	Isokinetic knee extensor strength and functional performance in healthy female soccer players. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 1998 , 8, 257-64	4.6	138
229	The development of a short measure of physical function for knee OA KOOS-Physical Function Shortform (KOOS-PS) - an OARSI/OMERACT initiative. <i>Osteoarthritis and Cartilage</i> , 2008 , 16, 542-50	6.2	138
228	Good Life with osteoArthritis in Denmark (GLA:D) evidence-based education and supervised neuromuscular exercise delivered by certified physiotherapists nationwide. <i>BMC Musculoskeletal Disorders</i> , 2017 , 18, 72	2.8	133
227	Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle aged patients: randomised controlled trial with two year follow-up. <i>BMJ, The</i> , 2016 , 354, i3740	5.9	133

226	Cross-cultural adaptation and validation of the French version of the Knee injury and Osteoarthritis Outcome Score (KOOS) in knee osteoarthritis patients. <i>Osteoarthritis and Cartilage</i> , 2008 , 16, 423-8	6.2	132
225	The role of pain and functional impairment in the decision to recommend total joint replacement in hip and knee osteoarthritis: an international cross-sectional study of 1909 patients. Report of the OARSI-OMERACT Task Force on total joint replacement. <i>Osteoarthritis and Cartilage</i> , 2011 , 19, 147-54	6.2	131
224	Validity, reliability and responsiveness of patient-reported outcome questionnaires when assessing hip and groin disability: a systematic review. <i>British Journal of Sports Medicine</i> , 2010 , 44, 1186-96	10.3	128
223	Validation of the Dutch version of the Hip disability and Osteoarthritis Outcome Score. <i>Osteoarthritis and Cartilage</i> , 2007 , 15, 104-9	6.2	127
222	Muscle weakness, afferent sensory dysfunction and exercise in knee osteoarthritis. <i>Nature Reviews Rheumatology</i> , 2011 , 7, 57-63	8.1	122
221	Validity and inter-rater reliability of medio-lateral knee motion observed during a single-limb mini squat. <i>BMC Musculoskeletal Disorders</i> , 2010 , 11, 265	2.8	122
220	Reduced functional performance in the lower extremity predicted radiographic knee osteoarthritis five years later. <i>Annals of the Rheumatic Diseases</i> , 2004 , 63, 402-7	2.4	121
219	The acutely ACL injured knee assessed by MRI: changes in joint fluid, bone marrow lesions, and cartilage during the first year. <i>Osteoarthritis and Cartilage</i> , 2009 , 17, 161-7	6.2	118
218	The development of a short measure of physical function for hip OA HOOS-Physical Function Shortform (HOOS-PS): an OARSI/OMERACT initiative. <i>Osteoarthritis and Cartilage</i> , 2008 , 16, 551-9	6.2	118
217	WOMAC Osteoarthritis Index--additional dimensions for use in subjects with post-traumatic osteoarthritis of the knee. Western Ontario and MacMaster Universities. <i>Osteoarthritis and Cartilage</i> , 1999 , 7, 216-21	6.2	117
216	A 5 year prospective study of patient-relevant outcomes after total knee replacement. <i>Osteoarthritis and Cartilage</i> , 2009 , 17, 601-6	6.2	115
215	Magnitude and meaningfulness of change in SF-36 scores in four types of orthopedic surgery. <i>Health and Quality of Life Outcomes</i> , 2008 , 6, 55	3	115
214	Muscle strength and functional performance in patients with anterior cruciate ligament injury treated with training and surgical reconstruction or training only: a two to five-year followup. <i>Arthritis and Rheumatism</i> , 2008 , 59, 1773-9		114
213	Patient-Reported Outcome (PRO) questionnaires for young to middle-aged adults with hip and groin disability: a systematic review of the clinimetric evidence. <i>British Journal of Sports Medicine</i> , 2015 , 49, 812	10.3	113
212	Is Knee Pain During Adolescence a Self-limiting Condition? Prognosis of Patellofemoral Pain and Other Types of Knee Pain. <i>American Journal of Sports Medicine</i> , 2016 , 44, 1165-71	6.8	109
211	Feasibility of neuromuscular training in patients with severe hip or knee OA: the individualized goal-based NEMEX-TJR training program. <i>BMC Musculoskeletal Disorders</i> , 2010 , 11, 126	2.8	109
210	Effectiveness and practice variation of rehabilitation after joint replacement. <i>Current Opinion in Rheumatology</i> , 2003 , 15, 160-2	5.3	104
209	Neuromuscular versus quadriceps strengthening exercise in patients with medial knee osteoarthritis and varus malalignment: a randomized controlled trial. <i>Arthritis and Rheumatology</i> , 2014 , 66, 950-9	9.5	98

208	Patient-reported outcomes for total hip and knee arthroplasty: commonly used instruments and attributes of a "good" measure. <i>Clinics in Geriatric Medicine</i> , 2012 , 28, 367-94	3.8	97
207	Comparative, validity and responsiveness of the HOOS-PS and KOOS-PS to the WOMAC physical function subscale in total joint replacement for osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2009 , 17, 843-7	6.2	96
206	The acutely ACL injured knee assessed by MRI: are large volume traumatic bone marrow lesions a sign of severe compression injury?. <i>Osteoarthritis and Cartilage</i> , 2008 , 16, 829-36	6.2	91
205	Knee osteoarthritis risk is increased 4-6 fold after knee injury - a systematic review and meta-analysis. <i>British Journal of Sports Medicine</i> , 2019 , 53, 1454-1463	10.3	88
204	Osteoarthritis: Models for appropriate care across the disease continuum. <i>Best Practice and Research in Clinical Rheumatology</i> , 2016 , 30, 503-535	5.3	87
203	Exercise during school hours when added to patient education improves outcome for 2 years in adolescent patellofemoral pain: a cluster randomised trial. <i>British Journal of Sports Medicine</i> , 2015 , 49, 406-12	10.3	86
202	Minimal clinically important improvement (MCII) and patient-acceptable symptom state (PASS) in total hip arthroplasty (THA) patients 1 year postoperatively. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2014 , 85, 39-48	4.3	86
201	Clinical update: treating osteoarthritis. <i>Lancet, The</i> , 2007 , 370, 2082-4	4.0	85
200	ICRS Recommendation Document: Patient-Reported Outcome Instruments for Use in Patients with Articular Cartilage Defects. <i>Cartilage</i> , 2011 , 2, 122-36	3	84
199	Knee extension and flexion muscle power after anterior cruciate ligament reconstruction with patellar tendon graft or hamstring tendons graft: a cross-sectional comparison 3 years post surgery. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2009 , 17, 162-9	5.5	83
198	Lower mechanical pressure pain thresholds in female adolescents with patellofemoral pain syndrome. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2013 , 43, 414-21	4.2	81
197	Serum levels of cartilage oligomeric matrix protein (COMP) increase temporarily after physical exercise in patients with knee osteoarthritis. <i>BMC Musculoskeletal Disorders</i> , 2006 , 7, 98	2.8	81
196	Foot orthoses for the treatment of plantar fasciitis. <i>Foot and Ankle International</i> , 2006 , 27, 606-11	3.3	81
195	High prevalence of daily and multi-site pain--a cross-sectional population-based study among 3000 Danish adolescents. <i>BMC Pediatrics</i> , 2013 , 13, 191	2.6	80
194	Muscle strength, functional performance, and self-reported outcomes four years after arthroscopic partial meniscectomy in middle-aged patients. <i>Arthritis and Rheumatism</i> , 2006 , 55, 946-52		80
193	Substantial disability 3 months after arthroscopic partial meniscectomy: A prospective study of patient-relevant outcomes. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2000 , 16, 619-26	5.4	80
192	The effect of eight weeks of exercise on knee adduction moment in early knee osteoarthritis--a pilot study. <i>Osteoarthritis and Cartilage</i> , 2007 , 15, 1163-70	6.2	74
191	Comparative responsiveness of measures of pain and function after total hip replacement. <i>Arthritis and Rheumatism</i> , 2001 , 45, 258-62		70

190	Cross-cultural adaptation and validation of Singapore English and Chinese versions of the Knee injury and Osteoarthritis Outcome Score (KOOS) in Asians with knee osteoarthritis in Singapore. <i>Osteoarthritis and Cartilage</i> , 2006 , 14, 1098-103	6.2	68
189	Six-week high-intensity exercise program for middle-aged patients with knee osteoarthritis: a randomized controlled trial [ISRCTN20244858]. <i>BMC Musculoskeletal Disorders</i> , 2005 , 6, 27	2.8	67
188	OMERACT/OARSI initiative to define states of severity and indication for joint replacement in hip and knee osteoarthritis. <i>Journal of Rheumatology</i> , 2007 , 34, 1432-5	4.1	66
187	Postoperative effects of neuromuscular exercise prior to hip or knee arthroplasty: a randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1130-7	2.4	64
186	Cross-cultural adaptation and validation of the French version of the Hip disability and Osteoarthritis Outcome Score (HOOS) in hip osteoarthritis patients. <i>Osteoarthritis and Cartilage</i> , 2010 , 18, 522-9	6.2	63
185	Validity and reliability of functional performance tests in meniscectomized patients with or without knee osteoarthritis. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2007 , 17, 120-7	4.6	59
184	Development of the Knee Injury and Osteoarthritis Outcome Score for children (KOOS-Child): comprehensibility and content validity. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2012 , 83, 666-73	4.3	57
183	Proportion of Patients Reporting Acceptable Symptoms or Treatment Failure and Their Associated KOOS Values at 6 to 24 Months After Anterior Cruciate Ligament Reconstruction: A Study From the Norwegian Knee Ligament Registry. <i>American Journal of Sports Medicine</i> , 2015 , 43, 1902-7	6.8	56
182	Immediate efficacy of neuromuscular exercise in patients with severe osteoarthritis of the hip or knee: a secondary analysis from a randomized controlled trial. <i>Journal of Rheumatology</i> , 2014 , 41, 1385-94 ¹	4.1	54
181	Variation in age and physical status prior to total knee and hip replacement surgery: a comparison of centers in Australia and Europe. <i>Arthritis and Rheumatism</i> , 2009 , 61, 166-73		54
180	Total knee replacement and non-surgical treatment of knee osteoarthritis: 2-year outcome from two parallel randomized controlled trials. <i>Osteoarthritis and Cartilage</i> , 2018 , 26, 1170-1180	6.2	53
179	Osteoarthritis 2012 year in review: rehabilitation and outcomes. <i>Osteoarthritis and Cartilage</i> , 2012 , 20, 1477-83	6.2	52
178	Neuromuscular exercise as treatment of degenerative knee disease. <i>Exercise and Sport Sciences Reviews</i> , 2015 , 43, 14-22	6.7	51
177	Effects of neuromuscular training (NEMEX-TJR) on patient-reported outcomes and physical function in severe primary hip or knee osteoarthritis: a controlled before-and-after study. <i>BMC Musculoskeletal Disorders</i> , 2013 , 14, 232	2.8	51
176	The challenge of recruiting patients with anterior cruciate ligament injury of the knee into a randomized clinical trial comparing surgical and non-surgical treatment. <i>Contemporary Clinical Trials</i> , 2007 , 28, 295-302	2.3	51
175	Establishing outcome measures in early knee osteoarthritis. <i>Nature Reviews Rheumatology</i> , 2019 , 15, 438-448	8.1	50
174	Measurement properties of performance-based outcome measures to assess physical function in young and middle-aged people known to be at high risk of hip and/or knee osteoarthritis: a systematic review. <i>Osteoarthritis and Cartilage</i> , 2014 , 22, 26-39	6.2	49
173	Lower extremity performance following ACL rehabilitation in the KANON-trial: impact of reconstruction and predictive value at 2 and 5 years. <i>British Journal of Sports Medicine</i> , 2013 , 47, 980-5	10.3	49

172	Hip and knee strength is not affected in 12-16 year old adolescents with patellofemoral pain--a cross-sectional population-based study. <i>PLoS ONE</i> , 2013 , 8, e79153	3.7	49
171	Increased pain sensitivity but normal function of exercise induced analgesia in hip and knee osteoarthritis--treatment effects of neuromuscular exercise and total joint replacement. <i>Osteoarthritis and Cartilage</i> , 2013 , 21, 1299-307	6.2	48
170	The efficacy of 12 weeks non-surgical treatment for patients not eligible for total knee replacement: a randomized controlled trial with 1-year follow-up. <i>Osteoarthritis and Cartilage</i> , 2015 , 23, 1465-75	6.2	47
169	Change in self-reported outcomes and objective physical function over 7 years in middle-aged subjects with or at high risk of knee osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2008 , 67, 505-10	2.4	47
168	Patterns of musculoskeletal pain in the population: a latent class analysis using a nationally representative interviewer-based survey of 4817 Danes. <i>European Journal of Pain</i> , 2013 , 17, 452-60	3.7	46
167	The patellofemoral pain and osteoarthritis subscale of the KOOS (KOOS-PF): development and validation using the COSMIN checklist. <i>British Journal of Sports Medicine</i> , 2018 , 52, 1130-1136	10.3	44
166	The effect of patient characteristics on variability in pain and function over two years in early knee osteoarthritis. <i>Health and Quality of Life Outcomes</i> , 2005 , 3, 59	3	44
165	How do middle-aged patients conceive exercise as a form of treatment for knee osteoarthritis?. <i>Disability and Rehabilitation</i> , 2006 , 28, 51-9	2.4	44
164	Impact of exercise on articular cartilage in people at risk of, or with established, knee osteoarthritis: a systematic review of randomised controlled trials. <i>British Journal of Sports Medicine</i> , 2019 , 53, 940-947	10.3	42
163	Meaningful Change Scores in the Knee Injury and Osteoarthritis Outcome Score in Patients Undergoing Anterior Cruciate Ligament Reconstruction. <i>American Journal of Sports Medicine</i> , 2018 , 46, 1120-1128	6.8	41
162	Psychometric properties of the Knee injury and Osteoarthritis Outcome Score for Children (KOOS-Child) in children with knee disorders. <i>British Journal of Sports Medicine</i> , 2014 , 48, 1437-46	10.3	41
161	Delaying ACL reconstruction and treating with exercise therapy alone may alter prognostic factors for 5-year outcome: an exploratory analysis of the KANON trial. <i>British Journal of Sports Medicine</i> , 2017 , 51, 1622-1629	10.3	39
160	Minimal important change values for the Oxford Knee Score and the Forgotten Joint Score at 1 year after total knee replacement. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018 , 89, 541-547	4.3	38
159	Care-seeking behaviour of adolescents with knee pain: a population-based study among 504 adolescents. <i>BMC Musculoskeletal Disorders</i> , 2013 , 14, 225	2.8	38
158	A 12-week exercise therapy program in middle-aged patients with degenerative meniscus tears: a case series with 1-year follow-up. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2012 , 42, 919-31	4.2	38
157	Neuromuscular activity and knee kinematics in adolescents with patellofemoral pain. <i>Medicine and Science in Sports and Exercise</i> , 2013 , 45, 1730-9	1.2	38
156	Effect of exercise therapy compared with arthroscopic surgery on knee muscle strength and functional performance in middle-aged patients with degenerative meniscus tears: a 3-mo follow-up of a randomized controlled trial. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2015 , 94, 460-73	2.6	37
155	Effects of functional exercise training on performance and muscle strength after meniscectomy: a randomized trial. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2009 , 19, 156-65	4.6	37

154	Effect of leisure time physical activity on severe knee or hip osteoarthritis leading to total joint replacement: a population-based prospective cohort study. <i>BMC Musculoskeletal Disorders</i> , 2012 , 13, 73	2.8	36
153	Validation of the Rheumatoid and Arthritis Outcome Score (RAOS) for the lower extremity. <i>Health and Quality of Life Outcomes</i> , 2003 , 1, 55	3	36
152	Pain trajectory and exercise-induced pain flares during 8 weeks of neuromuscular exercise in individuals with knee and hip pain. <i>Osteoarthritis and Cartilage</i> , 2016 , 24, 589-92	6.2	35
151	Comparison of neuromuscular and quadriceps strengthening exercise in the treatment of varus malaligned knees with medial knee osteoarthritis: a randomised controlled trial protocol. <i>BMC Musculoskeletal Disorders</i> , 2011 , 12, 276	2.8	35
150	Effectiveness and safety of strengthening, aerobic, and coordination exercises for patients with osteoarthritis. <i>Current Opinion in Rheumatology</i> , 2003 , 15, 141-4	5.3	35
149	Arthroscopic partial meniscectomy in middle-aged patients with mild or no knee osteoarthritis: a protocol for a double-blind, randomized sham-controlled multi-centre trial. <i>BMC Musculoskeletal Disorders</i> , 2013 , 14, 71	2.8	34
148	Choosing surgery: patients' preferences within a trial of treatments for anterior cruciate ligament injury. A qualitative study. <i>BMC Musculoskeletal Disorders</i> , 2009 , 10, 100	2.8	34
147	Examining the Minimal Important Difference of Patient-reported Outcome Measures for Individuals with Knee Osteoarthritis: A Model Using the Knee Injury and Osteoarthritis Outcome Score. <i>Journal of Rheumatology</i> , 2016 , 43, 395-404	4.1	33
146	Feasibility of 4 patient-reported outcome measures in a registry setting. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2012 , 83, 321-7	4.3	33
145	A 2-year prospective study of patient-relevant outcomes in patients operated on for knee osteoarthritis with tibial osteotomy. <i>BMC Musculoskeletal Disorders</i> , 2005 , 6, 18	2.8	33
144	The OMERACT-OARSI Core Domain Set for Measurement in Clinical Trials of Hip and/or Knee Osteoarthritis. <i>Journal of Rheumatology</i> , 2019 , 46, 981-989	4.1	33
143	Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. <i>British Journal of Sports Medicine</i> , 2015 , 49, 700	10.3	32
142	Early intervention for adolescents with patellofemoral pain syndrome--a pragmatic cluster randomised controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2012 , 13, 9	2.8	31
141	Cross-cultural translation and measurement properties of the Polish version of the Knee injury and Osteoarthritis Outcome Score (KOOS) following anterior cruciate ligament reconstruction. <i>Health and Quality of Life Outcomes</i> , 2013 , 11, 107	3	31
140	Agreement and reliability of functional performance and muscle power in patients with advanced osteoarthritis of the hip or knee. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2012 , 91, 401-10	2.6	31
139	Psychometric properties of the French translation of the reduced KOOS and HOOS (KOOS-PS and HOOS-PS). <i>Osteoarthritis and Cartilage</i> , 2009 , 17, 1604-8	6.2	29
138	Patient education with or without manual therapy compared to a control group in patients with osteoarthritis of the hip. A proof-of-principle three-arm parallel group randomized clinical trial. <i>Osteoarthritis and Cartilage</i> , 2013 , 21, 1494-503	6.2	28
137	Relationships between postural orientation and self reported function, hop performance and muscle power in subjects with anterior cruciate ligament injury. <i>BMC Musculoskeletal Disorders</i> , 2010 , 11, 143	2.8	28

136	Supervised neuromuscular exercise prior to hip and knee replacement: 12-month clinical effect and cost-utility analysis alongside a randomised controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2017 , 18, 5	2.8	27
135	A pragmatic approach to prevent post-traumatic osteoarthritis after sport or exercise-related joint injury. <i>Best Practice and Research in Clinical Rheumatology</i> , 2019 , 33, 158-171	5.3	27
134	Over-optimistic patient expectations of recovery and leisure activities after arthroscopic meniscus surgery. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2016 , 87, 615-621	4.3	27
133	Influence of Biomechanical Characteristics on Pain and Function Outcomes From Exercise in Medial Knee Osteoarthritis and Varus Malalignment: Exploratory Analyses From a Randomized Controlled Trial. <i>Arthritis Care and Research</i> , 2015 , 67, 1281-1288	4.7	27
132	GLA:D Back group-based patient education integrated with exercises to support self-management of back pain development, theories and scientific evidence. <i>BMC Musculoskeletal Disorders</i> , 2018 , 19, 418	2.8	27
131	Impact of a daily exercise dose on knee joint cartilage - a systematic review and meta-analysis of randomized controlled trials in healthy animals. <i>Osteoarthritis and Cartilage</i> , 2017 , 25, 1223-1237	6.2	26
130	A 12-item short form of the Knee injury and Osteoarthritis Outcome Score (KOOS-12): tests of reliability, validity and responsiveness. <i>Osteoarthritis and Cartilage</i> , 2019 , 27, 762-770	6.2	26
129	Self-reported musculoskeletal pain predicts long-term increase in general health care use: a population-based cohort study with 20-year follow-up. <i>Scandinavian Journal of Public Health</i> , 2014 , 42, 698-704	3	26
128	Psychometric properties of the Neck OutCome Score, Neck Disability Index, and Short Form-36 were evaluated in patients with neck pain. <i>Journal of Clinical Epidemiology</i> , 2016 , 79, 31-40	5.7	26
127	Half of 12-15-year-olds with knee pain still have pain after one year. <i>Danish Medical Journal</i> , 2013 , 60, A4725	3.8	25
126	Five-year prognosis and impact of adolescent knee pain: a prospective population-based cohort study of 504 adolescents in Denmark. <i>BMJ Open</i> , 2019 , 9, e024113	3	24
125	Effect of preoperative neuromuscular training (NEMEX-TJR) on functional outcome after total knee replacement: an assessor-blinded randomized controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2015 , 16, 101	2.8	24
124	Clinical outcomes assessment in clinical trials to assess treatment of femoroacetabular impingement: use of patient-reported outcome measures. <i>Journal of the American Academy of Orthopaedic Surgeons, The</i> , 2013 , 21 Suppl 1, S39-46	4.5	24
123	Reliability and reference values of two clinical measurements of dynamic and static knee position in healthy children. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2011 , 19, 2060-6	5.5	24
122	Knee kinematics and kinetics in former soccer players with a 16-year-old ACL injury--the effects of twelve weeks of knee-specific training. <i>BMC Musculoskeletal Disorders</i> , 2007 , 8, 35	2.8	24
121	Knee function and knee muscle strength in middle-aged patients with degenerative meniscal tears eligible for arthroscopic partial meniscectomy. <i>British Journal of Sports Medicine</i> , 2014 , 48, 784-8	10.3	23
120	Impact of Exercise Therapy on Molecular Biomarkers Related to Cartilage and Inflammation in Individuals at Risk of, or With Established, Knee Osteoarthritis: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. <i>Arthritis Care and Research</i> , 2019 , 71, 1504-1515	4.7	23
119	Young females with long-standing patellofemoral pain display impaired conditioned pain modulation, increased temporal summation of pain, and widespread hyperalgesia. <i>Pain</i> , 2018 , 159, 2530-2537	8.2	23

118	Efficacy of multimodal, systematic non-surgical treatment of knee osteoarthritis for patients not eligible for a total knee replacement: a study protocol of a randomised controlled trial. <i>BMJ Open</i> , 2012 , 2,	3	22
117	Similar group mean scores, but large individual variations, in patient-relevant outcomes over 2 years in meniscectomized subjects with and without radiographic knee osteoarthritis. <i>Health and Quality of Life Outcomes</i> , 2004 , 2, 38	3	22
116	Item selection for 12-item short forms of the Knee injury and Osteoarthritis Outcome Score (KOOS-12) and Hip disability and Osteoarthritis Outcome Score (HOOS-12). <i>Osteoarthritis and Cartilage</i> , 2019 , 27, 746-753	6.2	22
115	OARSI Clinical Trials Recommendations: Design and conduct of clinical trials for primary prevention of osteoarthritis by joint injury prevention in sport and recreation. <i>Osteoarthritis and Cartilage</i> , 2015 , 23, 815-25	6.2	21
114	A Randomized, Controlled Trial of Total Knee Replacement. <i>New England Journal of Medicine</i> , 2016 , 374, 692	59.2	21
113	Thigh muscle strength, functional capacity, and self-reported function in patients at high risk of knee osteoarthritis compared with controls. <i>Arthritis Care and Research</i> , 2010 , 62, 1244-51	4.7	21
112	Three steps to changing the narrative about knee osteoarthritis care: a call to action. <i>British Journal of Sports Medicine</i> , 2020 , 54, 256-258	10.3	21
111	The challenge of recruiting patients into a placebo-controlled surgical trial. <i>Trials</i> , 2014 , 15, 167	2.8	20
110	Group education and exercise is feasible in knee and hip osteoarthritis. <i>Danish Medical Journal</i> , 2012 , 59, A4554	3.8	20
109	Criteria used when deciding on eligibility for total knee arthroplasty--Between thinking and doing. <i>Knee</i> , 2016 , 23, 300-5	2.6	19
108	Muscle strength and functional performance in patients at high risk of knee osteoarthritis: a follow-up study. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2012 , 20, 1110-7	5.5	19
107	Muscle function is associated with future patient-reported outcomes in young adults with ACL injury. <i>BMJ Open Sport and Exercise Medicine</i> , 2016 , 2, e000154	3.4	19
106	The efficacy of non-surgical treatment on pain and sensitization in patients with knee osteoarthritis: a pre-defined ancillary analysis from a randomized controlled trial. <i>Osteoarthritis and Cartilage</i> , 2016 , 24, 108-16	6.2	17
105	Better outcome from arthroscopic partial meniscectomy than skin incisions only? A sham-controlled randomised trial in patients aged 35-55 years with knee pain and an MRI-verified meniscal tear. <i>BMJ Open</i> , 2018 , 8, e019461	3	17
104	The impact of physical activity level on the short- and long-term pain relief from supervised exercise therapy and education: a study of 12,796 Danish patients with knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2018 , 26, 1474-1478	6.2	16
103	Middle-aged patients with an MRI-verified medial meniscal tear report symptoms commonly associated with knee osteoarthritis. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2017 , 88, 664-669	4.3	16
102	The effect on knee-joint load of instruction in analgesic use compared with neuromuscular exercise in patients with knee osteoarthritis: study protocol for a randomized, single-blind, controlled trial (the EXERPHARMA trial). <i>Trials</i> , 2014 , 15, 444	2.8	16
101	Total knee replacement plus physical and medical therapy or treatment with physical and medical therapy alone: a randomised controlled trial in patients with knee osteoarthritis (the MEDIC-study). <i>BMC Musculoskeletal Disorders</i> , 2012 , 13, 67	2.8	16

100	It is good to feel better, but better to feel good: whether a patient finds treatment successful or not depends on the questions researchers ask. <i>British Journal of Sports Medicine</i> , 2019 , 53, 1474-1478	10.3	14
99	Reliability and validity of videotaped functional performance tests in ACL-injured subjects. <i>Physiotherapy Research International</i> , 2008 , 13, 119-30	1.8	14
98	Targeted prevention in primary care aimed at lifestyle-related diseases: a study protocol for a non-randomised pilot study. <i>BMC Family Practice</i> , 2018 , 19, 124	2.6	14
97	Predictors of long-term effect from education and exercise in patients with knee and hip pain. <i>Danish Medical Journal</i> , 2014 , 61, A4867	3.8	14
96	Self-reported Recovery is Associated With Improvement in Localized Hyperalgesia Among Adolescent Females With Patellofemoral Pain: Results From a Cluster Randomized Trial. <i>Clinical Journal of Pain</i> , 2016 , 32, 428-34	3.5	13
95	Republished research: treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. <i>British Journal of Sports Medicine</i> , 2013 , 47, 373	10.3	13
94	Development of a patient-reported outcome: the Neck OutCome Score (NOOS)--Content and construct validity. <i>Journal of Rehabilitation Medicine</i> , 2015 , 47, 844-53	3.4	13
93	Validity and responsiveness of the Clubfoot Assessment Protocol (CAP). A methodological study. <i>BMC Musculoskeletal Disorders</i> , 2006 , 7, 28	2.8	13
92	Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle aged patients: randomised controlled trial with two year follow-up. <i>British Journal of Sports Medicine</i> , 2016 , 50, 1473-1480	10.3	13
91	A 12-item short form of the Hip disability and Osteoarthritis Outcome Score (HOOS-12): tests of reliability, validity and responsiveness. <i>Osteoarthritis and Cartilage</i> , 2019 , 27, 754-761	6.2	13
90	Effect of exercise therapy on neuromuscular activity and knee strength in female adolescents with patellofemoral pain-An ancillary analysis of a cluster randomized trial. <i>Clinical Biomechanics</i> , 2016 , 34, 22-9	2.2	12
89	Response to Letter to the Editor entitled "Comments on OARSI guidelines for the non-surgical management of knee osteoarthritis". <i>Osteoarthritis and Cartilage</i> , 2014 , 22, 890-1	6.2	12
88	Clinical Outcomes Assessment in Clinical Trials to Assess Treatment of Femoroacetabular Impingement: Use of Patient-reported Outcome Measures. <i>Journal of the American Academy of Orthopaedic Surgeons</i> , 2013 , 21, S39-S46	4.5	12
87	Comparison of three sets of clinical classification criteria for knee osteoarthritis: a cross-sectional study of 13,459 patients treated in primary care. <i>Osteoarthritis and Cartilage</i> , 2020 , 28, 167-172	6.2	12
86	Uptake of the OMERACT-OARSI Hip and Knee Osteoarthritis Core Outcome Set: Review of Randomized Controlled Trials from 1997 to 2017. <i>Journal of Rheumatology</i> , 2019 , 46, 976-980	4.1	11
85	Vibratory perception threshold in young and middle-aged patients at high risk of knee osteoarthritis compared to controls. <i>Arthritis Care and Research</i> , 2012 , 64, 144-8	4.7	11
84	The iHOT-33: how valid is it?. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2012 , 28, 1194-5; author reply 1195-6	5.4	11
83	Association between knee alignment and knee pain in patients surgically treated for medial knee osteoarthritis by high tibial osteotomy. A one year follow-up study. <i>BMC Musculoskeletal Disorders</i> , 2009 , 10, 154	2.8	11

82	Neuromuscular function during a forward lunge in meniscectomized patients. <i>Medicine and Science in Sports and Exercise</i> , 2012 , 44, 1358-65	1.2	11
81	A Pragmatic Approach to the Implementation of Osteoarthritis Guidelines Has Fewer Potential Barriers Than Recommended Implementation Frameworks. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2019 , 49, 1-4	4.2	11
80	Implementing a national first-line management program for moderate-severe knee osteoarthritis in Australia: A budget impact analysis focusing on knee replacement avoidance. <i>Osteoarthritis and Cartilage Open</i> , 2020 , 2, 100070	1.5	10
79	The Association Between Knee Confidence and Muscle Power, Hop Performance, and Postural Orientation in People With Anterior Cruciate Ligament Injury. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2016 , 46, 477-82	4.2	10
78	Patients report inferior quality of care for knee osteoarthritis prior to assessment for knee replacement surgery - a cross-sectional study of 517 patients in Denmark. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2020 , 91, 82-87	4.3	10
77	Knee Confidence as It Relates to Self-reported and Objective Correlates of Knee Osteoarthritis: A Cross-sectional Study of 220 Patients. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2015 , 45, 765-77	4.2	9
76	Development of osteoarthritis in patients with degenerative meniscal tears treated with exercise therapy or surgery: a randomized controlled trial. <i>Osteoarthritis and Cartilage</i> , 2020 , 28, 897-906	6.2	9
75	Association of osteoarthritis risk factors with knee and hip pain in a population-based sample of 29-59-year olds in Denmark: a cross-sectional analysis. <i>BMC Musculoskeletal Disorders</i> , 2018 , 19, 300	2.8	9
74	Complex Tears, Extrusion, and Larger Excision Are Prognostic Factors for Worse Outcomes 1 and 2 Years After Arthroscopic Partial Meniscectomy for Degenerative Meniscal Tears: A Secondary Explorative Study of the Surgically Treated Group From the Odense-Oslo Meniscectomy Versus Exercise (OMEX) Trial. <i>American Journal of Sports Medicine</i> , 2019 , 47, 2402-2411	6.8	9
73	Neuromuscular Exercise post Partial Medial Meniscectomy: Randomized Controlled Trial. <i>Medicine and Science in Sports and Exercise</i> , 2015 , 47, 1557-66	1.2	9
72	Cost-effectiveness of total knee replacement in addition to non-surgical treatment: a 2-year outcome from a randomised trial in secondary care in Denmark. <i>BMJ Open</i> , 2020 , 10, e033495	3	9
71	3 steps to improve reporting and interpretation of patient-reported outcome scores in orthopedic studies. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2018 , 89, 1-2	4.3	9
70	Factors associated with the orthopaedic surgeon's decision to recommend total joint replacement in hip and knee osteoarthritis: an international cross-sectional study of 1905 patients. <i>Osteoarthritis and Cartilage</i> , 2018 , 26, 1311-1318	6.2	8
69	The effects of neuromuscular exercise on medial knee joint load post-arthroscopic partial medial meniscectomy: SCOPEX, a randomised control trial protocol. <i>BMC Musculoskeletal Disorders</i> , 2012 , 13, 233	2.8	8
68	Non-surgical treatment of hip osteoarthritis. Hip school, with or without the addition of manual therapy, in comparison to a minimal control intervention: protocol for a three-armed randomized clinical trial. <i>BMC Musculoskeletal Disorders</i> , 2011 , 12, 88	2.8	8
67	Neuromuscular function during stair descent in meniscectomized patients and controls. <i>Medicine and Science in Sports and Exercise</i> , 2011 , 43, 1272-9	1.2	8
66	Bridging Disciplines as a pathway to Finding New Solutions for Osteoarthritis a collaborative program presented at the 2019 Orthopaedic Research Society and the Osteoarthritis Research Society International. <i>Osteoarthritis and Cartilage Open</i> , 2020 , 2, 100026	1.5	8
65	Disease modification in OA - will we ever get there?. <i>Nature Reviews Rheumatology</i> , 2019 , 15, 133-135	8.1	7

64	Effect of pre-operative neuromuscular training on functional outcome after total knee replacement: a randomized-controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2013 , 14, 157	2.8	7
63	An 8-Week Neuromuscular Exercise Program for Patients With Mild to Moderate Knee Osteoarthritis: A Case Series Drawn From a Registered Clinical Trial. <i>Journal of Athletic Training</i> , 2017 , 52, 592-605	4	7
62	Cross-Cultural and Construct Validity of the Animated Activity Questionnaire. <i>Arthritis Care and Research</i> , 2017 , 69, 1349-1359	4.7	7
61	Surgical or non-surgical treatment of traumatic skeletal fractures in adults: systematic review and meta-analysis of benefits and harms. <i>Systematic Reviews</i> , 2020 , 9, 179	3	7
60	Program evaluation of GLA:D [®] Australia: Physiotherapist training outcomes and effectiveness of implementation for people with knee osteoarthritis. <i>Osteoarthritis and Cartilage Open</i> , 2021 , 3, 100175	1.5	7
59	Association between thigh muscle strength four years after partial meniscectomy and radiographic features of osteoarthritis 11 years later. <i>BMC Musculoskeletal Disorders</i> , 2019 , 20, 512	2.8	6
58	IKDC or KOOS? Which measures symptoms and disabilities most important to postoperative articular cartilage repair patients?. <i>American Journal of Sports Medicine</i> , 2009 , 37, 1042-3; author reply 1043	6.8	6
57	Principles of brain plasticity in improving sensorimotor function of the knee and leg in healthy subjects: a double-blind randomized exploratory trial. <i>BMC Musculoskeletal Disorders</i> , 2009 , 10, 99	2.8	6
56	Physical Capacity in Female Soccer Players - Does Age Make a Difference?. <i>Advances in Physiotherapy</i> , 2000 , 2, 39-48		6
55	Immediate effect of valgus bracing on knee joint moments in meniscectomised patients: An exploratory study. <i>Journal of Science and Medicine in Sport</i> , 2016 , 19, 964-969	4.4	6
54	Impact of treatment strategy and physical performance on future knee-related self-efficacy in individuals with ACL injury. <i>BMC Musculoskeletal Disorders</i> , 2018 , 19, 50	2.8	5
53	The effect of education and supervised exercise vs. education alone on the time to total hip replacement in patients with severe hip osteoarthritis. A randomized clinical trial protocol. <i>BMC Musculoskeletal Disorders</i> , 2013 , 14, 21	2.8	5
52	Principles of brain plasticity in improving sensorimotor function of the knee and leg in patients with anterior cruciate ligament injury: a double-blind randomized exploratory trial. <i>BMC Musculoskeletal Disorders</i> , 2012 , 13, 68	2.8	5
51	Translation, Cross-Cultural Adaptation, and Validation of the Activity Rating Scale for Disorders of the Knee. <i>Orthopaedic Journal of Sports Medicine</i> , 2017 , 5, 2325967117729361	3.5	5
50	The Danish Hip Arthroscopy Registry: Registration Completeness and Patient Characteristics Between Responders and Non-Responders. <i>Clinical Epidemiology</i> , 2020 , 12, 825-833	5.9	5
49	The 6-m timed hop test is a prognostic factor for outcomes in patients with meniscal tears treated with exercise therapy or arthroscopic partial meniscectomy: a secondary, exploratory analysis of the Odense-Oslo meniscectomy versus exercise (OMEX) trial. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2019 , 27, 2478-2487	5.5	5
48	Mapping EQ-5D-3L from the Knee Injury and Osteoarthritis Outcome Score (KOOS). <i>Quality of Life Research</i> , 2020 , 29, 265-274	3.7	5
47	Prevalence, Severity, and Correlates of Pain Flares in Response to a Repeated Sit-to-Stand Activity: A Cross-sectional Study of 14 902 Patients With Knee and Hip Osteoarthritis in Primary Care. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2020 , 50, 309-318	4.2	5

46	Patients use fewer analgesics following supervised exercise therapy and patient education: an observational study of 16 499 patients with knee or hip osteoarthritis. <i>British Journal of Sports Medicine</i> , 2021 , 55, 670-675	10.3	5
45	Patients with osteoarthritis are least likely to receive lifestyle advice compared with patients with diabetes and hypertension: A national health survey study from Denmark. <i>Osteoarthritis and Cartilage Open</i> , 2020 , 2, 100067	1.5	4
44	Exercise-induced hypoalgesia in young adult females with long-standing patellofemoral pain - A randomized crossover study. <i>European Journal of Pain</i> , 2019 , 23, 1780-1789	3.7	4
43	Exploring the effect of space and place on response to exercise therapy for knee and hip pain--a protocol for a double-blind randomised controlled clinical trial: the CONEX trial. <i>BMJ Open</i> , 2015 , 5, e007701	2	4
42	Room for improvement: a randomised controlled trial with nested qualitative interviews on space, place and treatment delivery. <i>British Journal of Sports Medicine</i> , 2019 , 53, 359-367	10.3	4
41	Infographic. Therapeutic exercise relieves pain and does not harm knee cartilage nor trigger inflammation. <i>British Journal of Sports Medicine</i> , 2020 , 54, 118-119	10.3	3
40	Associations between comorbidities and immediate and one-year outcomes following supervised exercise therapy and patient education - A cohort study of 24,513 individuals with knee or hip osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2021 , 29, 39-49	6.2	3
39	Individualized predictions of changes in knee pain, quality of life and walking speed following patient education and exercise therapy in patients with knee osteoarthritis - a prognostic model study. <i>Osteoarthritis and Cartilage</i> , 2020 , 28, 1191-1201	6.2	2
38	Arthroscopic meniscectomy versus non-surgical or sham treatment in patients with MRI confirmed degenerative meniscus lesions: a protocol for an individual participant data meta-analysis. <i>BMJ Open</i> , 2020 , 10, e031864	3	2
37	Infographic. Roadmap to managing a person with musculoskeletal pain irrespective of body region. <i>British Journal of Sports Medicine</i> , 2020 , 54, 554-555	10.3	2
36	Percentile curves for the knee injury and osteoarthritis outcome score in the middle-aged Dutch population. <i>Osteoarthritis and Cartilage</i> , 2020 , 28, 1046-1054	6.2	2
35	Association between sensory function and hop performance and self-reported outcomes in patients with anterior cruciate ligament injury. <i>Open Access Journal of Sports Medicine</i> , 2017 , 8, 1-8	2.9	2
34	Infographic. Risk profile for sport-related post-traumatic knee osteoarthritis. <i>British Journal of Sports Medicine</i> , 2020 , 54, 362-363	10.3	2
33	Reliability, validity and responsiveness of a new leisure index: the Patient-Specific Leisure Scale (PSLS). <i>Musculoskeletal Care</i> , 2009 , 7, 178-93	1.6	2
32	Analgesic use in adolescents with patellofemoral pain or Osgood-Schlatter Disease: a secondary cross-sectional analysis of 323 subjects. <i>Scandinavian Journal of Pain</i> , 2021 ,	1.9	2
31	Prevalence of multimorbid degenerative lumbar spinal stenosis with knee and/or hip osteoarthritis: protocol for a systematic review and meta-analysis. <i>Systematic Reviews</i> , 2020 , 9, 232	3	2
30	On a Trajectory for Success-9 in Every 10 People With a Degenerative Meniscus Tear Have Improved Knee Function Within 2 Years After Treatment: A Secondary Exploratory Analysis of a Randomized Controlled Trial. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2021 , 51, 289-297	4.2	2
29	Knee osteoarthritis after meniscectomy: Prevalence of radiographic changes after twenty-one years, compared with matched controls 1998 , 41, 687		2

28	Zukunftsweisendes Arthrose-Management. <i>Der Schmerzpatient</i> , 2019 , 2, 23-28	0	1
27	Exercise Therapy as Treatment for Patients with Osteoarthritis of the Knee 2013 , 25-33		1
26	Efficacy of the SOAR knee health program: protocol for a two-arm stepped-wedge randomized delayed-controlled trial.. <i>BMC Musculoskeletal Disorders</i> , 2022 , 23, 85	2.8	1
25	Association between received treatment elements and satisfaction with care for patients with knee osteoarthritis seen in general practice in Denmark. <i>Scandinavian Journal of Primary Health Care</i> , 2021 , 39, 257-264	2.7	1
24	Assessing baseline dependency of anchor-based minimal important change (MIC): don't stratify on the baseline score!. <i>Quality of Life Research</i> , 2021 , 30, 2773-2782	3.7	1
23	Prognosis and transition of multi-site pain during the course of 5 years: Results of knee pain and function from a prospective cohort study among 756 adolescents. <i>PLoS ONE</i> , 2021 , 16, e0250415	3.7	1
22	Cutoff Values to Interpret Short-term Treatment Outcomes After Arthroscopic Meniscal Surgery, Measured With the Knee Injury and Osteoarthritis Outcome Score. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2021 , 51, 281-288	4.2	1
21	AuthorsPreply to Joshi. <i>BMJ, The</i> , 2016 , 354, i4623	5.9	1
20	Agreement Between Self-Reported Information and Administrative Data on Comorbidities, Imaging and Treatment in Denmark - A Validation Study of 38,745 Patients with Knee or Hip Osteoarthritis. <i>Clinical Epidemiology</i> , 2021 , 13, 779-790	5.9	1
19	Comparative responsiveness of measures of pain and function after total hip replacement 2001 , 45, 258		1
18	Proportion of patients with hip osteoarthritis in primary care identified by differing clinical criteria: a cross-sectional study of 4699 patients. <i>Osteoarthritis and Cartilage Open</i> , 2020 , 2, 100111	1.5	0
17	Comments on: validation study of an electronic method of condensed outcomes tools reporting in orthopaedics (J Knee Surg 2013;26:445-452). <i>Journal of Knee Surgery</i> , 2014 , 27, 165-6	2.4	0
16	Feasibility of the SOAR (Stop OsteoARthritis) program. <i>Osteoarthritis and Cartilage Open</i> , 2022 , 4, 100239.5	3.5	0
15	Care-seeking behaviour of adolescents with patellofemoral pain: a retrospective cohort study. <i>F1000Research</i> , 11, 161	3.6	0
14	Impact of educational level and employment status on short-term and long-term pain relief from supervised exercise therapy and education: an observational study of 22 588 patients with knee and hip osteoarthritis. <i>BMJ Open</i> , 2021 , 11, e045156	3	0
13	Prevalence of multimorbid degenerative lumbar spinal stenosis with knee or hip osteoarthritis: a systematic review and meta-analysis.. <i>BMC Musculoskeletal Disorders</i> , 2022 , 23, 177	2.8	0
12	Cost-effectiveness of an 8-week supervised education and exercise therapy programme for knee and hip osteoarthritis: a pre-post analysis of 16 255 patients participating in Good Life with osteoArthritis in Denmark (GLA:D).. <i>BMJ Open</i> , 2021 , 11, e049541	3	0
11	A more correct interpretation. <i>Cmaj</i> , 2015 , 187, 358	3.5	

10	Response to Letter to the Editor: "Osteoarthritis progression after exercise therapy or meniscectomy in patients with degenerative meniscal tears". <i>Osteoarthritis and Cartilage</i> , 2020 , 28, 1148-1149	6.2
9	Expert panels: can they be trusted?. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2013 , 29, 1127-8	5.4
8	Authors Preply to Chitnavis. <i>BMJ, The</i> , 2015 , 350, h431	5.9
7	Changes in medial-to-lateral knee joint loading patterns assessed by Dual-Energy X-ray Absorptiometry following supervised neuromuscular exercise therapy and patient education in patients with knee osteoarthritis: an exploratory cohort study.. <i>Physiotherapy Theory and Practice</i> , 2022 , 1-10	1.5
6	Are validated questionnaires valid?. <i>Journal of Bone and Joint Surgery - Series A</i> , 2006 , 88, 448	5.6
5	PROMs for Osteoarthritis 2016 , 249-285	
4	Single postoperative infusion of zoledronic acid to improve patient-reported outcome after hip or knee replacement: study protocol for a randomised, controlled, double-blinded clinical trial. <i>BMJ Open</i> , 2020 , 10, e040985	3
3	Response to Letter to Editor: 2-year outcome from two parallel randomized controlled trials. Reporting considerations? <i>Osteoarthritis and Cartilage</i> , 2019 , 27, e1-e2	6.2
2	Calculate Patient-Reported Scores as Recommended so as Not to Threaten the Validity of the Measure: Letter to the Editor. <i>Orthopaedic Journal of Sports Medicine</i> , 2018 , 6, 2325967118816987	3.5
1	Prognostic factors of total hip replacement during a 2-year period in participants enrolled in supervised education and exercise therapy: a prognostic study of 3657 participants with hip osteoarthritis. <i>Arthritis Research and Therapy</i> , 2021 , 23, 235	5.7