

Liane Maldaner

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2034711/publications.pdf>

Version: 2024-02-01

25
papers

479
citations

687363

13
h-index

713466

21
g-index

25
all docs

25
docs citations

25
times ranked

676
citing authors

#	ARTICLE	IF	CITATIONS
1	Rapid determination of L-ascorbic acid content in vitamin C serums by ultra-high-performance liquid chromatography-tandem mass spectrometry. <i>International Journal of Cosmetic Science</i> , 2022, 44, 131-141.	2.6	12
2	A 1/4-QuEChERS method combined with UHPLC-MS/MS for the analysis of phenolic compounds in red pepper varieties. <i>Journal of Food Composition and Analysis</i> , 2022, 112, 104647.	3.9	4
3	Piceatannol: um estilbeno natural com um espectro amplo de atividades biológicas. <i>Research, Society and Development</i> , 2022, 11, e49211932221.	0.1	1
4	Determination of phenolic acids and flavonoids from <i>Myrciaria cauliflora</i> edible part employing vortex-assisted matrix solid-phase dispersion (VA-MSPD) and UHPLC-MS/MS. <i>Journal of Food Composition and Analysis</i> , 2021, 95, 103667.	3.9	13
5	Evaluation of the QuEChERS method for the determination of phenolic compounds in yellow (<i>Brassica alba</i>), brown (<i>Brassica juncea</i>), and black (<i>Brassica nigra</i>) mustard seeds. <i>Food Chemistry</i> , 2021, 340, 128162.	8.2	24
6	Determination of n-3 fatty acids in shrimp using a mini-scale extraction method and GC-FID analysis. <i>Journal of the Iranian Chemical Society</i> , 2021, 18, 375-383.	2.2	1
7	Pharmacokinetics of amoxicillin in obese and nonobese subjects. <i>British Journal of Clinical Pharmacology</i> , 2021, 87, 3227-3233.	2.4	9
8	An improved analytical strategy based on the QuEChERS method for piceatannol analysis in seeds of <i>Passiflora</i> species. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2021, 44, 699-710.	1.0	1
9	Phenolic Compounds from <i>Butia odorata</i> (Barb. Rodr.) Noblick Fruit and Its Antioxidant and Antitumor Activities. <i>Food Analytical Methods</i> , 2020, 13, 61-68.	2.6	14
10	Evaluation of Dispersive Solid-Phase Extraction (d-SPE) as a Clean-up Step for Phenolic Compound Determination of <i>Myrciaria cauliflora</i> Peel. <i>Food Analytical Methods</i> , 2020, 13, 155-165.	2.6	19
11	Rapid extraction method followed by a d-SPE clean-up step for determination of phenolic composition and antioxidant and antiproliferative activities from berry fruits. <i>Food Chemistry</i> , 2020, 309, 125694.	8.2	20
12	Modified QuEChERS method for phenolic compounds determination in mustard greens (<i>Brassica</i>) <i>Tj ETQqO O O rgBT /Overlock 10 Tf 50</i>	4.9	16
13	Determination of acrylamide in brewed coffee by dispersive liquid-liquid microextraction (DLLME) and ultra-performance liquid chromatography tandem mass spectrometry (UPLC-MS/MS). <i>Food Chemistry</i> , 2019, 282, 120-126.	8.2	66
14	Determination of phenolic compounds and antioxidant activity in passion fruit pulp (<i>Passiflora</i> spp.) using a modified QuEChERS method and UHPLC-MS/MS. <i>LWT - Food Science and Technology</i> , 2019, 100, 397-403.	5.2	52
15	Antioxidant Activity and Determination of Phenolic Compounds from <i>Eugenia involucrata</i> DC. Fruits by UHPLC-MS/MS. <i>Food Analytical Methods</i> , 2017, 10, 2718-2728.	2.6	31
16	Determination of antioxidant activity and phenolic compounds of <i>Muntingia calabura</i> Linn. peel by HPLC-DAD and UPLC-ESI-MS/MS. <i>International Journal of Food Science and Technology</i> , 2017, 52, 954-963.	2.7	23
17	Evaluation of an alternative fluorinated sorbent for dispersive solid-phase extraction clean-up of the quick, easy, cheap, effective, rugged, and safe method for pesticide residues analysis. <i>Journal of Chromatography A</i> , 2017, 1514, 36-43.	3.7	36
18	Determination of some organic contaminants in water samples by solid-phase extraction and liquid chromatography-tandem mass spectrometry. <i>Talanta</i> , 2012, 100, 38-44.	5.5	50

#	ARTICLE	IF	CITATIONS
19	UHPLC Uma abordagem atual: desenvolvimentos e desafios recentes. <i>Scientia Chromatographica</i> , 2012, 4, 197-207.	0.2	8
20	Preparation and Characterization of a Microwave-Immobilized Fluorinated Stationary Phase for RP-LC. <i>Chromatographia</i> , 2010, 72, 617-626.	1.3	9
21	Fases estacionárias modernas para cromatografia líquida de alta eficiência em fase reversa. <i>Quimica Nova</i> , 2010, 33, 1559-1568.	0.3	13
22	O estado da arte da cromatografia líquida de ultra eficiência. <i>Quimica Nova</i> , 2009, 32, 214-222.	0.3	26
23	Poly(methyltetradecylsiloxane) immobilized onto silica for extraction of multiclass pesticides from surface waters. <i>Analytica Chimica Acta</i> , 2007, 582, 34-40.	5.4	18
24	Determination of Phenolic Compounds in Red Sweet Pepper (<i>Capsicum annuum</i> L.) using a Modified QuEChERS Method and UHPLC-MS/MS Analysis and Its Relation to Antioxidant Activity. <i>Journal of the Brazilian Chemical Society</i> , 0, , .	0.6	13
25	Phenolic Composition of <i>Dipteryx alata</i> Vogel Pulp + Peel and Its Antioxidant and Cytotoxic Properties. <i>Journal of the Brazilian Chemical Society</i> , 0, , .	0.6	0