

Alexander R Schmidt

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/203781/publications.pdf>

Version: 2024-02-01

98
papers

2,848
citations

172457

29
h-index

223800

46
g-index

102
all docs

102
docs citations

102
times ranked

1866
citing authors

#	ARTICLE	IF	CITATIONS
1	Miocene Ethiopian amber: A new source of fossil cryptogams. <i>Journal of Systematics and Evolution</i> , 2022, 60, 932-954.	3.1	11
2	Fossil evidence of lichen grazing from Paleogene amber. <i>Review of Palaeobotany and Palynology</i> , 2022, , 104664.	1.5	1
3	The mid-Miocene Zhangpu biota reveals an outstandingly rich rainforest biome in East Asia. <i>Science Advances</i> , 2021, 7, .	10.3	51
4	Checklist of fossil liverworts suitable for calibrating phylogenetic reconstructions. <i>Bryophyte Diversity and Evolution</i> , 2021, 43, .	1.1	12
5	Re-appraisal of two fossil Frullaniaceae species (Marchantiophyta, Porellales) from the mid-Cretaceous Burmese amber. <i>Cretaceous Research</i> , 2021, 124, 104803.	1.4	9
6	Conservation, preparation and imaging of diverse ambers and their inclusions. <i>Earth-Science Reviews</i> , 2021, 220, 103653.	9.1	32
7	Liverworts from Cretaceous amber. <i>Cretaceous Research</i> , 2021, 128, 104987.	1.4	9
8	Fossil <i>Usnea</i> and similar fruticose lichens from Palaeogene amber. <i>Lichenologist</i> , 2020, 52, 319-324.	0.8	4
9	Comment on the letter of the Society of Vertebrate Paleontology (SVP) dated April 21, 2020 regarding "Fossils from conflict zones and reproducibility of fossil-based scientific data" Myanmar amber. <i>Palaontologische Zeitschrift</i> , 2020, 94, 431-437.	1.6	28
10	Extinction and dawn of the modern world in the Carnian (Late Triassic). <i>Science Advances</i> , 2020, 6, .	10.3	116
11	Revealing the diversity of amber source plants from the Early Cretaceous Crato Formation, Brazil. <i>BMC Evolutionary Biology</i> , 2020, 20, 107.	3.2	17
12	<i>Selaginella</i> was hyperdiverse already in the Cretaceous. <i>New Phytologist</i> , 2020, 228, 1176-1182.	7.3	18
13	Crustose lichens with lichenicolous fungi from Paleogene amber. <i>Scientific Reports</i> , 2019, 9, 10360.	3.3	7
14	<i>Heinrichsia cheilanthoides</i> gen. et sp. nov., a fossil fern in the family Pteridaceae (Polypodiales) from the Cretaceous amber forests of Myanmar. <i>Journal of Systematics and Evolution</i> , 2019, 57, 329-338.	3.1	14
15	How diverse were ferns in the Baltic amber forest?. <i>Journal of Systematics and Evolution</i> , 2019, 57, 305-328.	3.1	13
16	Caspary's fungi from Baltic amber: historic specimens and new evidence. <i>Papers in Palaeontology</i> , 2019, 5, 365-389.	1.5	18
17	Morphological stasis in the first myxomycete from the Mesozoic, and the likely role of cryptobiosis. <i>Scientific Reports</i> , 2019, 9, 19730.	3.3	8
18	Amber inclusions from New Zealand. <i>Gondwana Research</i> , 2018, 56, 135-146.	6.0	31

#	ARTICLE	IF	CITATIONS
19	Fossil evidence of eupolypod ferns in the mid-Cretaceous of Myanmar. <i>Plant Systematics and Evolution</i> , 2018, 304, 1-13.	0.9	18
20	Production and preservation of resins – Past and present. <i>Biological Reviews</i> , 2018, 93, 1684-1714.	10.4	113
21	Morphological Convergence in Forest Microfungi Provides a Proxy for Paleogene Forest Structure. , 2018, , 527-549.		13
22	Calicioid lichens and fungi in amber – Tracing extant lineages back to the Paleogene. <i>Geobios</i> , 2018, 51, 469-479.	1.4	11
23	The Carnian Pluvial Episode and the first global appearance of amber. <i>Journal of the Geological Society</i> , 2018, 175, 1012-1018.	2.1	20
24	<i>Frullania grabenhorstii</i> sp. nov., a fossil liverwort (Jungermanniopsida: Frullaniaceae) with perianth from Bitterfeld amber. <i>Bryophyte Diversity and Evolution</i> , 2018, 40, 91.	1.1	9
25	Marine microorganisms as amber inclusions: insights from coastal forests of New Caledonia. <i>Fossil Record</i> , 2018, 21, 213-221.	1.4	11
26	Jochen Heinrichs March 14, 1969 – April 22, 2018. <i>Cryptogamie, Bryologie</i> , 2018, 39, 407-412.	0.2	0
27	The leafy liverwort genus <i>Lejeunea</i> (Porellales, Jungermanniopsida) in Miocene Dominican amber. <i>Review of Palaeobotany and Palynology</i> , 2017, 238, 144-150.	1.5	11
28	A fossil genus of the Frullaniaceae (Porellales, Jungermanniopsida) from the mid-Cretaceous of Myanmar. <i>Cretaceous Research</i> , 2017, 74, 223-226.	1.4	18
29	Diversity and ecological adaptations in Palaeogene lichens. <i>Nature Plants</i> , 2017, 3, 17049.	9.3	35
30	Resin exudation and resinicolous communities on <i>Araucaria humboldtensis</i> in New Caledonia. <i>Arthropod-Plant Interactions</i> , 2017, 11, 495-505.	1.1	8
31	Diverse early dwarf mistletoes (<i>Arceuthobium</i>), ecological keystones of the Eocene Baltic amber biota. <i>American Journal of Botany</i> , 2017, 104, 694-718.	1.7	17
32	Parasitaxus parasitized: novel infestation of <i>Parasitaxus usta</i> (Podocarpaceae). <i>Arthropod-Plant Interactions</i> , 2017, 11, 507-514.	1.1	0
33	The first fossil of Lindsaeaceae (Polypodiales) from the Cretaceous amber forest of Myanmar. <i>Cretaceous Research</i> , 2017, 72, 8-12.	1.4	24
34	A fossil species of the enigmatic early polypod fern genus <i>Cystodium</i> (Cystodiaceae) in Cretaceous amber from Myanmar. <i>Scientific Reports</i> , 2017, 7, 14615.	3.3	14
35	<i>Chaenothecopsis schefflerae</i> (Ascomycota: Mycocaliciales): a widespread fungus on semi-hardened exudates of endemic New Zealand Araliaceae. <i>New Zealand Journal of Botany</i> , 2017, 55, 387-406.	1.1	6
36	Problems related to the taxonomic placement of incompletely preserved amber fossils: transfer of the Paleogene liverwort <i>Cylindrocolea dimorpha</i> ; (Cephaloziellaceae) to the extant <i>Odontoschisma</i> sect. <i>lwatsukia</i> ; (Cephaloziaceae). <i>Fossil Record</i> , 2017, 20, 147-157.	1.4	6

#	ARTICLE	IF	CITATIONS
37	A Burmese amber fossil of <i>Radula</i> (Porellales, Jungermanniopsida) provides insights into the Cretaceous evolution of epiphytic lineages of leafy liverworts. <i>Fossil Record</i> , 2017, 20, 201-213.	1.4	18
38	Revision of the leafy liverwort genus <i>Radula</i> (Porellales, Jungermanniopsida) in Baltic and Bitterfeld amber. <i>Review of Palaeobotany and Palynology</i> , 2016, 235, 157-164.	1.5	12
39	A Caribbean epiphyte community preserved in Miocene Dominican amber. <i>Earth and Environmental Science Transactions of the Royal Society of Edinburgh</i> , 2016, 107, 321-331.	0.3	12
40	Diversity of lichen-associated filamentous fungi preserved in European Paleogene amber. <i>Earth and Environmental Science Transactions of the Royal Society of Edinburgh</i> , 2016, 107, 311-320.	0.3	8
41	<i>Sciadopitys</i> cladodes from Eocene Baltic amber. <i>Botanical Journal of the Linnean Society</i> , 2016, 180, 258-268.	1.6	24
42	<i>Resinogalea humboldtensis</i> gen. et sp. nov., a New Resiniculous Fungus from New Caledonia, Placed in <i>Bruceomycetaceae</i> (Ascomycota). <i>Annales Botanici Fennici</i> , 2016, 53, 205-215.	0.1	8
43	Burmese amber fossils bridge the gap in the Cretaceous record of polypod ferns. <i>Perspectives in Plant Ecology, Evolution and Systematics</i> , 2016, 18, 70-78.	2.7	40
44	Lichen-associated fungi from Paleogene amber. <i>New Phytologist</i> , 2016, 209, 896-898.	7.3	16
45	Graminids from Eocene Baltic amber. <i>Review of Palaeobotany and Palynology</i> , 2016, 233, 161-168.	1.5	14
46	Crown Group <i>Lejeuneaceae</i> and <i>Pleurocarpus</i> Mosses in Early Eocene (Ypresian) Indian Amber. <i>PLoS ONE</i> , 2016, 11, e0156301.	2.5	20
47	<i>Notoscyphus grollei</i> sp. nov. in Bitterfeld amber rather than the extant <i>Notoscyphus lutescens</i> (Lehm.) Tj ETQq1 1 0,784314 rgBT /Over	0.3	1
48	Transfer of the Eocene <i>Jungermannia berendtii</i> Grolle to <i>Solenostoma</i> . <i>Cryptogamie, Bryologie</i> , 2015, 36, 285-288.	0.2	6
49	Transfer of the Miocene <i>Lejeunea palaeomexicana</i> Grolle to <i>Ceratolejeunea</i> . <i>Cryptogamie, Bryologie</i> , 2015, 36, 335-341.	0.2	9
50	Plant-feeding mite diversity in Triassic amber (Acari: Tetrapodili). <i>Journal of Systematic Palaeontology</i> , 2015, 13, 129-151.	1.5	49
51	A new Dominican amber fossil of the derived fern genus <i>Pleopeltis</i> confirms generic stasis in the epiphytic fern diversity of the West Indies. <i>Organisms Diversity and Evolution</i> , 2015, 15, 277-283.	1.6	22
52	The enigmatic hyphomycete <i>Torula</i> sensu Caspary revisited. <i>Review of Palaeobotany and Palynology</i> , 2015, 219, 183-193.	1.5	12
53	The leafy liverwort <i>Notoscyphus balticus</i> sp. nov. (Jungermanniales) in Eocene Baltic amber. <i>Review of Palaeobotany and Palynology</i> , 2015, 217, 39-44.	1.5	15
54	<i>Lejeuneaceae</i> (Marchantiophyta) from a species-rich taphocoenosis in Miocene Mexican amber, with a review of liverworts fossilised in amber. <i>Review of Palaeobotany and Palynology</i> , 2015, 221, 59-70.	1.5	36

#	ARTICLE	IF	CITATIONS
55	Carnivorous leaves from Baltic amber. Proceedings of the National Academy of Sciences of the United States of America, 2015, 112, 190-195.	7.1	22
56	The intriguing marine diatom genus <i>Corethron</i> in Late Cretaceous amber from Vendôme (France). Cretaceous Research, 2015, 52, 64-72.	1.4	14
57	Alectoroid Morphologies in Paleogene Lichens: New Evidence and Re-Evaluation of the Fossil <i>Alectoria succini</i> Mägdefrau. PLoS ONE, 2015, 10, e0129526.	2.5	36
58	Molecular and Morphological Evidence Challenges the Records of the Extant Liverwort <i>Ptilidium pulcherrimum</i> in Eocene Baltic Amber. PLoS ONE, 2015, 10, e0140977.	2.5	17
59	Species-level determination of closely related araucarian resins using FTIR spectroscopy and its implications for the provenance of New Zealand amber. PeerJ, 2015, 3, e1067.	2.0	23
60	The first fossil of a bolbitidoid fern belongs to the early-divergent lineages of <i>Elaphoglossum</i> (Dryopteridaceae). American Journal of Botany, 2014, 101, 1466-1475.	1.7	31
61	The extant liverwort <i>Gackstroemia</i> (Lepidolaenaceae, Porellales) in Cretaceous amber from Myanmar. Review of Palaeobotany and Palynology, 2014, 203, 48-52.	1.5	30
62	Amber fossils of sooty moulds. Review of Palaeobotany and Palynology, 2014, 200, 53-64.	1.5	37
63	An acrocarpous moss in Cretaceous amber from Myanmar. Cretaceous Research, 2014, 51, 260-265.	1.4	13
64	A fossil species of <i>Ceratolejeunea</i> (Lejeuneaceae, Porellales) preserved in Miocene Mexican amber. Bryologist, 2014, 117, 10-14.	0.6	20
65	Bryophytes of the Burmese amber forest: Amending and expanding the circumscription of the Cretaceous moss genus <i>Vetiplanaxis</i> . Review of Palaeobotany and Palynology, 2014, 209, 1-10.	1.5	26
66	Estimating the Phanerozoic history of the Ascomycota lineages: Combining fossil and molecular data. Molecular Phylogenetics and Evolution, 2014, 78, 386-398.	2.7	197
67	The Bromeliaceae tank dweller <i>Bromeliophila</i> (Lejeuneaceae, Porellales) is a member of the <i>Cyclolejeunea-Prionolejeunea</i> clade. Plant Systematics and Evolution, 2014, 300, 63-73.	0.9	14
68	<p align="left">Chaenothecopsis neocaledonica sp. nov.: The first resinicolous mycocalicioid fungus from an araucarian conifer. Phytotaxa, 2014, 173, 49.	0.3	14
69	Epiphytic leafy liverworts diversified in angiosperm-dominated forests. Scientific Reports, 2014, 4, 5974.	3.3	104
70	Exploring the impact of fossil constraints on the divergence time estimates of derived liverworts. Plant Systematics and Evolution, 2013, 299, 585-601.	0.9	38
71	Stuck in time – a new <i>Chaenothecopsis</i> species with proliferating ascomata from <i>Cunninghamia</i> resin and its fossil ancestors in European amber. Fungal Diversity, 2013, 58, 199-213.	12.3	29
72	Molecular Phylogeny of the Leafy Liverwort <i>Lejeunea</i> (Porellales): Evidence for a Neotropical Origin, Uneven Distribution of Sexual Systems and Insufficient Taxonomy. PLoS ONE, 2013, 8, e82547.	2.5	53

#	ARTICLE	IF	CITATIONS
73	Arthropods in amber from the Triassic Period. Proceedings of the National Academy of Sciences of the United States of America, 2012, 109, 14796-14801.	7.1	132
74	Tramps, narrow endemics and morphologically cryptic species in the epiphyllous liverwort <i>Diplasiolejeunea</i> . Molecular Phylogenetics and Evolution, 2012, 65, 582-594.	2.7	59
75	The leafy liverwort <i>Frullania</i> (Jungermanniopsida) in the Cretaceous amber forest of Myanmar. Review of Palaeobotany and Palynology, 2012, 169, 21-28.	1.5	39
76	Ectomycorrhizas from a Lower Eocene angiosperm forest. New Phytologist, 2011, 192, 988-996.	7.3	47
77	<i>Kaolakia borealis</i> nov. gen. et sp. (Porellales, Jungermanniopsida): A leafy liverwort from the Cretaceous of Alaska. Review of Palaeobotany and Palynology, 2011, 165, 235-240.	1.5	28
78	Microbes in Resinous Habitats: A Compilation from Modern and Fossil Resins. Lecture Notes in Earth Sciences, 2011, , 391-407.	0.5	17
79	One species or at least eight? Delimitation and distribution of <i>Frullania tamarisci</i> (L.) Dumort. s. l. (Jungermanniopsida, Porellales) inferred from nuclear and chloroplast DNA markers. Molecular Phylogenetics and Evolution, 2010, 56, 1105-1114.	2.7	99
80	Testate Amoebae from a Cretaceous Forest Floor Microbiocoenosis of France. Journal of Eukaryotic Microbiology, 2010, 57, 245-248.	1.7	23
81	Cretaceous African life captured in amber. Proceedings of the National Academy of Sciences of the United States of America, 2010, 107, 7329-7334.	7.1	85
82	Exceptional preservation of marine diatoms in upper Albian amber. Geology, 2009, 37, 83-86.	4.4	37
83	Evidence for marine microfossils from amber. Proceedings of the National Academy of Sciences of the United States of America, 2008, 105, 17426-17429.	7.1	60
84	<i>Palaeoanellus dimorphus</i> gen. et sp. nov. (Deuteromycotina): a Cretaceous predatory fungus. American Journal of Botany, 2008, 95, 1328-1334.	1.7	28
85	Aquatic organisms as amber inclusions and examples from a modern swamp forest. Proceedings of the National Academy of Sciences of the United States of America, 2007, 104, 16581-16585.	7.1	55
86	Evidence of Cenozoic Matoniaceae from Baltic and Bitterfeld amber. Review of Palaeobotany and Palynology, 2007, 144, 145-156.	1.5	20
87	AN ARCHAIC SLIME MOULD IN BALTIC AMBER. Palaeontology, 2006, 49, 1013-1017.	2.2	9
88	A microworld in Triassic amber. Nature, 2006, 444, 835-835.	27.8	88
89	LEPTOTRICHITES RESINATUS NEW GENUS AND SPECIES: A FOSSIL SHEATHED BACTERIUM IN ALPINE CRETACEOUS AMBER. Journal of Paleontology, 2005, 79, 175-184.	0.8	39
90	Diverse fossil amoebae in German Mesozoic amber. Palaeontology, 2004, 47, 185-197.	2.2	47

#	ARTICLE	IF	CITATIONS
91	The oldest fossil myxogastroid slime mould. <i>Mycological Research</i> , 2003, 107, 123-126.	2.5	22
92	Sooty moulds from European Tertiary amber, with notes on the systematic position of <i>Rosaria</i> (â€Cyanobacteriaâ€™). <i>Mycological Research</i> , 2003, 107, 251-256.	2.5	36
93	The Mesozoic amber of Schliersee (southern Germany) is Cretaceous in age. <i>Cretaceous Research</i> , 2001, 22, 423-428.	1.4	41
94	A Fossil <i>Scapania</i> (Hepaticae) with Perianth and Capsule in Bitterfeld Amber (Eocene) from Germany. <i>Bryologist</i> , 2001, 104, 362-366.	0.6	11
95	<i>Rosaria succina</i> spec. nov. - a fossil cyanobacterium from Tertiary amber. <i>Journal of Basic Microbiology</i> , 2000, 40, 327-332.	3.3	7
96	A conifer seedling with two herbicolous fungi from the Baltic amber forest. <i>Botanical Journal of the Linnean Society</i> , 0, 155, 449-456.	1.6	11
97	Uncovering the natural variability of araucariacean exudates from <i>ex situ</i> and <i>in situ</i> tree populations in New Caledonia using FTIR spectroscopy. , 0, 4, e17.		2
98	An Exceptionally Preserved Terrestrial Record of LIP Effects on Plants in the Carnian (Upper Triassic) Amber-Bearing Section of the Dolomites, Italy. <i>Frontiers in Earth Science</i> , 0, 10, .	1.8	7