Yuji Mizuno

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2038765/publications.pdf

Version: 2024-02-01

304743 434195 2,691 34 22 31 citations h-index g-index papers 34 34 34 2478 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Î ² -Blockers are associated with increased B-type natriuretic peptide levels differently in men and women in heart failure with preserved ejection fraction. American Journal of Physiology - Heart and Circulatory Physiology, 2022, 323, H276-H284.	3.2	6
2	B-Type Natriuretic Peptide in Heart Failure With Preserved Ejection Fraction ― Relevance to Age-Related Left Ventricular Modeling in Japanese ―. Circulation Journal, 2017, 81, 1006-1013.	1.6	20
3	Differences and Interactions between Risk Factors for Coronary Spasm and Atherosclerosis -Smoking, Aging, Inflammation, and Blood Pressure Internal Medicine, 2014, 53, 2663-2670.	0.7	24
4	Coronary spastic angina is associated with insulin resistance – possible involvement of endothelial dysfunction. Coronary Artery Disease, 2013, 24, 559-565.	0.7	5
5	Coronary Spasm Preferentially Occurs at Branch Points. Circulation: Cardiovascular Interventions, 2009, 2, 97-104.	3.9	39
6	Coronary artery spasmâ€"Clinical features, diagnosis, pathogenesis, and treatment. Journal of Cardiology, 2008, 51, 2-17.	1.9	330
7	Effects of a 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase Inhibitor, Fluvastatin, on Coronary Spasm After Withdrawal of Calcium-Channel Blockers. Journal of the American College of Cardiology, 2008, 51, 1742-1748.	2.8	163
8	Coronary Spasm is Associated With Chronic Low-Grade Inflammation. Circulation Journal, 2007, 71, 1074-1078.	1.6	70
9	Decreases in serum thioredoxin and hyperglycemia by acidity water footbath after lunch in type II diabetics. Journal of Molecular and Cellular Cardiology, 2007, 42, S58-S59.	1.9	O
10	Cardiac Pituitary-adrenal Axis in the Hypertensive Human Heart. Journal of Cardiac Failure, 2006, 12, S149.	1.7	0
11	Low Plasma Levels of Natriuretic Peptides in Obesity. Journal of Cardiac Failure, 2006, 12, S170.	1.7	O
12	Low-Grade Inflammation, Thrombogenicity, and Atherogenic Lipid Profile in Cigarette Smokers. Circulation Journal, 2006, 70, 8-13.	1.6	97
13	Plasma Level of B-Type Natriuretic Peptide as a Prognostic Marker After Acute Myocardial Infarction. Circulation, 2004, 110, 1387-1391.	1.6	134
14	Aldosterone Is Produced From Ventricles in Patients With Essential Hypertension. Hypertension, 2002, 39, 958-962.	2.7	72
15	B-type natriuretic peptide as a marker of the effects of enalapril in patients with heart failure. American Journal of Medicine, 2002, 112, 716-720.	1.5	65
16	B-type natriuretic peptide after percutaneous transluminal septal myocardial ablation. International Journal of Cardiology, 2002, 83, 151-158.	1.7	5
17	Effects of perindopril on aldosterone production in the failing human heart. American Journal of Cardiology, 2002, 89, 1197-1200.	1.6	16
18	Paraoxonase gene Gln192Arg (Q192R) polymorphism is associated with coronary artery spasm. Human Genetics, 2002, 110, 89-94.	3.8	74

#	Article	IF	CITATIONS
19	Aldosterone Induces Angiotensin-Converting-Enzyme Gene Expression in Cultured Neonatal Rat Cardiocytes. Circulation, 2001, 104, 137-139.	1.6	202
20	Aldosterone Production Is Activated in Failing Ventricle in Humans. Circulation, 2001, 103, 72-77.	1.6	327
21	A T-786→C mutation in the 5′-flanking region of the endothelial nitric oxide synthase gene and coronary arterial vasomotility. American Journal of Cardiology, 2000, 85, 710-714.	1.6	42
22	Tâ^'786â†'C mutation in the 5′-flanking region of the endothelial nitric oxide synthase gene is associated with myocardial infarction, especially without coronary organic stenosis. American Journal of Cardiology, 2000, 86, 628-634.	1.6	139
23	Plasma levels of A- and B-type natriuretic peptides in patients with hypertrophic cardiomyopathy or idiopathic dilated cardiomyopathy. American Journal of Cardiology, 2000, 86, 1036-1040.	1.6	70
24	The Plasma Levels of Dehydroepiandrosterone Sulfate Are Decreased in Patients with Chronic Heart Failure in Proportion to the Severity*. Journal of Clinical Endocrinology and Metabolism, 2000, 85, 1834-1840.	3.6	85
25	The effect of heparin on tissue factor and tissue factor pathway inhibitor in patients with acute myocardial infarction. International Journal of Cardiology, 2000, 75, 267-274.	1.7	10
26	Comparison of the risk factors for coronary artery spasm with those for organic stenosis in a Japanese population: role of cigarette smoking. International Journal of Cardiology, 2000, 72, 121-126.	1.7	99
27	Interaction on metabolic clearance between A-type and B-type natriuretic peptides in patients with heart failure. Metabolism: Clinical and Experimental, 2000, 49, 1228-1233.	3.4	12
28	Remnant Lipoprotein Levels in Fasting Serum Predict Coronary Events in Patients With Coronary Artery Disease. Circulation, 1999, 99, 2858-2860.	1.6	270
29	Cardiac angiotensin-converting enzyme activity in myocardial infarction. American Journal of Cardiology, 1999, 84, 774-778.	1.6	12
30	Effect of Interleukin- $1\hat{l}^2$ on Cardiac Hypertrophy and Production of Natriuretic Peptides in Rat Cardiocyte Culture. Journal of Molecular and Cellular Cardiology, 1999, 31, 1997-2006.	1.9	74
31	Effects of angiotensin-converting enzyme inhibitor on plasma b-type natriuretic peptide levels in patients with acute myocardial infarction. Journal of Cardiac Failure, 1997, 3, 287-293.	1.7	16
32	Increased plasma tissue factor levels in acute myocardial infarction. American Heart Journal, 1997, 134, 253-259.	2.7	146
33	The effects of the angiotensin-converting enzyme inhibitor imidapril on plasma plasminogen activator inhibitor activity in patients with acute myocardial infarction. American Heart Journal, 1997, 134, 961-966.	2.7	43
34	A Case of 5-Fluorouracil Cardiotoxicity Simulating Acute Myocardial Infarction Japanese Circulation Journal, 1995, 59, 303-307.	1.0	24