

Paul Dupree

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/2045776/paul-dupree-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

177
papers

14,747
citations

70
h-index

119
g-index

205
ext. papers

17,046
ext. citations

7.4
avg, IF

6.23
L-index

#	Paper	IF	Citations
177	Importance of Water in Maintaining Softwood Secondary Cell Wall Nanostructure. <i>Biomacromolecules</i> , 2021 , 22, 4669-4680	6.9	7
176	Characterisation of the enzyme transport path between shipworms and their bacterial symbionts. <i>BMC Biology</i> , 2021 , 19, 233	7.3	1
175	The nucleotide sugar transporter GONST2 is a functional homolog of GONST1. <i>Plant Direct</i> , 2021 , 5, e00309	3.9	2
174	Xylan Structure and Dynamics in Native Grass Cell Walls Investigated by Solid-State NMR Spectroscopy. <i>ACS Omega</i> , 2021 , 6, 15460-15471	3.9	5
173	Unlocking the structural features for the xylobiohydrolase activity of an unusual GH11 member identified in a compost-derived consortium. <i>Biotechnology and Bioengineering</i> , 2021 , 118, 4052-4064	4.9	2
172	Plant cell wall architecture guided design of CBM3-GH11 chimeras with enhanced xylanase activity using a tandem repeat left-handed β -prism scaffold. <i>Computational and Structural Biotechnology Journal</i> , 2021 , 19, 1108-1118	6.8	4
171	Spontaneous rearrangement of acetylated xylan on hydrophilic cellulose surfaces. <i>Cellulose</i> , 2021 , 28, 3327-3345	5.5	5
170	Two conifer GUX clades are responsible for distinct glucuronic acid patterns on xylan. <i>New Phytologist</i> , 2021 , 231, 1720-1733	9.8	2
169	Galactoglucomannan structure of Arabidopsis seed-coat mucilage in GDP-mannose synthesis impaired mutants. <i>Physiologia Plantarum</i> , 2021 , 173, 1244-1252	4.6	0
168	Loss of TaIRX9b gene function in wheat decreases chain length and amount of arabinoxylan in grain but increases cross-linking. <i>Plant Biotechnology Journal</i> , 2020 , 18, 2316-2327	11.6	7
167	Cell wall remodeling under salt stress: Insights into changes in polysaccharides, feruloylation, lignification, and phenolic metabolism in maize. <i>Plant, Cell and Environment</i> , 2020 , 43, 2172-2191	8.4	25
166	Three Decades of Advances in Arabinogalactan-Protein Biosynthesis. <i>Frontiers in Plant Science</i> , 2020 , 11, 610377	6.2	18
165	Carbohydrate Gel Electrophoresis. <i>Methods in Molecular Biology</i> , 2020 , 2149, 33-44	1.4	
164	BdGT43B2 functions in xylan biosynthesis and is essential for seedling survival in. <i>Plant Direct</i> , 2020 , 4, e00216	3.3	5
163	Calcium Binding by Arabinogalactan Polysaccharides Is Important for Normal Plant Development. <i>Plant Cell</i> , 2020 , 32, 3346-3369	11.6	27
162	Oligosaccharide Binding and Thermostability of Two Related AA9 Lytic Polysaccharide Monoxygenases. <i>Biochemistry</i> , 2020 , 59, 3347-3358	3.2	11
161	Vascular Plants Are Globally Significant Contributors to Marine Carbon Fluxes and Sinks. <i>Annual Review of Marine Science</i> , 2020 , 12, 469-497	15.4	31

160	Correlative FLIM-confocal-Raman mapping applied to plant lignin composition and autofluorescence. <i>Micron</i> , 2019 , 126, 102733	2.3	5
159	Development of an oligosaccharide library to characterise the structural variation in glucuronoarabinoxylan in the cell walls of vegetative tissues in grasses. <i>Biotechnology for Biofuels</i> , 2019 , 12, 109	7.8	9
158	An engineered GH1 β -glucosidase displays enhanced glucose tolerance and increased sugar release from lignocellulosic materials. <i>Scientific Reports</i> , 2019 , 9, 4903	4.9	24
157	Two members of the DUF579 family are responsible for arabinogalactan methylation in <i>Arabidopsis</i> . <i>Plant Direct</i> , 2019 , 3, e00117	3.3	14
156	Molecular architecture of softwood revealed by solid-state NMR. <i>Nature Communications</i> , 2019 , 10, 4978	7.4	87
155	Structural Imaging of Native Cryo-Preserved Secondary Cell Walls Reveals the Presence of Macrofibrils and Their Formation Requires Normal Cellulose, Lignin and Xylan Biosynthesis. <i>Frontiers in Plant Science</i> , 2019 , 10, 1398	6.2	19
154	Covalent interactions between lignin and hemicelluloses in plant secondary cell walls. <i>Current Opinion in Biotechnology</i> , 2019 , 56, 97-104	11.4	103
153	Water deficit and abscisic acid treatments increase the expression of a glucomannan mannosyltransferase gene (GMMT) in <i>Aloe vera</i> Burm. F. <i>Phytochemistry</i> , 2019 , 159, 90-101	4	5
152	An ancient family of lytic polysaccharide monooxygenases with roles in arthropod development and biomass digestion. <i>Nature Communications</i> , 2018 , 9, 756	17.4	135
151	Identification of an algal xylan synthase indicates that there is functional orthology between algal and plant cell wall biosynthesis. <i>New Phytologist</i> , 2018 , 218, 1049-1060	9.8	35
150	A Transcriptomic Analysis of Xylan Mutants Does Not Support the Existence of a Secondary Cell Wall Integrity System in <i>Arabidopsis</i> . <i>Frontiers in Plant Science</i> , 2018 , 9, 384	6.2	15
149	Characterization of a β -galactosidase from <i>Bacillus subtilis</i> with transgalactosylation activity. <i>International Journal of Biological Macromolecules</i> , 2018 , 120, 279-287	7.9	15
148	Glycosyltransferases of the GT43 Family 2018 , 251-263		
147	Hemocyanin facilitates lignocellulose digestion by wood-boring marine crustaceans. <i>Nature Communications</i> , 2018 , 9, 5125	17.4	16
146	The Patterned Structure of Galactoglucomannan Suggests It May Bind to Cellulose in Seed Mucilage. <i>Plant Physiology</i> , 2018 , 178, 1011-1026	6.6	30
145	A surface endogalactanase in <i>Bacteroides thetaiotaomicron</i> confers keystone status for arabinogalactan degradation. <i>Nature Microbiology</i> , 2018 , 3, 1314-1326	26.6	57
144	UUAT1 Is a Golgi-Localized UDP-Uronic Acid Transporter That Modulates the Polysaccharide Composition of <i>Arabidopsis</i> Seed Mucilage. <i>Plant Cell</i> , 2017 , 29, 129-143	11.6	36
143	Effects of Xylan Side-Chain Substitutions on Xylan-Cellulose Interactions and Implications for Thermal Pretreatment of Cellulosic Biomass. <i>Biomacromolecules</i> , 2017 , 18, 1311-1321	6.9	46

142	An even pattern of xylan substitution is critical for interaction with cellulose in plant cell walls. <i>Nature Plants</i> , 2017 , 3, 859-865	11.5	114
141	Structural and electronic determinants of lytic polysaccharide monooxygenase reactivity on polysaccharide substrates. <i>Nature Communications</i> , 2017 , 8, 1064	17.4	101
140	The wood from the trees: The use of timber in construction. <i>Renewable and Sustainable Energy Reviews</i> , 2017 , 68, 333-359	16.2	424
139	Removal of glucuronic acid from xylan is a strategy to improve the conversion of plant biomass to sugars for bioenergy. <i>Biotechnology for Biofuels</i> , 2017 , 10, 224	7.8	33
138	Evolution of Xylan Substitution Patterns in Gymnosperms and Angiosperms: Implications for Xylan Interaction with Cellulose. <i>Plant Physiology</i> , 2016 , 171, 2418-31	6.6	89
137	Golgi-localized STELLO proteins regulate the assembly and trafficking of cellulose synthase complexes in Arabidopsis. <i>Nature Communications</i> , 2016 , 7, 11656	17.4	62
136	An efficient arabinoxylan-debranching L-arabinofuranosidase of family GH62 from <i>Aspergillus nidulans</i> contains a secondary carbohydrate binding site. <i>Applied Microbiology and Biotechnology</i> , 2016 , 100, 6265-6277	5.7	18
135	The molecular basis of polysaccharide cleavage by lytic polysaccharide monooxygenases. <i>Nature Chemical Biology</i> , 2016 , 12, 298-303	11.7	205
134	Structural Modifications of Fructans in <i>Aloe barbadensis</i> Miller (Aloe Vera) Grown under Water Stress. <i>PLoS ONE</i> , 2016 , 11, e0159819	3.7	28
133	Folding of xylan onto cellulose fibrils in plant cell walls revealed by solid-state NMR. <i>Nature Communications</i> , 2016 , 7, 13902	17.4	207
132	Xylan decoration patterns and the plant secondary cell wall molecular architecture. <i>Biochemical Society Transactions</i> , 2016 , 44, 74-8	5.1	46
131	Structure and boosting activity of a starch-degrading lytic polysaccharide monooxygenase. <i>Nature Communications</i> , 2015 , 6, 5961	17.4	205
130	L-Fucose-containing arabinogalactan-protein in radish leaves. <i>Carbohydrate Research</i> , 2015 , 415, 1-11	2.9	19
129	Glycan complexity dictates microbial resource allocation in the large intestine. <i>Nature Communications</i> , 2015 , 6, 7481	17.4	241
128	Probing the molecular architecture of <i>Arabidopsis thaliana</i> secondary cell walls using two- and three-dimensional ¹³ C solid state nuclear magnetic resonance spectroscopy. <i>Biochemistry</i> , 2015 , 54, 2335-45	3.2	54
127	Monoclonal antibodies indicate low-abundance links between heteroxylan and other glycans of plant cell walls. <i>Planta</i> , 2015 , 242, 1321-34	4.7	43
126	Aspen Tension Wood Fibers Contain $\beta(1\rightarrow 4)$ -Galactans and Acidic Arabinogalactans Retained by Cellulose Microfibrils in Gelatinous Walls. <i>Plant Physiology</i> , 2015 , 169, 2048-63	6.6	54
125	Suppression of xylan endotransglycosylase PtXyn10A affects cellulose microfibril angle in secondary wall in aspen wood. <i>New Phytologist</i> , 2015 , 205, 666-81	9.8	44

124	An unusual xylan in Arabidopsis primary cell walls is synthesised by GUX3, IRX9L, IRX10L and IRX14. <i>Plant Journal</i> , 2015 , 83, 413-26	6.9	57
123	Action of an endo- β 1,3(4)-glucanase on cellobiosyl unit structure in barley β 1,3:1,4-glucan. <i>Bioscience, Biotechnology and Biochemistry</i> , 2015 , 79, 1810-7	2.1	8
122	KONJAC1 and 2 Are Key Factors for GDP-Mannose Generation and Affect l-Ascorbic Acid and Glucomannan Biosynthesis in Arabidopsis. <i>Plant Cell</i> , 2015 , 27, 3397-409	11.6	30
121	Lignocellulose degradation mechanisms across the Tree of Life. <i>Current Opinion in Chemical Biology</i> , 2015 , 29, 108-19	9.7	317
120	Secondary cell wall composition and candidate gene expression in developing willow (<i>Salix purpurea</i>) stems. <i>Planta</i> , 2014 , 239, 1041-53	4.7	7
119	The role of carbon starvation in the induction of enzymes that degrade plant-derived carbohydrates in <i>Aspergillus niger</i> . <i>Fungal Genetics and Biology</i> , 2014 , 72, 34-47	3.9	62
118	Identification of a sphingolipid β glucuronosyltransferase that is essential for pollen function in Arabidopsis. <i>Plant Cell</i> , 2014 , 26, 3314-25	11.6	59
117	Label-free protein quantification for plant Golgi protein localization and abundance. <i>Plant Physiology</i> , 2014 , 166, 1033-43	6.6	31
116	Evidence that GH115 β glucuronidase activity, which is required to degrade plant biomass, is dependent on conformational flexibility. <i>Journal of Biological Chemistry</i> , 2014 , 289, 53-64	5.4	46
115	The Golgi localized bifunctional UDP-rhamnose/UDP-galactose transporter family of Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 11563-8	11.5	81
114	Enzymatic fragmentation of carbohydrate moieties of radish arabinogalactan-protein and elucidation of the structures. <i>Bioscience, Biotechnology and Biochemistry</i> , 2014 , 78, 818-31	2.1	19
113	The pattern of xylan acetylation suggests xylan may interact with cellulose microfibrils as a twofold helical screw in the secondary plant cell wall of Arabidopsis thaliana. <i>Plant Journal</i> , 2014 , 79, 492-506	6.9	181
112	Characterisation of FUT4 and FUT6 (β 1 \rightarrow 2)-fucosyltransferases reveals that absence of root arabinogalactan fucosylation increases Arabidopsis root growth salt sensitivity. <i>PLoS ONE</i> , 2014 , 9, e93291	3.7	46
111	Lignin biosynthesis perturbations affect secondary cell wall composition and saccharification yield in Arabidopsis thaliana. <i>Biotechnology for Biofuels</i> , 2013 , 6, 46	7.8	194
110	Secondary cell wall characterization in a BY-2 inductive system. <i>Plant Cell, Tissue and Organ Culture</i> , 2013 , 115, 223-232	2.7	10
109	Identification of an additional protein involved in mannan biosynthesis. <i>Plant Journal</i> , 2013 , 73, 105-17	6.9	33
108	A proteomic approach identifies many novel palmitoylated proteins in Arabidopsis. <i>New Phytologist</i> , 2013 , 197, 805-814	9.8	104
107	Development and application of a high throughput carbohydrate profiling technique for analyzing plant cell wall polysaccharides and carbohydrate active enzymes. <i>Biotechnology for Biofuels</i> , 2013 , 6, 94	7.8	30

106	A galactosyltransferase acting on arabinogalactan protein glycans is essential for embryo development in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2013 , 76, 128-37	6.9	64
105	Abnormal glycosphingolipid mannosylation triggers salicylic acid-mediated responses in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2013 , 25, 1881-94	11.6	74
104	A β -glucuronosyltransferase from <i>Arabidopsis thaliana</i> involved in biosynthesis of type III arabinogalactan has a role in cell elongation during seedling growth. <i>Plant Journal</i> , 2013 , 76, 1016-29	6.9	60
103	Palmitoylation in plants: new insights through proteomics. <i>Plant Signaling and Behavior</i> , 2013 , 8,	2.5	12
102	β -galactosyl Yariv reagent binds to the β 1,3-galactan of arabinogalactan proteins. <i>Plant Physiology</i> , 2013 , 161, 1117-26	6.6	101
101	GUX1 and GUX2 glucuronyltransferases decorate distinct domains of glucuronoxylan with different substitution patterns. <i>Plant Journal</i> , 2013 , 74, 423-34	6.9	124
100	Studies of enzymatic cleavage of cellulose using polysaccharide analysis by carbohydrate gel electrophoresis (PACE). <i>Methods in Enzymology</i> , 2012 , 510, 51-67	1.7	8
99	Structural characterization of <i>Arabidopsis</i> leaf arabinogalactan polysaccharides. <i>Plant Physiology</i> , 2012 , 160, 653-66	6.6	93
98	Putative glycosyltransferases and other plant Golgi apparatus proteins are revealed by LOPIT proteomics. <i>Plant Physiology</i> , 2012 , 160, 1037-51	6.6	119
97	Glycosyl transferases in family 61 mediate arabinofuranosyl transfer onto xylan in grasses. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 989-93	11.5	185
96	<i>Arabidopsis</i> genes IRREGULAR XYLEM (IRX15) and IRX15L encode DUF579-containing proteins that are essential for normal xylan deposition in the secondary cell wall. <i>Plant Journal</i> , 2011 , 66, 401-13	6.9	110
95	Carbohydrate gel electrophoresis. <i>Methods in Molecular Biology</i> , 2011 , 715, 81-92	1.4	8
94	Determination of the N-glycosylation patterns of seed proteins: applications to determine the authenticity and substantial equivalence of genetically modified (GM) crops. <i>Journal of Agricultural and Food Chemistry</i> , 2011 , 59, 8779-88	5.7	12
93	Insights into the oxidative degradation of cellulose by a copper metalloenzyme that exploits biomass components. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 15079-84	11.5	692
92	Absence of branches from xylan in <i>Arabidopsis</i> gux mutants reveals potential for simplification of lignocellulosic biomass. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 17409-14	11.5	221
91	Phylogenetic and biochemical evidence supports the recruitment of an ADP-glucose translocator for the export of photosynthate during plastid endosymbiosis. <i>Molecular Biology and Evolution</i> , 2010 , 27, 2691-701	8.3	39
90	VASCULAR-RELATED NAC-DOMAIN6 and VASCULAR-RELATED NAC-DOMAIN7 effectively induce transdifferentiation into xylem vessel elements under control of an induction system. <i>Plant Physiology</i> , 2010 , 153, 906-14	6.6	189
89	Proteomic complex detection using sedimentation (ProCoDeS): screening for proteins in stable complexes and their candidate interaction partners. <i>Biochemical Society Transactions</i> , 2010 , 38, 923-7	5.1	2

88	The environmental and economic sustainability of potential bioethanol from willow in the UK. <i>Bioresource Technology</i> , 2010 , 101, 9612-23	11	55
87	Carbohydrate structural analysis of wheat flour arabinogalactan protein. <i>Carbohydrate Research</i> , 2010 , 345, 2648-56	2.9	84
86	Chemical and in situ characterization of macromolecular components of the cell walls from the green seaweed <i>Codium fragile</i> . <i>Glycobiology</i> , 2009 , 19, 212-28	5.8	74
85	Characterization of IRX10 and IRX10-like reveals an essential role in glucuronoxylan biosynthesis in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2009 , 57, 732-46	6.9	229
84	Cell wall glucomannan in <i>Arabidopsis</i> is synthesised by CSLA glycosyltransferases, and influences the progression of embryogenesis. <i>Plant Journal</i> , 2009 , 60, 527-38	6.9	132
83	ECA3, a Golgi-localized P2A-type ATPase, plays a crucial role in manganese nutrition in <i>Arabidopsis</i> . <i>Plant Physiology</i> , 2008 , 146, 116-28	6.6	128
82	Sub-cellular localization of membrane proteins. <i>Proteomics</i> , 2008 , 8, 3991-4011	4.8	61
81	Oligosaccharide relative quantitation using isotope tagging and normal-phase liquid chromatography/mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2008 , 22, 2723-30	2.2	43
80	Proteomic complex detection using sedimentation. <i>Analytical Chemistry</i> , 2007 , 79, 2078-83	7.8	27
79	Structure elucidation of arabinoxylan isomers by normal phase HPLC-MALDI-TOF/TOF-MS/MS. <i>Carbohydrate Research</i> , 2007 , 342, 724-35	2.9	85
78	Comparison of five xylan synthesis mutants reveals new insight into the mechanisms of xylan synthesis. <i>Plant Journal</i> , 2007 , 52, 1154-68	6.9	271
77	Plant organelle proteomics. <i>Current Opinion in Plant Biology</i> , 2007 , 10, 594-9	9.9	48
76	A multifunctional hybrid glycosyl hydrolase discovered in an uncultured microbial consortium from ruminant gut. <i>Applied Microbiology and Biotechnology</i> , 2007 , 74, 113-24	5.7	64
75	UDP-glucose 4-epimerase isoforms UGE2 and UGE4 cooperate in providing UDP-galactose for cell wall biosynthesis and growth of <i>Arabidopsis thaliana</i> . <i>Plant Cell</i> , 2007 , 19, 1565-79	11.6	98
74	The ectopically parting cells 1-2 (<i>epc1-2</i>) mutant exhibits an exaggerated response to abscisic acid. <i>Journal of Experimental Botany</i> , 2007 , 58, 1813-23	7	14
73	A novel bioinformatics approach identifies candidate genes for the synthesis and feruloylation of arabinoxylan. <i>Plant Physiology</i> , 2007 , 144, 43-53	6.6	154
72	Dynamic response of prevacuolar compartments to brefeldin a in plant cells. <i>Plant Physiology</i> , 2006 , 142, 1442-59	6.6	57
71	Galactoglucomannans increase cell population density and alter the protoxylem/metaxylem tracheary element ratio in xylogenic cultures of <i>Zinnia</i> . <i>Plant Physiology</i> , 2006 , 142, 696-709	6.6	39

70	Methods of quantitative proteomics and their application to plant organelle characterization. <i>Journal of Experimental Botany</i> , 2006 , 57, 1493-9	7	70
69	Resolution of the structural isomers of partially methylesterified oligogalacturonides by polysaccharide analysis using carbohydrate gel electrophoresis. <i>Glycobiology</i> , 2006 , 16, 29-35	5.8	213
68	Mapping the Arabidopsis organelle proteome. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006 , 103, 6518-23	11.5	464
67	Plant endoplasmic reticulum supports the protein secretory pathway and has a role in proliferating tissues. <i>Plant Journal</i> , 2006 , 48, 657-73	6.9	43
66	Quantitative proteomic approach to study subcellular localization of membrane proteins. <i>Nature Protocols</i> , 2006 , 1, 1778-89	18.8	70
65	Enzymatic fingerprinting of Arabidopsis pectic polysaccharides using polysaccharide analysis by carbohydrate gel electrophoresis (PACE). <i>Planta</i> , 2006 , 224, 163-74	4.7	26
64	Deficiency of adenosine kinase activity affects the degree of pectin methyl-esterification in cell walls of Arabidopsis thaliana. <i>Planta</i> , 2006 , 224, 1401-14	4.7	17
63	A unique family of proteins associated with internalized membranes in protein storage vacuoles of the Brassicaceae. <i>Plant Journal</i> , 2005 , 41, 429-41	6.9	34
62	An investigation of pectin methylesterification patterns by two independent methods: capillary electrophoresis and polysaccharide analysis using carbohydrate gel electrophoresis. <i>Carbohydrate Research</i> , 2005 , 340, 1193-9	2.9	22
61	Diversity of the exoproteome of Fusarium graminearum grown on plant cell wall. <i>Current Genetics</i> , 2005 , 48, 366-79	2.9	120
60	COBRA, an Arabidopsis extracellular glycosyl-phosphatidyl inositol-anchored protein, specifically controls highly anisotropic expansion through its involvement in cellulose microfibril orientation. <i>Plant Cell</i> , 2005 , 17, 1749-63	11.6	260
59	Analysis of detergent-resistant membranes in Arabidopsis. Evidence for plasma membrane lipid rafts. <i>Plant Physiology</i> , 2005 , 137, 104-16	6.6	395
58	SETH1 and SETH2, two components of the glycosylphosphatidylinositol anchor biosynthetic pathway, are required for pollen germination and tube growth in Arabidopsis. <i>Plant Cell</i> , 2004 , 16, 229-40	11.6	150
57	Localization of organelle proteins by isotope tagging (LOPIT). <i>Molecular and Cellular Proteomics</i> , 2004 , 3, 1128-34	7.6	255
56	Unusual microbial xylanases from insect guts. <i>Applied and Environmental Microbiology</i> , 2004 , 70, 3609-17	4.8	130
55	An evolutionary route to xylanase process fitness. <i>Protein Science</i> , 2004 , 13, 494-503	6.3	103
54	Arabidopsis thaliana expresses multiple Golgi-localised nucleotide-sugar transporters related to GONST1. <i>Molecular Genetics and Genomics</i> , 2004 , 272, 397-410	3.1	60
53	The use of isotope-coded affinity tags (ICAT) to study organelle proteomes in Arabidopsis thaliana. <i>Biochemical Society Transactions</i> , 2004 , 32, 520-3	5.1	54

52	The modular architecture of Cellvibrio japonicus mannanases in glycoside hydrolase families 5 and 26 points to differences in their role in mannan degradation. <i>Biochemical Journal</i> , 2003 , 371, 1027-43	3.8	95
51	Localisation and characterisation of cell wall mannan polysaccharides in Arabidopsis thaliana. <i>Planta</i> , 2003 , 218, 27-36	4.7	107
50	Analysis of methylated and unmethylated polygalacturonic acid structure by polysaccharide analysis using carbohydrate gel electrophoresis. <i>Analytical Biochemistry</i> , 2003 , 321, 174-82	3.1	27
49	Identification of glycosylphosphatidylinositol-anchored proteins in Arabidopsis. A proteomic and genomic analysis. <i>Plant Physiology</i> , 2003 , 132, 568-77	6.6	322
48	Glycosylphosphatidylinositol lipid anchoring of plant proteins. Sensitive prediction from sequence- and genome-wide studies for Arabidopsis and rice. <i>Plant Physiology</i> , 2003 , 133, 1691-701	6.6	168
47	The Arabidopsis ppi1 mutant is specifically defective in the expression, chloroplast import, and accumulation of photosynthetic proteins. <i>Plant Cell</i> , 2003 , 15, 1859-71	11.6	136
46	AtCSLA7, a cellulose synthase-like putative glycosyltransferase, is important for pollen tube growth and embryogenesis in Arabidopsis. <i>Plant Physiology</i> , 2003 , 131, 547-57	6.6	98
45	Polysaccharide analysis using carbohydrate gel electrophoresis: a method to study plant cell wall polysaccharides and polysaccharide hydrolases. <i>Analytical Biochemistry</i> , 2002 , 300, 53-68	3.1	144
44	Two-dimensional gel electrophoresis: recent advances in sample preparation, detection and quantitation. <i>Current Opinion in Chemical Biology</i> , 2002 , 6, 46-50	9.7	223
43	Prediction of glycosylphosphatidylinositol-anchored proteins in Arabidopsis. A genomic analysis. <i>Plant Physiology</i> , 2002 , 129, 486-99	6.6	164
42	GARNet, the Genomic Arabidopsis Resource Network. <i>Trends in Plant Science</i> , 2002 , 7, 145-7	13.1	10
41	Identification and Characterization of GONST1, a Golgi-Localized GDP-Mannose Transporter in Arabidopsis. <i>Plant Cell</i> , 2001 , 13, 2283	11.6	
40	Quantitative and reproducible two-dimensional gel analysis using Phoretix 2D Full. <i>Electrophoresis</i> , 2001 , 22, 2075-85	3.6	91
39	Identification and characterization of GONST1, a golgi-localized GDP-mannose transporter in Arabidopsis. <i>Plant Cell</i> , 2001 , 13, 2283-95	11.6	133
38	A proteomic analysis of organelles from Arabidopsis thaliana. <i>Electrophoresis</i> , 2000 , 21, 3488-99	3.6	117
37	Glycosylphosphatidylinositol-anchored cell-surface proteins from Arabidopsis. <i>Electrophoresis</i> , 1999 , 20, 2027-35	3.6	125
36	Use of a proteome strategy for tagging proteins present at the plasma membrane. <i>Plant Journal</i> , 1998 , 16, 633-41	6.9	125
35	The plant Golgi apparatus. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1998 , 1404, 259-70	4.9	95

34	Targeting of active sialyltransferase to the plant Golgi apparatus. <i>Plant Cell</i> , 1998 , 10, 1759-68	11.6	172
33	Plant embryogenesis: cell division forms a pattern. <i>Current Biology</i> , 1996 , 6, 683-5	6.3	9
32	Guilty by insolubility - does a protein's detergent insolubility reflect a caveolar location?. <i>Trends in Cell Biology</i> , 1995 , 5, 187-189	18.3	
31	Isolation of a murine cDNA clone encoding Rab19, a novel tissue-specific small GTPase. <i>Gene</i> , 1995 , 155, 257-60	3.8	20
30	Co-operative regulation of endocytosis by three Rab5 isoforms. <i>FEBS Letters</i> , 1995 , 366, 65-71	3.8	132
29	A deficiency of the small GTPase rab8 inhibits membrane traffic in developing neurons. <i>Molecular and Cellular Biology</i> , 1995 , 15, 918-24	4.8	85
28	Use of antisense oligonucleotides to study Rab function in vivo. <i>Methods in Enzymology</i> , 1995 , 257, 302-127		4
27	Guilty by insolubility--does a protein's detergent insolubility reflect a caveolar location?. <i>Trends in Cell Biology</i> , 1995 , 5, 187-9	18.3	82
26	VIP21-Caveolin, a protein of the trans-Golgi network and caveolae. <i>FEBS Letters</i> , 1994 , 346, 88-91	3.8	78
25	Isolation of a mouse cDNA encoding Rab23, a small novel GTPase expressed predominantly in the brain. <i>Gene</i> , 1994 , 138, 207-11	3.8	39
24	Expression of exogenous proteins in mammalian cells with the Semliki Forest virus vector. <i>Methods in Cell Biology</i> , 1994 , 43 Pt A, 43-53	1.8	39
23	Mapping of Ras-related GTP-binding proteins by GTP overlay following two-dimensional gel electrophoresis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994 , 91, 7874-8	11.5	53
22	Cloning and subcellular localization of novel rab proteins reveals polarized and cell type-specific expression. <i>Journal of Cell Science</i> , 1994 , 107, 3437-3448	5.3	77
21	Protein Sorting and Glycolipid-Enriched Detergent-Insoluble Complexes in Epithelial Cells 1994 , 107-114		0
20	VIP21, Caveolae and Sorting in the Trans-Golgi Network of Epithelial Cells 1994 , 45-51		
19	Protein transport to the dendritic plasma membrane of cultured neurons is regulated by rab8p. <i>Journal of Cell Biology</i> , 1993 , 123, 47-55	7.3	113
18	Caveolae and sorting in the trans-Golgi network of epithelial cells.. <i>EMBO Journal</i> , 1993 , 12, 1597-1605	13	344
17	Molecular cloning and subcellular localization of three GTP-binding proteins of the rab subfamily. <i>Journal of Cell Science</i> , 1993 , 106 (Pt 4), 1249-61	5.3	42

16	Molecular cloning and subcellular localization of three GTP-binding proteins of the rab subfamily. <i>Journal of Cell Science</i> , 1993 , 106, 1249-1261	5.3	108
15	Caveolae and sorting in the trans-Golgi network of epithelial cells. <i>EMBO Journal</i> , 1993 , 12, 1597-605	13	142
14	Compartmentalization of rab Proteins in Mammalian Cells. <i>Handbook of Experimental Pharmacology</i> , 1993 , 423-445	3.2	1
13	VIP21, a 21-kD membrane protein is an integral component of trans-Golgi-network-derived transport vesicles. <i>Journal of Cell Biology</i> , 1992 , 118, 1003-14	7.3	49 ⁸
12	Sequence of a canine cDNA clone encoding a Ran/TC4 GTP-binding protein. <i>Gene</i> , 1992 , 120, 325-6	3.8	11
11	Internal fragmentation of proteins in polyacrylamide matrices for microsequencing. <i>The Protein Journal</i> , 1992 , 11, 356-356		1
10	Biogenesis of cell-surface polarity in epithelial cells and neurons. <i>Cold Spring Harbor Symposia on Quantitative Biology</i> , 1992 , 57, 611-9	3.9	44
9	Regulation of Expression of the Pea Plastocyanin Gene 1992 , 23-29		
8	Expression of photosynthesis gene-promoter fusions in leaf epidermal cells of transgenic tobacco plants. <i>Plant Journal</i> , 1991 , 1, 115-120	6.9	32
7	A scaffold-associated DNA region is located downstream of the pea plastocyanin gene. <i>Plant Cell</i> , 1991 , 3, 1239-50	11.6	50
6	A Scaffold-Associated DNA Region Is Located Downstream of the Pea Plastocyanin Gene. <i>Plant Cell</i> , 1991 , 3, 1239	11.6	3
5	Expression of a Pea Gene for Ferredoxin-NADP+ Oxidoreductase in Transgenic Tobacco Plants 1990 , 2531-2534		
4	GONST2 transports GDP-Mannose for sphingolipid glycosylation in the Golgi apparatus of Arabidopsis		1
3	Scission of Glucosidic Bonds by a <i>Lentinus similis</i> Lytic Polysaccharide Monooxygenases Is Strictly Dependent on H ₂ O ₂ while the Oxidation of Saccharide Products Depends on O ₂ . <i>ACS Catalysis</i> , 13848-13859	13.1	2
2	Expression of a glucomannan mannosyltransferase gene (GMMT) from Aloe vera is induced by water deficit and abscisic acid		1
1	Glycosyltransferases of the GT43 Family 251-263		