
Cemal Can Bilgin

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2052313/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Species level classification of Mediterranean sparse forests-maquis formations using Sentinel-2 imagery. Geocarto International, 2022, 37, 1587-1606. | 3.5 | 4 |
| 2 | Connecting the data landscape of longâ€ŧerm ecological studies: The SPIâ€Birds data hub. Journal of Animal Ecology, 2021, 90, 2147-2160. | 2.8 | 25 |
| 3 | Unsustainable harvest of water frogs in southern Turkey for the European market. Oryx, 2021, 55, 364-372. | 1.0 | 11 |
| 4 | Variable kinship patterns in Neolithic Anatolia revealed by ancient genomes. Current Biology, 2021, 31, 2455-2468.e18. | 3.9 | 47 |
| 5 | Song overlapping, noise, and territorial aggression in great tits. Behavioral Ecology, 2020, 31, 807-814. | 2.2 | 15 |
| 6 | Genetic diversity is retained in a bottlenecked Cinereous Vulture population in Turkey. Ibis, 2019, 161, 793-805. | 1.9 | 4 |
| 7 | Extremely Low Mitochondrial DNA Diversity in a near Threatened Species, the Cinereous Vulture Aegypius monachus in Turkey. Ardea, 2019, 107, 85. | 0.6 | 2 |
| 8 | Nest site selection patterns of a local Egyptian Vulture <i>Neophron percnopterus</i> population in Turkey. Bird Conservation International, 2017, 27, 568-581. | 1.3 | 15 |
| 9 | Comparison of three different primer sets for sexing birds. Journal of Veterinary Diagnostic Investigation, 2017, 29, 59-63. | 1.1 | 59 |
| 10 | Strong population structure in a species manipulated by humans since the Neolithic: the European fallow deer (Dama dama dama). Heredity, 2017, 119, 16-26. | 2.6 | 30 |
| 11 | Archaeogenomic analysis of the first steps of Neolithization in Anatolia and the Aegean. Proceedings of the Royal Society B: Biological Sciences, 2017, 284, 20172064. | 2.6 | 26 |
| 12 | Exceptional maternal lineage diversity in brown bears (<i>Ursus arctos</i>) from Turkey. Zoological Journal of the Linnean Society, 2016, 176, 463-477. | 2.3 | 44 |
| 13 | An overview of biodiversity and conservation status of steppes of the Anatolian Biogeographical Region. Biodiversity and Conservation, 2016, 25, 2491-2519. | 2.6 | 30 |
| 14 | IMPACT OF LARGE SCALE DAM CONSTRUCTION ON MOVEMENT CORRIDORS OF MAMMALS IN ARTVIN, NORTH-EASTERN TURKEY. Applied Ecology and Environmental Research, 2016, 14, 489-507. | 0.5 | 4 |
| 15 | Genetic evidence for human-mediated introduction of Anatolian water frogs (<i>Pelophylax</i> cf. <i>bedriagae</i>) to Cyprus (Amphibia: Ranidae). Zoology in the Middle East, 2015, 61, 125-132. | 0.6 | 12 |
| 16 | Effects of landscape, land use and vegetation on bird community composition and diversity in Inner Anatolian steppes. Agriculture, Ecosystems and Environment, 2014, 182, 37-46. | 5.3 | 21 |
| 17 | First record of a melanistic golden jackal (Canis aureus, Canidae) from Turkey. Mammalia, 2013, 77, . | 0.7 | 7 |
| 18 | Post-Fledging Movements of Cinereous Vultures <i>Aegypius monachus</i> in Turkey Revealed by GPS Telemetry, Ardea, 2012, 100, 149-156. | 0.6 | 28 |

CEMAL CAN BILGIN

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Genetic data reveal that water frogs of Cyprus (genus <i>Pelophylax</i>) are an endemic species of Messinian origin. Zoosystematics and Evolution, 2012, 88, 261-283. | 1.1 | 37 |
| 20 | Identification of Prime Butterfly Areas in Turkey using systematic conservation planning: Challenges and opportunities. Biological Conservation, 2012, 150, 86-93. | 4.1 | 9 |
| 21 | Phylogeographic patterns of genetic diversity in eastern Mediterranean water frogs were determined by geological processes and climate change in the Late Cenozoic. Journal of Biogeography, 2010, 37, 2111-2124. | 3.0 | 101 |
| 22 | Discordance between ventral colour and mtDNA haplotype in the water frog Rana (ridibunda) caralitana, 1988 Arıkan. Amphibia - Reptilia, 2010, 31, 9-20. | 0.5 | 15 |
| 23 | Genetic Divergence and Evolution of Reproductive Isolation in Eastern Mediterranean Water Frogs. , 2010, , 373-403. | | 35 |
| 24 | Human–Brown Bear Conflicts in Artvin, Northeastern Turkey: Encounters, Damage, and Attitudes. Ursus, 2008, 19, 146-153. | 0.5 | 67 |
| 25 | Intergration of environmental variables with satellite images in regional scale vegetation classification. International Journal of Remote Sensing, 2007, 28, 2397-2397. | 2.9 | 1 |
| 26 | Stopover Ecology of Some Passerines at Ankara (Central Turkey). Ring, 2005, 27, 127-136. | 0.4 | 4 |
| 27 | A numerical taxonomic study of the genus Acantholimon Boiss. (Plumbaginaceae) in Ankara Province (Turkey). Israel Journal of Plant Sciences, 2001, 49, 298-300. | 0.5 | 8 |
| 28 | First record of the Common Myna <i>(Acridotheres tristis)</i> from Ankara, Turkey. Zoology in the Middle East, 1996, 13, 25-26. | 0.6 | 2 |
| 29 | First record of <i>Musca domestica calleva</i> Walker, 1849 (Diptera: Muscidae) from Southern Turkey. Zoology in the Middle East, 1995, 11, 81-85. | 0.6 | 0 |
| 30 | The Purple Gallinule, <i>Porphyrio porphyrio</i> , in the Eastern Mediterranean. Zoology in the Middle East, 1989, 3, 19-30. | 0.6 | 2 |